



POSTER PRESENTATION

Open Access

Altered innate functions of myeloid dendritic cells in ANCA-associated vasculitis

Cécile Braudeau^{1,2}, Antoine Néel^{2,3}, Marie Rimbart^{1,2}, Mohamed Hamidou^{2,4}, Régis Josien^{1,2,4*}

From 7th European Workshop on Immune-Mediated Inflammatory Diseases Noordwijk aan Zee, the Netherlands. 28-30 November 2012

Background

Dendritic cells (DC) are critical effectors of innate and adaptive immunity, acting both as sentinels that detect the presence of pathogens and as key antigen-presenting cells that regulate the adaptive immune response. Therefore, DC play a crucial role in the control of autoimmune responses. We previously showed that blood DC numbers were strongly reduced in ANCA-associated vasculitis (AAV) likely due to their recruitment in tissues. Here, we assessed the ex vivo responsiveness of blood DC from AAV-patients to Toll-like receptors (TLRs) stimulation.

Materials and methods

Blood samples from 10 untreated patients with AAV during flares and before any immunosuppressive treatment (AP) were analyzed, along with 9 AAV patients in remission (RP) and 11 age-matched healthy controls (HC). Intracellular cytokine (IL-12, TNF- α , IFN- α) production by blood DC was assessed by 8-colors flow cytometry after stimulation by Toll-like receptors of whole blood samples.

Results

We found that myeloid DC (mDC) from patients in acute phase exhibited a decreased IL-12 production after TLR3, 4 and 7/8 stimulation compared to patients in remission and healthy controls. These mDC also produced less TNF- α after TLR3 stimulation. Moreover, we observed a reduction in the frequency TNF α -producing plasmacytoid DC (pDC) upon TLR7/8 triggering in AP patients compared to RP patients and HC.

Conclusion

Our data show that circulating mDC from patients with AAV exhibited an altered response to several TLR ligands, with a notable decrease in IL-12 production.

These unexpected results suggest the innate functions of DC especially in response to pathogens are impaired during AAV.

Author details

¹CHU Nantes, Laboratoire d'Immunologie, Nantes, France. ²INSERM Center of Research in Transplantation and Immunology, UMR 1064, Nantes, France. ³CHU Nantes, Service de Médecine Interne, Nantes, France. ⁴Université de Nantes, Faculté de Médecine, Nantes, France.

Published: 28 November 2012

doi:10.1186/1479-5876-10-S3-P20

Cite this article as: Braudeau et al.: Altered innate functions of myeloid dendritic cells in ANCA-associated vasculitis. *Journal of Translational Medicine* 2012 **10**(Suppl 3):P20.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹CHU Nantes, Laboratoire d'Immunologie, Nantes, France
Full list of author information is available at the end of the article