



HAL
open science

Inducible Treg cell populations as cell based-therapy for rheumatoid arthritis

Delphine Martire, Julie Quentin, Anne-Laure Mausset-Bonnefont, H el ene Asnagli, Nathalie Belmonte, Arnaud Foussat, Christian Jorgensen, Pascale Louis-Plence

► **To cite this version:**

Delphine Martire, Julie Quentin, Anne-Laure Mausset-Bonnefont, H el ene Asnagli, Nathalie Belmonte, et al.. Inducible Treg cell populations as cell based-therapy for rheumatoid arthritis. 7th European Workshop on Immune-Mediated Inflammatory Diseases, Nov 2012, Noordwijk aan Zee, Netherlands. pp.P55. inserm-00758221

HAL Id: inserm-00758221

<https://www.hal.inserm.fr/inserm-00758221>

Submitted on 28 Nov 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.



POSTER PRESENTATION

Open Access

Inducible Treg cell populations as cell based-therapy for rheumatoid arthritis

Delphine Martire¹, Julie Quentin¹, Anne-Laure Mausset-Bonnefont¹, H el ene Asnagli², Nathalie Belmonte², Arnaud Foussat², Christian Jorgensen¹, Pascale Louis-Plence^{1*}

From 7th European Workshop on Immune-Mediated Inflammatory Diseases Noordwijk aan Zee, the Netherlands. 28-30 November 2012

Background

Adoptive cell transfer of Treg cells is a promising approach to restore tolerance in autoimmune disease.

However the various type of Tregs, their doses of injection and their *in vivo*-suppressive mechanism need to be precisely define to clearly establish which Tregs will be able to dampen efficiently the immune response in the various settings.

In our study, we compared the therapeutic potential of induced CD25⁺FoxP3⁺ and two IL10-secreting Tregs: Tr1 and CD49b-induced Tregs.

Materials and methods

CD49b Treg cells were generated in naive mice following repetitive injections of iDC. The purification was based on the negative selection of CD4 T cells isolated from the spleen and liver of the iDC-vaccinated mice.

Cell sorting experiments were realized to obtain 98% pure CD49b⁺ T or CD25⁺ cells. Collagen type II (bCII) specific Tr1 clones were obtained from TCR transgenic mice and expanded *in vitro*. Selected clones showed *in vitro* antigen specificity, Tr1 cytokine profile and IL10- and TGF -dependent suppressive activity.

Results

Several doses of CD49b or Tr1 cells were injected i.v. at day 28 in established collagen-induced arthritis. Clinical signs of arthritis were scored, as well as biological parameters such as the level of anti-bCII antibodies in sera and the cytokine profile of bCII specific T cells.

We defined for both Treg cell populations the dose effect in curative settings experiments. One single dose of 3x10⁶ or 1x10⁶ of Tr1 cell administration could reduce

the incidence and severity of CIA. Interestingly, higher dose of 10M of Tr1 cells did not improve the disease. In the same manner, the dose of 10⁵ CD4CD49b⁺ or CD25⁺ cells reverse clinical symptom with a lack of efficacy of higher doses. The cytokinic profile of the Tr1 cells was also investigated in inflammatory settings as well as the impact of the various Treg cells on the proliferation of effector cells *in vivo*.

Conclusions

Our results suggest that even if the Treg cells present some similarities, we need to precisely define the dose and type of Treg that will be efficient in each experimental setting.

Author details

¹Inserm U844, Universit  Montpellier 1, CHU Lapeyronnie, Montpellier, France. ²TxCell, Valbonne-Sophia Antipolis, France.

Published: 28 November 2012

doi:10.1186/1479-5876-10-S3-P55

Cite this article as: Martire et al.: Inducible Treg cell populations as cell based-therapy for rheumatoid arthritis. *Journal of Translational Medicine* 2012 **10**(Suppl 3):P55.

¹Inserm U844, Universit  Montpellier 1, CHU Lapeyronnie, Montpellier, France
Full list of author information is available at the end of the article