



HAL
open science

Two large-scale analyses of Ty1 LTRretrotransposon de novo insertion events indicate that Ty1 targets nucleosomal DNA near the H2A/H2B interface.

Antoine Bridier-Nahmias, Pascale Lesage

► To cite this version:

Antoine Bridier-Nahmias, Pascale Lesage. Two large-scale analyses of Ty1 LTRretrotransposon de novo insertion events indicate that Ty1 targets nucleosomal DNA near the H2A/H2B interface.. Mobile DNA, 2012, 3 (1), pp.22. 10.1186/1759-8753-3-22 . inserm-00776299

HAL Id: inserm-00776299

<https://inserm.hal.science/inserm-00776299>

Submitted on 15 Jan 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

COMMENTARY

Open Access

Two large-scale analyses of Ty1 LTR-retrotransposon *de novo* insertion events indicate that Ty1 targets nucleosomal DNA near the H2A/H2B interface

Antoine Bridier-Nahmias^{1,2,3*} and Pascale Lesage^{1,2,3}

Abstract

Background: Over the years, a number of reports have revealed that Ty1 integration occurs in a 1-kb window upstream of Pol III-transcribed genes with an approximate 80-bp periodicity between each integration hotspot and that this targeting requires active Pol III transcription at the site of integration. However, the molecular bases of Ty1 targeting are still not understood.

Findings: The publications by Baller et al. and Mularoni et al. in the April issue of *Genome Res.* report the first high-throughput sequencing analysis of Ty1 *de novo* insertion events. Their observations converge to the same conclusion, that Ty1 targets a specific surface of the nucleosome at the H2A/H2B interface.

Conclusion: This discovery is important, and should help identifying factor(s) involved in Ty1 targeting. Recent data on transposable elements and retroviruses integration site choice obtained by large-scale analyses indicate that transcription and chromatin structure play an important role in this process. The studies reported in this commentary add a new evidence of the importance of chromatin in integration selectivity that should be of interest for everyone interested in transposable elements integration.

Keywords: LTR retrotransposon, Ty1, Selective integration, Large-scale analysis

Findings

Retrotransposons are major components of eukaryotic genomes. They represent, for example, half of the human genome, up to 80% of some plant's genomes and 3% of the compact genome of Yeast *S. cerevisiae*. They have a central role in shaping genomes, and have been shown to be a powerful force of evolution and to play a positive role in long-term adaptation. However, they can also be deleterious in the short-term, since their integration into the host genome can inactivate or deregulate gene expression or even induce large chromosomal rearrangements by homologous recombination of distant copies. LTR-

retrotransposons are structurally and functionally related to retroviruses but their life cycle is exclusively intracellular since they do not encode an envelope glycoprotein. They replicate by reverse transcribing their RNA into cDNA, which is ultimately integrated into the host genome by the element-encoded integrase (IN). The non-random distribution of LTR-retrotransposons and retroviruses into genomes suggests that these elements actively select their integration sites (for review, [1]).


For the past 20 years, studies on Ty3 and Ty5 LTR-retrotransposons of *S. cerevisiae* have led to better understand the molecular bases of their targeted integration (Figure 1). It has been established that an interaction between Ty3 IN and the Brf1 subunit of TFIIB is sufficient to target Ty3 integration to Pol III transcription initiation sites *in vitro* [2]. Likewise, *in vivo*, Ty5 preferential integration into silent telomeric heterochromatin depends on the interaction between Ty5 IN and the Sir4 protein, a

* Correspondence: antoine.bridier@inserm.fr

¹CNRS/P7 UMR7212, INSERMU 944, Laboratoire Pathologie et Virologie Moléculaire Institut Universitaire d'Hématologie, 1 avenue Claude Vellefaux, F75010, Paris, France

²Inserm U944, Institut Universitaire d'Hématologie, Hôpital St Louis, Paris, France

Full list of author information is available at the end of the article




structural component of silent chromatin [3]. In the distant Yeast *S. pombe*, interaction between Tf1 IN and the Atf1 transcription factor plays a direct and specific role in targeting Tf1 integration in the *fbp1* gene promoter [4]. These studies have converged on a common targeting mechanism, based on tethering of integration complexes to the cell genome through interaction between IN and cellular proteins bound at favored insertion sites. It is noteworthy that this model may also account for the selectivity of the integration of retroviruses, since HIV-1 integration in active transcription units relies on the interaction of its IN with the LEDGF/p75 transcription factor (reviewed in [5]).

Discovered in 1979 [6], Ty1 is the most abundant and active LTR-retrotransposon in *S. cerevisiae*. It was first noticed in 1982 that Ty1 insertions are located adjacent to several tRNA genes [7]. Over the years, a number of reports have found that Ty1 integration occurs in a 1-kb window upstream of Pol III-transcribed genes with an approximate 80-bp periodicity between each integration hotspot and that this targeting requires active Pol III transcription at the site of integration (Figure 1) [8-10]. However, despite 30 years of active research, the molecular bases of Ty1 targeting are still not understood. Thus, the recent articles of Baller *et al.* and Mularoni *et al.* are an important advance for understanding the selection of Ty1 integration sites, by showing that Ty1 targets a specific surface of the nucleosome [11,12].

Both studies used a deep-sequencing approach to get insights into Ty1 integration selectivity. To discriminate *de novo* insertion events from resident Ty1 elements, both used a short tag sequence introduced in one LTR of a galactose-inducible donor Ty1 element, such that after a complete retrotransposition cycle the tag would be recovered in both LTRs (Figure 2), and this short tag was used to specifically sequence newly transposed sequences. In the Mularoni *et al.*'s study, sequences were generated by an Illumina GAI1 apparatus. Of a total of 7,990,112 reads, 1,154,281 were characterized as non-redundant Ty1 insertions. Baller *et al.* used a donor Ty1 element, which contained the *his3AI* reporter construct conferring histidine prototrophy to the cells after Ty1 retrotransposition, and recovered only His⁺ insertions. A single 454 run produced from 13,000 to 111,000 reads. While Mularoni *et al.* analyzed 10- to 100-fold more integration events than Baller *et al.*, the latter analyzed integration profiles in diploid and haploid wild-type cells and in a panel of mutants affecting DNA-related processes known to increase integration frequency (*rrm3Δ*, *rtt109Δ*, *hos2Δ*) and integration in coding sequences and transcription units (*rad6Δ*). *RTT109* and *HOS2* encode histone-modifying enzymes, while *RRM3* encodes a helicase and *RAD6* an ubiquitin-conjugating enzyme.

Both reports describe the same general insertion profile pattern in wild-type cells. Independently of cell ploidy, a



vast majority (~90%) of insertions were observed as predicted in the 5' region of class III genes. However, Ty1 did not target class III genes equally. All but two tRNA genes, tE(UUC)C and tI(AAU)L1, received many insertions. Other class III genes, such as *SNR6*, *RPRI*, *SNR52*, *SCR1* and the repeated locus *RDN5* received insertions as well. However, two class III loci of unknown function, *RNA170* and *ZOD1*, were not targeted, probably because the Pol III transcription machinery was not efficiently recruited at these loci. Insertions in Pol II-transcribed genes were rare, representing about 5% of total events. Most of them occurred near a Pol III gene, with a strong preference for the region closest to the class III gene. Considering ORFs which are more distant to tRNA genes (~5 kb), the few recovered insertions occurred at the gene 5' end. Although insertions into mitochondrial sequences (mtDNA) were reported by Mularoni *et al.*, they were not detected by Baller *et al.*, probably because of a smaller dataset size or because those insertions did not confer histidine prototrophy and were, consequently, not selected for sequencing. Mularoni *et al.* suggest that these insertions might come from shattered mitochondria and could occur in the nucleus or even in the cytoplasm.


The novel and most striking observation of both studies is that Ty1 integration is positively correlated with nucleosome occupancy. An important role for chromatin in the selection process of insertion sites was already suspected after the discovery of an intriguing periodicity of ~80 bp between each integration hotspots that relied on the ATP-dependent chromatin remodeling factor Isw2 [10]. By comparing their deep-sequencing results with genome-wide nucleosome positioning data sets, they have discovered two hotspots per nucleosome, separated by about 70 bp. These observations were made for the first three nucleosomes directly upstream of a class III gene

and the integration events were aligned with the nucleosome H2A/H2B interface (Figure 3).

In *hos2Δ* and *rtt109Δ* mutant strains analyzed by Baller *et al.*, the pattern of Ty1 insertion events was not significantly different from that in wild-type cells. In contrast, integration events in verified ORFs increased significantly in *rrm3Δ* and *rad6Δ* mutant strains (by two- and three-fold, respectively), although the integration pattern upstream of tRNA genes was unmodified, leading to the conclusion that the determining factors for specific nucleosomal targeting upstream of class III genes were not affected in these mutants.

High-throughput sequencing of insertion events has provided a saturated profile of target activity for several retrotransposons and contributed to better understand their integration preferences. For example, integration of the LTR retrotransposon of *S. pombe* Tf1 has been shown to be strongly biased for Pol II promoters with a clear preference for stress-induced promoters [13], and another report, in which 10,000 events have been analyzed, confirms all previous *in vitro* evidences on Ty3 integration at Pol III sites [14]. A high-throughput sequencing strategy has also been largely used to map the insertions profiles of different retroviruses (for review [15]). The reports of Mularoni *et al.* and Baller *et al.* reveal that Ty1 integration upstream of class III genes is strongly correlated with the chromatin structure at these loci and preferentially targets a specific nucleosomal DNA segment. Interestingly, the preference for nucleosome-rich regions is not a conserved feature of retroelements since it has been shown that elements such as Ty5 and Hermes (when expressed in yeast) prefer nucleosome-free regions [16,17].

Although these two studies clearly contribute to better understanding of Ty1 targeting, they do not characterize



whether a specific nucleosomal DNA conformation, a specific histone modification, or a nucleosome-bound factor enriched at sites of Pol III transcription, determine Ty1 preferred target sites, nor do they elucidate the role of RNA polymerase III and its co-factors in Ty1 targeting. Thus, further work is required to completely decipher the molecular bases of Ty1 targeting.

Abbreviations

H2A: Histone protein H2A; H2B: Histone protein H2B; His+: Histidine prototrophy; HIV-1: Human Immunodeficiency Virus 1; IN: Integrase; LTR: Long Terminal Repeat; ORF: Open Reading Frame; Pol III: RNA Polymerase III; Ty: Transposon in yeast.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

AB-N, wrote the manuscript PL, supervised the final manuscript. All authors read and approved the final manuscript.

Acknowledgements

We thank JC Gluckman for critical editing of the manuscript.

Funding

A B-N is supported by a graduate student fellowship from the CNRS (bourse des ingénieurs).

Work in the laboratory of PL is supported by the French National Research Agency against AIDS (ANRS) and intramural fundings from CNRS and INSERM.

Author details

¹CNRS/P7 UMR7212, INSERMU 944, Laboratoire Pathologie et Virologie Moléculaire Institut Universitaire d'Hématologie, 1 avenue Claude Vellefaux, F75010, Paris, France. ²Inserm U944, Institut Universitaire d'Hématologie, Hôpital St Louis, Paris, France. ³Université Paris Diderot, Sorbonne Paris Cité, Hôpital St Louis, Paris, France.

Received: 25 June 2012 Accepted: 29 August 2012

Published: 17 December 2012

References

1. Delelis O, Zamborlini A, Thierry S, Saib A: **Chromosomal tethering and proviral integration.** *Biochim Biophys Acta* 2010, **1799**:207–216.
2. Yieh L, Kassavetis G, Geiduschek EP, Sandmeyer SB: **The Brf and TATA-binding protein subunits of the RNA polymerase III transcription factor IIIB mediate position-specific integration of the gypsy-like element, Ty3.** *J Biol Chem* 2000, **275**:29800–29807.
3. Xie W, Gai X, Zhu Y, Zappulla DC, Sternglanz R, et al: **Targeting of the yeast Ty5 retrotransposon to silent chromatin is mediated by interactions between integrase and Sir4p.** *Mol Cell Biol* 2001, **21**:6606–6614.
4. Leem YE, Ripmaster TL, Kelly FD, Ebina H, Heincelman ME, et al: **Retrotransposon Tf1 is targeted to Pol II promoters by transcription activators.** *Mol Cell* 2008, **30**:98–107.
5. Engelman A, Cherepanov P: **The lentiviral integrase binding protein LEDGF/p75 and HIV-1 replication.** *PLoS Pathog* 2008, **4**:e1000046.
6. Cameron JR, Loh EY, Davis RW: **Evidence for transposition of dispersed repetitive DNA families in yeast.** *Cell* 1979, **16**:739–751.
7. Eigel A, Feldmann H: **Ty1 and delta elements occur adjacent to several tRNA genes in yeast.** *EMBO J* 1982, **1**:1245–1250.
8. Devine SE, Boeke JD: **Integration of the Yeast Retrotransposon Ty1 Is Targeted to Regions Upstream of Genes Transcribed by RNA-Polymerase-III.** *Genes Dev* 1996, **10**:620–633.
9. Bachman N, Eby Y, Boeke JD: **Local definition of Ty1 target preference by long terminal repeats and clustered tRNA genes.** *Genome Res* 2004, **14**:1232–1247.
10. Bachman N, Gelbart ME, Tsukiyama T, Boeke JD: **TFIIIB subunit Bdp1p is required for periodic integration of the Ty1 retrotransposon and targeting of lsw2p to *S. cerevisiae* tDNAs.** *Genes Dev* 2005, **19**:955–964.

11. Mularoni L, Zhou Y, Bowen T, Gangadharan S, Wheelan SJ, et al: **Retrotransposon Ty1 integration targets specifically positioned asymmetric nucleosomal DNA segments in tRNA hotspots.** *Genome Res* 2012, **22**:693–703.
12. Baller JA, Gao J, Stamenova R, Curcio MJ, Voytas DF: **A nucleosomal surface defines an integration hotspot for the *Saccharomyces cerevisiae* Ty1 retrotransposon.** *Genome Res* 2012, **22**:704–713.
13. Guo Y, Levin HL: **High-throughput sequencing of retrotransposon integration provides a saturated profile of target activity in *Schizosaccharomyces pombe*.** *Genome Res* 2010, **20**:239–248.
14. Qi X, Daily K, Nguyen K, Wang H, Mayhew D, et al: **Retrotransposon profiling of RNA polymerase III initiation sites.** *Genome Res* 2012, **22**:681–692.
15. Desfarges S, Ciuffi A: **Retroviral integration site selection.** *Viruses* 2010, **2**:111–130.
16. Baller JA, Gao J, Voytas DF: **Access to DNA establishes a secondary target site bias for the yeast retrotransposon Ty5.** *Proc Natl Acad Sci U S A* 2011, **108**:20351–20356.
17. Gangadharan S, Mularoni L, Fain-Thornton J, Wheelan SJ, Craig NL: **DNA transposon Hermes inserts into DNA in nucleosome-free regions in vivo.** *Proc Natl Acad Sci U S A* 2010, **107**:21966–21972.

doi:10.1186/1759-8753-3-22

Cite this article as: Bridier-Nahmias and Lesage: Two large-scale analyses of Ty1 LTR-retrotransposon *de novo* insertion events indicate that Ty1 targets nucleosomal DNA near the H2A/H2B interface. *Mobile DNA* 2012 **3**:22.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

