

Fasting in mood disorders: neurobiology and effectiveness. A review of the literature.

Guillaume Fond, Alexandra Macgregor, Marion Leboyer, Andreas Michalsen

► **To cite this version:**

Guillaume Fond, Alexandra Macgregor, Marion Leboyer, Andreas Michalsen. Fasting in mood disorders: neurobiology and effectiveness. A review of the literature.. *Psychiatry Research*, Elsevier, 2013, 209 ((3)), pp.253-8. 10.1016/j.psychres.2012.12.018 . inserm-00789122

HAL Id: inserm-00789122

<https://www.hal.inserm.fr/inserm-00789122>

Submitted on 15 Feb 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Study	Fasting protocol	Study design and population	Mood evaluation	Major findings
Teng 2011	Three months clinical trial on CR (reduction of 300 to 500 kcal/day) combined with two days/week of MF.	Randomized controlled trial Healthy individuals N=25	BDI- II QOL Pittsburgh Sleep Quality Index Perceived Stress Scale	significant energy increase in QOL questionnaire (P<0.05). No significant changes in depression and stress level, but baseline levels were very low. No change in sleep quality.
Farooq 2010	MF during Ramadan	Open-labelled bipolar affective disorder N=62	HDRS YMRS	significant decrease in HDRS and YMRS during Ramadan (F=34.12, P<0.01, for HDRS and F=15.6, P<0.001 for YMRS)
Michalsen 2009	8-day modified medical fasting treatment (total energy intake <350 kcal/d)	Controlled Healthy subjects N=108	Daily ratings of mood (VAS)	Significant mood improvement (p=0.036)
Tavakkoli 2008	MF during Ramadan	Controlled trial with irritable bowel syndrome N=60	HADS QOL	Decrease of anxiety score (from 12.7 (6.0) to 9.8 (4.4) after fasting (p=0.026)
Lepage 2008	CR based on patient's self-declaration	Observational study Female with CR vs non-CR N=486	CES-D BSI	Significant group differences on the BSI, F(3, 289) = 4.84, p < .005, while results for the CES-D neared significance, F(3, 309) = 3.03, p = .03.
Kanazawa 2006	Complete fasting during 10 days	Controlled trial Irritable Bowel Syndrome N=58	Clinical interview	Fasting significantly improved anorexia (p = .02) and anxiety (p < .001), and interference with life in general (p < .001).

Michalsen 2006	8-day modified fast (300 kcal/day)	Controlled trial Chronic Pain Syndromes N=36	Daily ratings of mood (VAS)	Mood ratings increased significantly in the late phase of fasting (P < 0.01) but were not related to weight-loss, leptin- depletion or cortisol increase.
Suzuki 1976	Complete fasting during 10 days	Observation study Depression N=36	Clinical interview	Reported remission rate of 86%

Table 1. Clinical studies evaluating mood in populations under fasting condition. BDI : Beck Depression Inventory. HDRS : Hamilton Depression Rating Scale . HADS : Hospitalisation Anxiety and Depression Scale. YMRS : Young Mania Rating Scale. CES-D : Center for Epidemiological Studies- Depression Scale. QOL : quality of life. CR : caloric restriction. MF : Muslim fasting. BSI Brief Symptom Inventory. VAS : Visual Analogue Scale.