

**Job strain as a risk factor for leisure-time physical inactivity: an individual-participant meta-analysis of up to 170,000 men and women: the IPD-Work Consortium.**

Eleonor Fransson, Katriina Heikkilä, Solja Nyberg, Marie Zins, Hugo Westerlund, Peter Westerholm, Ari Väänänen, Marianna Virtanen, Jussi Vahtera, Töres Theorell, et al.

► **To cite this version:**

Eleonor Fransson, Katriina Heikkilä, Solja Nyberg, Marie Zins, Hugo Westerlund, et al.. Job strain as a risk factor for leisure-time physical inactivity: an individual-participant meta-analysis of up to 170,000 men and women: the IPD-Work Consortium.. American Journal of Epidemiology, Oxford University Press (OUP), 2012, 176 (12), pp.1078-89. <10.1093/aje/kws336>. <inserm-00806190>

**HAL Id: inserm-00806190**

**<https://www.hal.inserm.fr/inserm-00806190>**

Submitted on 29 Mar 2013

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Systematic Reviews and Meta- and Pooled Analyses

### Job Strain as a Risk Factor for Leisure-Time Physical Inactivity: An Individual-Participant Meta-Analysis of Up to 170,000 Men and Women

The IPD-Work Consortium

Eleonor I. Fransson\*, Katriina Heikkilä, Solja T. Nyberg, Marie Zins, Hugo Westerlund, Peter Westerholm, Ari Väänänen, Marianna Virtanen, Jussi Vahtera, Töres Theorell, Sakari Suominen, Archana Singh-Manoux, Johannes Siegrist, Séverine Sabia, Reiner Rugulies, Jaana Pentti, Tuula Oksanen, Maria Nordin, Martin L. Nielsen, Michael G. Marmot, Linda L. Magnusson Hanson, Ida E. H. Madsen, Thorsten Lunau, Constanze Leineweber, Meena Kumari, Anne Kouvonen, Aki Koskinen, Markku Koskenvuo, Anders Knutsson, France Kittel, Karl-Heinz Jöckel, Matti Joensuu, Irene L. Houtman, Wendela E. Hoofman, Marcel Goldberg, Goedele A. Geuskens, Jane E. Ferrie, Raimund Erbel, Nico Dragano, Dirk De Bacquer, Els Clays, Annalisa Casini, Hermann Burr, Marianne Borritz, Sébastien Bonenfant, Jakob B. Bjorner, Lars Alfredsson, Mark Hamer, G. David Batty, and Mika Kivimäki

\* Correspondence to Dr. Eleonor Fransson, Institute of Environmental Medicine, Karolinska Institutet, Box 210, SE-171 77 Stockholm, Sweden (e-mail: [eleonor.fransson@ki.se](mailto:eleonor.fransson@ki.se)).

Initially submitted January 3, 2012; accepted for publication April 17, 2012.

Unfavorable work characteristics, such as low job control and too high or too low job demands, have been suggested to increase the likelihood of physical inactivity during leisure time, but this has not been verified in large-scale studies. The authors combined individual-level data from 14 European cohort studies (baseline years from 1985–1988 to 2006–2008) to examine the association between unfavorable work characteristics and leisure-time physical inactivity in a total of 170,162 employees (50% women; mean age, 43.5 years). Of these employees, 56,735 were reexamined after 2–9 years. In cross-sectional analyses, the odds for physical inactivity were 26% higher (odds ratio = 1.26, 95% confidence interval: 1.15, 1.38) for employees with high-strain jobs (low control/high demands) and 21% higher (odds ratio = 1.21, 95% confidence interval: 1.11, 1.31) for those with passive jobs (low control/low demands) compared with employees in low-strain jobs (high control/low demands). In prospective analyses restricted to physically active participants, the odds of becoming physically inactive during follow-up were 21% and 20% higher for those with high-strain (odds ratio = 1.21, 95% confidence interval: 1.11, 1.32) and passive (odds ratio = 1.20, 95% confidence interval: 1.11, 1.30) jobs at baseline. These data suggest that unfavorable work characteristics may have a spillover effect on leisure-time physical activity.

cohort studies; exercise; physical activity; psychosocial factors; working population

Abbreviations: Belstress, the Belgian Job Stress Study I; CI, confidence interval; DWECS, Danish Work Environment Cohort Study; FPS, Finnish Public Sector Study; Gazel, the Gaz et Electricité Cohort Study; HeSSup, Health and Social Support Study; HNR, Heinz Nixdorf Recall Study; IPAW, Intervention Project on Absence and Well-being; IPD-Work, Individual-Participant-Data Meta-Analysis in Working Populations; OR, odds ratio; POLS, Permanent Onderzoek LeefSituatie; PUMA, Burnout, Motivation, and Job Satisfaction Study; SES, socioeconomic status; SLOSH, Swedish Longitudinal Occupational Survey of Health; WOLF N, Work, Lipids, and Fibrinogen Study Norrland; WOLF S, Work, Lipids, and Fibrinogen Study Stockholm.

**Editor's note:** An invited commentary on this article appears on page 1090.

Physical inactivity is associated with increased risk of premature death and morbidity due to chronic disease, including cardiovascular disease, type 2 diabetes, and some cancers (1–8). According to the World Health Organization, almost 2 million deaths per year worldwide are attributable to physical inactivity (9). Despite numerous public health campaigns to increase regular physical activity in populations, reductions in sedentary lifestyle have been relatively modest. In the United States, for example, the proportion of the population that reported no leisure-time physical activity has decreased only 3 percentage points during the last 10 years, from 28% in 1998 to 25% in 2008 (10). For this reason, there is a need for increased understanding of factors that influence participation in leisure-time physical activities.

It has been hypothesized that stressful jobs characterized by high psychological demands and low control (also known as high-strain jobs) result in fatigue and greater need for recovery, increasing the likelihood of leisure-time passivity and sedentary behavior (11, 12). Another hypothesis proposes that passive, unchallenging jobs with few demands and little control over work can lead to reduced self-efficacy, which in turn may result in more passive lifestyles (11, 12). To date, however, empirical evidence for both hypotheses remains limited. Studies from Finland, Japan, the United States, Canada, the United Kingdom, and Sweden have provided support for a link between job strain and physical inactivity (13–21), although in some studies the association was attenuated after adjustment for covariates (14, 16, 18). In the Whitehall II Study of British civil servants, participants working in passive jobs were particularly likely to be physically inactive during their leisure time (22). However, other studies have failed to observe an association between high strain or passive jobs and leisure-time physical activity (12, 23). Heterogeneity in the association has also been observed by sex, education, and ethnicity (14, 17–19, 21).

To better characterize the associations between high-strain or passive jobs and leisure-time physical inactivity, we pooled data from 14 independent European cohort studies including over 170,000 men and women. Our aim was to examine whether leisure-time physical inactivity is more common among employees working in high-strain or passive jobs compared with those in low-strain jobs. As a subset of the participating studies had repeat data on both physical activity and work characteristics, we were also able to examine the temporal order of the association, that is, whether work characteristics predict leisure-time physical activity at follow-up, or, alternatively, if physical activity predicts moving into a high strain or passive job over the follow-up period.

## MATERIALS AND METHODS

This study is part of the Individual-Participant-Data Meta-Analysis in Working Populations (IPD-Work) Consortium of European cohort studies. Originally established

during the annual Four Centers Meeting in London, November 8, 2008, the collaboration has been joined by new cohort studies since. The overall aim of the IPD-Work Consortium is to establish reliable estimates of the effects of psychosocial risk factors at work on chronic disease, mental health, disability, and mortality, based on acquisition and synthesis of extensive individual-level data from multiple published and unpublished studies.

## Studies and participants

We pooled data from 14 prospective cohort studies based in 8 European countries: Belgium (the Belgian Job Stress Study I (Belstress)) (24, 25); Denmark (Danish Work Environment Cohort Study (DWECS) (26), Intervention Project on Absence and Well-being (IPAW) (27), Burnout, Motivation, and Job Satisfaction Study (PUMA) (28)); Finland (Finnish Public Sector Study (FPS) (29), Health and Social Support Study (HeSSup) (30), Still Working (31)); France (the Gaz et Electricité Cohort Study (Gazel) (32)); Germany (Heinz Nixdorf Recall Study (HNR) (33)); the Netherlands (Permanent Onderzoek LeefSituatie (POLS) (34)); Sweden (Swedish Longitudinal Occupational Survey of Health (SLOSH) (35, 36), Work, Lipids, and Fibrinogen Study Norrland (WOLF N) (37, 38) and Stockholm (WOLF S) (37, 39)); and the United Kingdom (Whitehall II Study) (40, 41). The years for the baseline data collection in the respective studies varied from 1985–1988 (Whitehall II) to 2006–2008 (SLOSH). Details of the design, recruitment, and ethical approval for the participating studies are presented in Web Appendix I available at <http://aje.oxfordjournals.org/>. Participants with complete data on leisure-time physical activity, the demand-control measures, sex, and age were included in the cross-sectional analyses in this study, which yielded an analytical sample of 85,132 employed men and 85,030 employed women. The prospective analyses were based on data from 56,735 participants.

## Work characteristics

Work characteristics were defined by using the Demand-Control Model, first proposed by Karasek (42) and further developed and tested by Karasek and Theorell (11). A description of the self-administered multiitem measures of job demands and job control in each participating study has been provided elsewhere (43). Briefly, all questions in the job demands and job control scales had Likert-type response formats. Mean response scores for the job demands items and for the job control items were computed for each participant. We then used the study-specific median scores as cutpoints for high and low demands (“high demands” being defined as scores strictly above the study-specific median score) and high and low job control (“low control” being defined as scores strictly below the study-specific median score). Four categories of jobs were defined: 1) low-strain jobs (low demands, high control); 2) passive jobs (low demands, low control); 3) active jobs (high demands, high control); and 4) high-strain jobs (high demands, low control). We also

evaluated the separate associations between job demands or job control and leisure-time physical inactivity using the study-specific quintiles for job demands and job control, respectively. Participants with missing data on more than half of the items for job demands or job control were excluded from the analysis ( $n = 1,793$ , 1% of the total population).

### Leisure-time physical inactivity

Physical activity was measured by self-report in all studies. The questions used to assess leisure-time physical activity differed between studies. Some studies had only questions on sports activities and exercise, while for other studies information was also available for other types of leisure-time physical activities, such as walking and cycling. As our main aim was to evaluate the association between work characteristics and leisure-time physical inactivity, we constructed a measure of physical inactivity defined as no or very little moderate or vigorous leisure-time physical activity or exercise based on the best available information in each study. Examples of definitions of physical inactivity are “no weekly leisure-time physical activity,” “no or very little exercise, only occasional walks,” and “sport activities a few times per year or less.” The

definitions of leisure-time physical inactivity in all the studies included in the analyses are presented in Table 1.

### Covariates

Sex and age were obtained from 1) either registers or self-reports during a medical examination (DWECS, FPS, Gazel, HNR, IPAW, PUMA, SLOSH, Still Working, WOLF N, and WOLF S) or 2) a questionnaire (Belstress, HeSSup, POLS, and Whitehall II). Age was treated as a continuous variable in the analyses. In addition, we included socioeconomic status (SES) as a covariate because SES may be related to both physical activity and psychosocial working conditions. SES was based on information on occupation obtained from register data (DWECS, FPS, Gazel, PUMA, and Still Working) or self-reports (Belstress, HNR, IPAW, POLS, SLOSH, WOLF N, WOLF S, Whitehall II). In HeSSup, SES was based on self-reported education. SES was classified as low, intermediate, or high. Self-employed participants and participants with missing data on SES were categorized as “others.” We included smoking status as an additional covariate because smoking is considered to be the leading preventable cause of illness, disability, and premature death, and previous findings suggest that job strain is associated with current smoking (44). Smoking

**Table 1.** Definitions of Leisure-Time Physical Inactivity Among the IPD-Work Consortium of European Cohort Studies (Baseline Years From 1985–1988 to 2006–2008)

| Study         | Leisure-Time Physical Inactivity   |
|---------------|--|
| Belstress     | No weekly physical activity  |
| DWECS         | Almost completely physically passive or light physical activity for less than 2 hours/week (e.g., reading, television, cinema)     |
| FPS           | Less than 0.5 hour of each (brisk walking, jogging, or running) per week   |
| Gazel         | No sport activities  |
| HeSSup        | Less than 0.5 hour of each (brisk walking, jogging, or running) per week   |
| HNR           | Less than 0.5 hour of moderate or vigorous physical activity per week  |
| IPAW          | Almost completely physically passive or light physical activity for less than 2 hours per week (e.g., reading, television, cinema) |
| POLS          | No exercise and less than 1 hour walking and less than 1 hour cycling for fun per week   |
| PUMA          | Almost completely physically passive or light physical activity for less than 2 hours per week (e.g., reading, television, cinema) |
| SLOSH         | No or very little exercise, only occasional walks  |
| Still Working | Sport activities less than a couple of times per month   |
| Whitehall II  | No moderate or vigorous exercise   |
| WOLF N        | No or very little exercise, only occasional walks  |
| WOLF S        | No or very little exercise, only occasional walks  |

Abbreviations: Belstress, the Belgian Job Stress Study I; DWECS, Danish Work Environment Cohort Study; FPS, Finnish Public Sector Study; Gazel, the Gaz et Electricité Cohort Study; HeSSup, Health and Social Support Study; HNR, Heinz Nixdorf Recall Study; IPAW, Intervention Project on Absence and Well-being; IPD-Work, individual-participant-data meta-analysis in working populations; POLS, Permanent Onderzoek LeefSituatie; PUMA, Burnout, Motivation, and Job Satisfaction Study; SLOSH, Swedish Longitudinal Occupational Survey of Health; WOLF N, Work, Lipids, and Fibrinogen Study Norrland; WOLF S, Work, Lipids, and Fibrinogen Study Stockholm.

status was self-reported in all studies and categorized as “never smoker,” “former smoker,” and “current smoker.”

### Statistical methods

Individual-level data were available for the following 10 studies: Belstress, FPS, Gazel, HeSSup, HNR, SLOSH, Still Working, Whitehall II, WOLF N, and WOLF S. Syntax and instructions for statistical analysis were provided for the investigators in the other studies (DWECS, IPAW, POLS, and PUMA), and they themselves calculated the study-specific results.

One- and two-stage meta-analyses of individual-participant data approaches were used (45–47). In the main cross-sectional analysis, we used 2-stage meta-analysis as we wanted to include all available cohort studies but had only aggregate data from 4 cohort studies (DWECS, IPAW, POLS, and PUMA). Stratified analyses were conducted by using 1-stage meta-analysis, excluding the 4 studies with only aggregate data.

In the 2-stage meta-analysis of the cross-sectional associations between work characteristics and physical inactivity, effect estimates and their standard errors were estimated by using logistic regression, separately for each study. The study-specific results were then pooled by random-effects meta-analysis (48). We calculated summary odds ratios and their 95% confidence intervals for individuals who were categorized as having passive, active, or high-strain jobs, comparing them with individuals with low-strain jobs. We adjusted the odds ratios for sex and age and for sex, age, SES, and smoking. Heterogeneity among study-specific estimates was assessed by using the  $I^2$  statistic (49).

It has been argued that the prevalence ratio is more appropriate than the odds ratio when evaluating the cross-sectional association between 2 variables, as the odds ratio tends to inflate the association if the prevalence of the outcome is high (50, 51). Therefore, we ran additional 2-stage individual-level meta-analyses using log binomial regression to estimate the pooled prevalence ratios of leisure-time physical inactivity in relation to work characteristics in the 10 studies where we had direct access to individual data.

In the 1-stage meta-analysis, we pooled all available individual-level data into 1 data set. To examine the robustness of the cross-sectional associations between the work characteristics and physical inactivity, we conducted subgroup analyses separately for men and women; participants aged less than 50 years and those aged 50 years or older; participants from high, intermediate, low, and “other” SES groups; and never smokers, former smokers, and current smokers. We also evaluated the separate effect of job demands and job control on leisure-time physical inactivity, using study-specific quintiles to categorize job demands and job control in 1-stage meta-analyses.

In addition, we used 1-stage individual-level meta-analysis to examine prospective associations between work characteristics and leisure-time physical inactivity in the 6 cohort studies (Belstress, FPS, HeSSup, SLOSH, Whitehall II, and WOLF N; total  $n = 56,735$ ) in which the work characteristics and physical activity measures had been repeated

2–9 years later and we had direct access to the data. In all studies, the same definition of work characteristics and physical activity was used at baseline and follow-up. These analyses were based on data from 56,735 participants. In the prospective analyses, we examined whether work characteristics at baseline predicted physical inactivity at follow-up in participants who were physically active at baseline, and if work characteristics at baseline predicted physical activity at follow-up in those who were inactive at baseline. To study potential reverse causality, we examined the association between physical inactivity at baseline and the likelihood of having a high-strain job versus having a low-strain, passive, or active job at follow-up among those in non-high-strain jobs at baseline. Corresponding analyses were undertaken to examine the odds of having a passive, active, or low-strain job at follow-up. In these analyses, we fitted a mixed-effects logistic regression model with study as the random effect and age, sex, SES, and smoking as covariates.

To study the effect of sample size on our findings, we ran sensitivity analysis taking 1% and 10% random samples from the pooled data set of 10 studies where we had access to individual data ( $n = 132,704$ ).

Study-specific logistic regression models were fitted with PROC GENMOD in SAS, version 9, software (SAS Institute, Inc., Cary, North Carolina) (Belstress, DWECS, FPS, Gazel, HeSSuP, HNR, IPAW, PUMA, SLOSH, Still Working, Whitehall II, WOLF N and WOLF S) or SPSS, version 17, statistical software (SPSS, Inc., Chicago, Illinois) (POLS). Meta-analysis was conducted by using R, version 2.11, library Meta (R Foundation for Statistical Computing, Vienna, Austria). Study-specific log binomial regression models were also fitted with PROC GENMOD in SAS, version 9. One-stage meta-analyses were fitted with SAS, version 9, PROC GLIMMIX.

## RESULTS

The characteristics of the study population are presented in Table 2. The mean age of the participants was 43.5 years, and 50% were women. The prevalence of leisure-time physical inactivity was 21% in the total sample, ranging from 7% in PUMA to 38% in Gazel. The proportion of participants with a high-strain job varied from 13% in the WOLF N study to 20% in SLOSH, while the prevalence of passive jobs ranged from 19% in DWECS to 34% in PUMA.

### Cross-sectional analyses

The overall prevalence of physical inactivity was 18.6%, 23.5%, 18.9%, and 23.9% among those with low-strain, passive, active, and high-strain jobs, respectively. There was strong evidence in the pooled analyses that participants with high-strain (odds ratio (OR) adjusted for age and sex = 1.36, 95% confidence interval (CI): 1.25, 1.48) and passive (OR = 1.34, 95% CI: 1.23, 1.47) jobs were more likely to be physically inactive during leisure time, compared with those working in low-strain jobs (Figure 1). Further adjustment for SES and smoking attenuated these




**Table 2.** Study Population Characteristics Among the IPD-Work Consortium of European Cohort Studies (Baseline Years From 1985–1988 to 2006–2008)<sup>a</sup>

| Study         | Total No. | Age, years |       | Female |    | Work Characteristics, % |         |        |             | Physical Inactivity |    |
|---------------|-----------|------------|-------|--------|----|-------------------------|---------|--------|-------------|---------------------|----|
|               |           | Mean       | Range | No.    | %  | Low Strain              | Passive | Active | High Strain | No.                 | %  |
| Belstress     | 20,397    | 45.4       | 33–61 | 4,775  | 23 | 26                      | 24      | 31     | 19          | 4,527               | 22 |
| DWECS         | 5,565     | 41.8       | 18–69 | 2,602  | 47 | 33                      | 19      | 26     | 22          | 841                 | 15 |
| FPS           | 46,588    | 44.6       | 17–64 | 37,544 | 81 | 33                      | 28      | 23     | 16          | 9,360               | 20 |
| Gazel         | 10,628    | 50.3       | 43–58 | 2,897  | 27 | 32                      | 20      | 33     | 14          | 4,001               | 38 |
| HeSSup        | 16,339    | 39.6       | 20–54 | 9,079  | 56 | 30                      | 27      | 26     | 18          | 3,601               | 22 |
| HNR           | 1,829     | 53.4       | 45–73 | 747    | 41 | 33                      | 30      | 25     | 12          | 226                 | 12 |
| IPAW          | 1,965     | 41.2       | 18–68 | 1,302  | 66 | 30                      | 20      | 32     | 18          | 151                 | 8  |
| POLS          | 24,753    | 38.3       | 15–85 | 10,169 | 41 | 32                      | 28      | 24     | 16          | 4,669               | 19 |
| PUMA          | 1,806     | 42.6       | 18–69 | 1,486  | 82 | 33                      | 34      | 18     | 15          | 130                 | 7  |
| SLOSH         | 10,853    | 47.6       | 19–68 | 5,848  | 54 | 28                      | 25      | 27     | 20          | 2,072               | 19 |
| Still Working | 8,969     | 40.8       | 18–65 | 2,044  | 23 | 34                      | 31      | 20     | 15          | 1,748               | 19 |
| Whitehall II  | 10,133    | 44.4       | 34–56 | 3,315  | 33 | 24                      | 33      | 29     | 14          | 1,652               | 16 |
| WOLF N        | 4,686     | 44.1       | 19–65 | 779    | 17 | 32                      | 24      | 31     | 13          | 1,254               | 27 |
| WOLF S        | 5,651     | 41.5       | 19–70 | 2,443  | 43 | 25                      | 33      | 26     | 16          | 1,321               | 23 |

Abbreviations: Belstress, the Belgian Job Stress Study I; DWECS, Danish Work Environment Cohort Study; FPS, Finnish Public Sector Study; Gazel, the Gaz et Electricité Cohort Study; HeSSup, Health and Social Support Study; HNR, Heinz Nixdorf Recall Study; IPAW, Intervention Project on Absence and Well-being; IPD-Work, individual-participant-data meta-analysis in working populations; POLS, Permanent Onderzoek LeefSituatie; PUMA, Burnout, Motivation, and Job Satisfaction Study; SLOSH, Swedish Longitudinal Occupational Survey of Health; WOLF N, Work, Lipids, and Fibrinogen Study Norrland; WOLF S, Work, Lipids, and Fibrinogen Study Stockholm.

<sup>a</sup> Participants with valid measures on work characteristics as defined by the Job Demand-Control Model, leisure-time physical activity, age, and sex.



**Figure 1.** Pooled results from cross-sectional 2-stage meta-analysis from the IPD-Work Consortium of European cohort studies (baseline years from 1985–1988 to 2006–2008). Odds ratios for leisure-time physical inactivity by job category are defined according to the Demand-Control Model as low strain (low demands, high control), passive (low demands, low control), active (high demands, high control), and high strain (high demands, low control). A, adjusted for sex and age ( $n=170,162$ ); B, adjusted for sex, age, socioeconomic status, and smoking ( $n=163,242$ ). CI, confidence interval; IPD-Work, individual-participant-data meta-analysis in working populations; OR, odds ratio.

associations, but the odds ratios remained statistically significant (OR = 1.26, 95% CI: 1.15, 1.38 and OR = 1.21, 95% CI: 1.11, 1.31, respectively). There was some heterogeneity between the studies in the meta-analyses of high-strain and passive jobs,  $I^2 = 77.5\%$  and  $76.7\%$ , respectively (the study-specific odds ratios are shown in Web Appendix II, Web Figure 1), supporting the use of a random-effects rather than a fixed-effect model. When the analysis was restricted to the 10 studies where we had direct access to the data, the sex-, age-, SES-, and smoking-adjusted odds ratios for leisure-time physical inactivity remained virtually the same as in the analysis including all 14 cohorts, with odds ratios of 1.26 (95% CI: 1.13, 1.40), 1.08 (95% CI: 1.01, 1.16), and 1.21 (95% CI: 1.10, 1.34), for the high-strain, active, and passive groups, respectively.

We repeated the 2-stage meta-analysis, estimating study-specific prevalence ratios using log binomial regression models and pooling them to give summary prevalence ratios across the 10 studies. These models yielded an age- and sex-adjusted summary prevalence ratio for leisure-time physical inactivity of 1.27 (95% CI: 1.18, 1.37) for high-strain jobs, 1.01 (95% CI: 0.95, 1.07) for active jobs, and 1.27 (95% CI: 1.17, 1.38) for passive jobs, compared with low-strain jobs. Additional adjustment for SES and smoking resulted in prevalence ratios of 1.18 (95% CI: 1.09, 1.28), 1.06 (95% CI: 1.00, 1.12), and 1.15 (95% CI: 1.07, 1.24), respectively.

**Table 3.** Cross-sectional Associations Between Work Characteristics<sup>a</sup> and Leisure-Time Physical Inactivity in Different Subgroups Among the IPD-Work Consortium of European Cohort Studies (Baseline Years From 1985–1988 to 2006–2008)

|                                    | No.    | Leisure-Time Physical Inactivity, % | Odds Ratio <sup>b</sup> | 95% CI     |
|------------------------------------|--------|-------------------------------------|-------------------------|------------|
| All ( <i>n</i> = 132,704)          |        |                                     |                         |            |
| Low strain                         | 39,903 | 19                                  | 1                       | Referent   |
| Passive                            | 35,870 | 25                                  | 1.29                    | 1.24, 1.33 |
| Active                             | 35,105 | 20                                  | 1.06                    | 1.02, 1.10 |
| High strain                        | 21,826 | 25                                  | 1.32                    | 1.27, 1.38 |
| Stratified by sex                  |        |                                     |                         |            |
| Men ( <i>n</i> = 65,043)           |        |                                     |                         |            |
| Low strain                         | 21,025 | 21                                  | 1                       | Referent   |
| Passive                            | 15,637 | 26                                  | 1.27                    | 1.21, 1.34 |
| Active                             | 19,623 | 20                                  | 1.05                    | 1.00, 1.11 |
| High strain                        | 8,758  | 27                                  | 1.36                    | 1.28, 1.44 |
| Women ( <i>n</i> = 67,661)         |        |                                     |                         |            |
| Low strain                         | 18,878 | 18                                  | 1                       | Referent   |
| Passive                            | 20,233 | 24                                  | 1.27                    | 1.21, 1.34 |
| Active                             | 15,482 | 19                                  | 1.08                    | 1.02, 1.14 |
| High strain                        | 13,068 | 24                                  | 1.28                    | 1.21, 1.35 |
| Stratified by age                  |        |                                     |                         |            |
| Age <50 years ( <i>n</i> = 86,650) |        |                                     |                         |            |
| Low strain                         | 25,830 | 17                                  | 1                       | Referent   |
| Passive                            | 23,537 | 23                                  | 1.28                    | 1.22, 1.34 |
| Active                             | 23,068 | 18                                  | 1.06                    | 1.01, 1.11 |
| High strain                        | 14,215 | 23                                  | 1.30                    | 1.23, 1.37 |
| Age ≥50 years ( <i>n</i> = 46,054) |        |                                     |                         |            |
| Low strain                         | 14,073 | 22                                  | 1                       | Referent   |
| Passive                            | 12,333 | 28                                  | 1.28                    | 1.20, 1.35 |
| Active                             | 12,037 | 23                                  | 1.05                    | 0.99, 1.12 |
| High strain                        | 7,611  | 28                                  | 1.34                    | 1.25, 1.43 |
| Stratified by SES                  |        |                                     |                         |            |
| Low SES ( <i>n</i> = 36,346)       |        |                                     |                         |            |
| Low strain                         | 8,483  | 24                                  | 1                       | Referent   |
| Passive                            | 15,267 | 28                                  | 1.23                    | 1.15, 1.31 |
| Active                             | 4,675  | 24                                  | 1.05                    | 0.96, 1.14 |
| High strain                        | 7,921  | 29                                  | 1.31                    | 1.22, 1.41 |

Table continues

To study the robustness of the associations further, we conducted 1-stage meta-analyses stratified by sex, age, SES, and smoking. The pattern of higher odds ratios for physical inactivity among those with high-strain or passive jobs was observed across all the subgroups examined (Table 3).

When levels of job demands and job control were analyzed separately, a clear association between job control and physical inactivity was observed, with odds for physical inactivity increasing at lower levels of job control. The association between job demands and physical inactivity was weaker, and there was evidence of an association only

in the highest quintile (Web Appendix II, Web Table 1; available at <http://aje.oxfordjournals.org/>).

### Prospective analysis

In the prospective analysis based on data from 6 studies, we observed increased odds of becoming physically inactive at follow-up among those who at baseline had high-strain (OR = 1.21, 95% CI: 1.11, 1.32) or passive (OR = 1.20, 95% CI: 1.11, 1.30) jobs compared with those who had low-strain jobs (Table 4). This analysis was restricted to those who were physically active (i.e., it

**Table 3.** Continued

|                                      | No.    | Leisure-Time<br>Physical Inactivity, % | Odds Ratio <sup>b</sup> | 95% CI     |
|--------------------------------------|--------|--|-------------------------|------------|
| Medium SES ( <i>n</i> = 63,530)      |        |  |                         |            |
| Low strain                           | 18,777 | 19                                     | 1                       | Referent   |
| Passive                              | 17,403 | 22                                     | 1.29                    | 1.23, 1.36 |
| Active                               | 15,862 | 20                                     | 1.05                    | 1.00, 1.11 |
| High strain                          | 11,488 | 23                                     | 1.33                    | 1.26, 1.41 |
| High SES ( <i>n</i> = 30,026)        |        |  |                         |            |
| Low strain                           | 11,706 | 17                                     | 1                       | Referent   |
| Passive                              | 2,483  | 20                                     | 1.25                    | 1.11, 1.40 |
| Active                               | 13,801 | 18                                     | 1.12                    | 1.05, 1.20 |
| High strain                          | 2,036  | 20                                     | 1.31                    | 1.16, 1.47 |
| Other SES ( <i>n</i> = 2,802)        |        |  |                         |            |
| Low strain                           | 937    | 19                                     | 1                       | Referent   |
| Passive                              | 717    | 26                                     | 1.36                    | 1.07, 1.73 |
| Active                               | 767    | 23                                     | 1.20                    | 0.95, 1.52 |
| High strain                          | 381    | 30                                     | 1.69                    | 1.28, 2.23 |
| Stratified by smoking                |        |  |                         |            |
| Never smokers ( <i>n</i> = 57,849)   |        |  |                         |            |
| Low strain                           | 17,285 | 17                                     | 1                       | Referent   |
| Passive                              | 15,124 | 22                                     | 1.34                    | 1.27, 1.43 |
| Active                               | 15,970 | 18                                     | 1.11                    | 1.05, 1.18 |
| High strain                          | 9,470  | 23                                     | 1.41                    | 1.32, 1.50 |
| Former smokers ( <i>n</i> = 45,076)  |        |  |                         |            |
| Low strain                           | 14,359 | 18                                     | 1                       | Referent   |
| Passive                              | 11,910 | 22                                     | 1.22                    | 1.15, 1.30 |
| Active                               | 11,996 | 18                                     | 1.02                    | 0.95, 1.08 |
| High strain                          | 6,811  | 22                                     | 1.25                    | 1.17, 1.35 |
| Current smokers ( <i>n</i> = 29,779) |        |  |                         |            |
| Low strain                           | 8,259  | 26                                     | 1                       | Referent   |
| Passive                              | 8,836  | 32                                     | 1.29                    | 1.20, 1.38 |
| Active                               | 7,139  | 26                                     | 1.04                    | 0.97, 1.12 |
| High strain                          | 5,545  | 32                                     | 1.29                    | 1.20, 1.40 |

Abbreviations: CI, confidence interval; IPD-Work, individual-participant-data meta-analysis in working populations; SES, socioeconomic status.

<sup>a</sup> Work characteristics defined according to the Demand-Control Model as low strain (low demands, high control), passive (low demands, low control), active (high demands, high control), and high strain (high demands, low control).

<sup>b</sup> Adjusted for age, sex, SES, and smoking. Study treated as random effect in the logistic model.

excluded the physically inactive) at baseline. In a further analysis—this time restricted to those who were physically inactive at baseline—we did not observe any clear association between work characteristics at baseline and becoming physically active at follow-up.

Our test of reverse causality showed physical inactivity at baseline to be associated with slightly increased odds of having a high-strain or passive job and with decreased odds of having an active or low-strain job at follow-up (Table 5).

### Sensitivity analysis and effect of sample size

Many studies of job strain have been based on sample sizes of 1,000–3,000; very few include more than 10,000 participants. Figure 2 shows that it is not possible to observe the association between job strain and physical inactivity in a 1% random sample (*n* = 1,327) of the pooled data. However, this association becomes significant in the 10% random sample comprising over 10,000 participants



**Table 4.** Prospective Associations Between Work Characteristics<sup>a</sup> at Baseline and Leisure-Time Physical Activity or Inactivity at Follow-up Among the IPD-Work Consortium of European Cohort Studies (Baseline Years From 1985–1988 to 2006–2008)<sup>b</sup>

| Baseline Population and Exposure at Baseline | No.    | Odds Ratio <sup>c</sup> | 95% CI     | Cases at Follow-up |    |
|--|--------|-------------------------|------------|--------------------|----|
|  |        |                         |            | No.                | %  |
| Physical activity at baseline (n = 45,927)   |        |                         |            |                    |    |
| Low strain                                   | 14,551 | 1 <sup>d</sup>          | Referent   | 1,685              | 12 |
| Passive                                      | 11,973 | 1.20 <sup>d</sup>       | 1.11, 1.30 | 1,806              | 15 |
| Active                                       | 12,334 | 1.07 <sup>d</sup>       | 0.99, 1.15 | 1,483              | 12 |
| High strain                                  | 7,059  | 1.21 <sup>d</sup>       | 1.11, 1.32 | 1,049              | 15 |
| Physical inactivity at baseline (n = 10,808) |        |                         |            |                    |    |
| Low strain                                   | 2,861  | 1 <sup>e</sup>          | Referent   | 1,416              | 49 |
| Passive                                      | 3,432  | 1.00 <sup>e</sup>       | 0.90, 1.11 | 1,634              | 48 |
| Active                                       | 2,545  | 1.10 <sup>e</sup>       | 0.98, 1.22 | 1,315              | 52 |
| High strain                                  | 1,970  | 0.98 <sup>e</sup>       | 0.87, 1.10 | 946                | 48 |

Abbreviations: Belstress, the Belgian Job Stress Study I; CI, confidence interval; IPD-Work, individual-participant-data meta-analysis in working populations.

<sup>a</sup> Work characteristics defined according to the Demand-Control Model as low strain (low demands, high control), passive (low demands, low control), active (high demands, high control), and high strain (high demands, low control).

<sup>b</sup> Studies and follow-up times: Belstress (4–8 years), Finnish Public Sector Study (2–4 years), Health and Social Support Study (5 years), Swedish Longitudinal Occupational Survey of Health (2 years), Whitehall II Study (3–9 years), and Work, Lipids, and Fibrinogen Study Norrland (3–7 years).

<sup>c</sup> Adjusted for age, sex, socioeconomic status, and smoking.

<sup>d</sup> Outcome at follow-up: physical inactivity.

<sup>e</sup> Outcome at follow-up: physical activity.

(n = 13,270). When repeating the random sampling procedure 5 times, it was observed that the estimated odds ratios vary substantially over the 5 different 1% samples. When using 10% samples, the estimates start to stabilize, but several estimates are still nonsignificant, as compared with the full sample size (Web Appendix II, Web Figures 2 and 3).

## DISCUSSION

We found robust cross-sectional and prospective associations between unfavorable work characteristics and leisure-time physical inactivity, with 21%–26% higher odds for inactivity among participants working in high-strain and passive jobs compared with those with low-strain jobs. Prospective analyses showed that high-strain and passive jobs also predicted change from a physically active to an inactive lifestyle. We found some support for a bidirectional association, as leisure-time physical inactivity at baseline to some extent predicted change in work characteristics; for example, physically inactive employees were more likely to

move into a high-strain or passive job compared with their physically active counterparts.

Individual-level meta-analysis of published and unpublished data, such as that used in the present study, is recognized as a strong study design as it reduces the possibility of publication bias that can limit the generalizability of evidence from single studies and literature-based meta-analyses (52). Our cross-sectional results are based on the largest data set to date on work characteristics (n > 170,000) and are in agreement with those of several previous studies. For example, an increased likelihood of low leisure-time physical activity among those with high strain and passive jobs compared with those in low-strain jobs was observed in 3,900 Swedish men (21) and 3,500 male white-collar workers in Canada (19), although no statistically significant associations were observed among women in these studies. Bennett et al. (17) found that people who reported job strain spent approximately 1 hour less in physical activities per week, compared with those who did not report job strain in a sample of 1,700 white individuals in Massachusetts. Furthermore, in a small study (n = 241) by Payne et al. (20), it was observed that employees reporting high strain did less exercise than employees in low-strain jobs (20). Choi et al. found that low-strain and active jobs were associated with a more physically active leisure-time compared with passive and high-strain jobs in 2,000 middle-aged American workers (18); Lallukka et al. (14) observed that a physically active leisure-time was more common in those with low strain and active jobs among 1,200 Finnish men and in those with low-strain jobs among the 5,000 Finnish women. However, in the latter 2 studies, the associations did not reach statistical significance at conventional levels in multivariable-adjusted models.

We observed similar odds ratios for physical inactivity in the high-strain and passive job groups. High-strain and passive jobs are both characterized by low control and, indeed, a subsidiary analysis revealed that the association between job control and leisure-time physical activity is much stronger than the association between job demands and physical activity. This is in agreement with some previous studies linking low control to a low level of physical activity (18, 53, 54), and it indicates that the association between work characteristics and leisure-time physical inactivity may be driven by the control dimension rather than by job demands. Our findings suggest that it makes little difference whether low control is combined with either high job demands (representing job strain) or low job demands (representing passive work).

Some earlier studies have failed to find an association between work characteristics and physical activity (12, 23). This inconsistency may be due to different definitions of physical activity but also to differences in the categorization of the psychosocial work characteristics. Furthermore, the smaller sample sizes in previous studies are likely to have introduced random error into the estimates, resulting in insufficient statistical power to detect relatively weak associations, such as those observed in the present meta-analysis (Figure 2; Web Appendix II).

Longitudinal data from 6 of the participating studies provided us with the opportunity to analyze temporal aspects

**Table 5.** Prospective Associations Between Leisure-Time Physical Activity or Inactivity at Baseline and Work Characteristics<sup>a</sup> at Follow-up Among the IPD-Work Consortium of European Cohort Studies (Baseline Years From 1985–1988 to 2006–2008)<sup>b</sup>

| Baseline Population and Exposure at Baseline                             | No.    | Odds Ratio <sup>c</sup> | 95% CI     | Cases at Follow-up |    |
|--|--------|-------------------------|------------|--------------------|----|
|  |        |                         |            | No.                | %  |
| All, except those with high-strain jobs at baseline ( <i>n</i> = 47,706) |        |                         |            |                    |    |
| Physical activity  | 38,868 | 1 <sup>d</sup>          | Referent   | 3,847              | 10 |
| Physical inactivity  | 8,838  | 1.15 <sup>d</sup>       | 1.07, 1.24 | 1,039              | 12 |
| All, except those with active jobs at baseline ( <i>n</i> = 41,846)      |        |                         |            |                    |    |
| Physical activity  | 33,583 | 1 <sup>e</sup>          | Referent   | 5,595              | 17 |
| Physical inactivity  | 8,263  | 0.89 <sup>e</sup>       | 0.83, 0.96 | 1,150              | 14 |
| All, except those with passive jobs at baseline ( <i>n</i> = 41,330)     |        |                         |            |                    |    |
| Physical activity  | 33,954 | 1 <sup>f</sup>          | Referent   | 4,763              | 14 |
| Physical inactivity  | 7,376  | 1.12 <sup>f</sup>       | 1.04, 1.20 | 1,196              | 16 |
| All, except those with low-strain jobs at baseline ( <i>n</i> = 39,323)  |        |                         |            |                    |    |
| Physical activity  | 31,376 | 1 <sup>g</sup>          | Referent   | 6,881              | 22 |
| Physical inactivity  | 7,947  | 0.89 <sup>g</sup>       | 0.84, 0.95 | 1,549              | 19 |

Abbreviations: Belstress, the Belgian Job Stress Study I; CI, confidence interval; IPD-Work, individual-participant-data meta-analysis in working populations.

<sup>a</sup> Work characteristics defined according to the Demand-Control Model as low strain (low demands, high control), passive (low demands, low control), active (high demands, high control), and high strain (high demands, low control).

<sup>b</sup> Studies and follow-up times: Belstress (4–8 years), Finnish Public Sector Study (2–4 years), Health and Social Support Study (5 years), Swedish Longitudinal Occupational Survey of Health (2 years), Whitehall II Study (3–9 years), and Work, Lipids, and Fibrinogen Study Norrland (3–7 years).

<sup>c</sup> Adjusted for age, sex, socioeconomic status, and smoking.

<sup>d</sup> Outcome at follow-up: high-strain job.

<sup>e</sup> Outcome at follow-up: active job.


<sup>f</sup> Outcome at follow-up: passive job.

<sup>g</sup> Outcome at follow-up: low-strain job.

of the link between work characteristics and leisure-time physical activity. This is in contrast to the vast majority of previous studies that have been based on cross-sectional data. Our main results support the idea that unfavorable work characteristics affect leisure-time physical activity. However, the association might be bidirectional because leisure-time physical inactivity also predicted, albeit weakly, adverse changes in work characteristics during follow-up. Certain personality traits may influence both participation in physical activity and the probability of having a job with more favorable characteristics. In a Finnish study, for example, it was observed that sustained involvement in physical activity in adolescence and young adult age was associated with reduced likelihood of reporting high strain jobs in early working life (15, 55), an association that was partly explained by personality traits (55).

This meta-analysis also has some limitations. First, data were based on multiexposure-multioutcome cohort studies that were not specifically designed to measure the impact of work characteristics on physical activity. Second, although we used a validated measure of work characteristics harmonized across all the studies (43), the number of items

and the wording of measures varied somewhat between the studies included. This may be one source of the heterogeneity observed between studies and one which may lead to some over- or underestimation of the magnitude of the associations. Furthermore, leisure-time physical activity was self-reported in all the studies, and this may have given rise to a degree of misclassification. However, we believe it is reasonable to assume that these misclassifications were largely independent of the work characteristics and, thus, if they had any effect, would rather attenuate than inflate the associations investigated in our study. Third, we observed no differences in the association between work characteristics and leisure-time physical activity by sex, age, SES, smoking status, or time of the study, but further research is needed to examine whether issues not assessed in this meta-analysis, such as social relations, physically demanding work, or sedentary work, economic circumstances, cultural contexts, and length of exposure to work characteristics, might modify this association. Fourth, our data were obtained from studies conducted in Scandinavia, Continental Europe, and the United Kingdom; it is unclear whether these findings are generalizable to other countries and



**Figure 2.** Estimated odds ratios and 95% confidence intervals for leisure-time physical inactivity in low-strain (L), passive (P), active (A), and high-strain (H) jobs based on different sample sizes from the IPD-Work Consortium of European cohort studies (baseline years from 1985–1988 to 2006–2008). Low-strain jobs are set as the referent category. The odds ratios are adjusted for sex, age, socioeconomic status, and smoking. A, 1% random sample ( $n=1,327$ ); B, 10% random sample ( $n=13,270$ ); C, total sample ( $n=132,704$ ). IPD-Work, individual-participant-data meta-analysis in working populations.

regions, such as Southern Europe, the United States, and Asia.

In conclusion, results from pooled data from over 170,000 participants in 14 European cohort studies provided consistent support for the hypothesis that unfavorable work characteristics have a spill-over effect on leisure-time physical activity. These results suggest that interventions to increase physical activity in the population may benefit from taking workplace factors into account.

## ACKNOWLEDGMENTS

Author affiliations: Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden (Eleonor I. Fransson, Lars Alfredsson); School of Health Sciences, Jönköping University, Jönköping, Sweden (Eleonor I. Fransson); Stress Research Institute, Stockholm University, Stockholm, Sweden (Eleonor I. Fransson, Hugo Westerlund, Töres Theorell, Linda L. Magnusson Hanson, Constanze Leineweber); Finnish Institute of Occupational Health, Helsinki and Turku, Finland (Katriina Heikkilä, Solja T. Nyberg, Ari Väänänen, Marianna Virtanen, Jussi Vahtera, Jaana Pentti, Tuula Oksanen, Aki Koskinen, Matti Joensuu, Mika Kivimäki); Versailles-Saint Quentin University, Versailles, France (Marie Zins, Marcel Goldberg, Sébastien Bonenfant); Centre for Research in Epidemiology

and Population Health, Institut National de la Santé et de la Recherche Médicale U1018, Villejuif, France (Marie Zins, Archana Singh-Manoux, Marcel Goldberg, Sébastien Bonenfant); Department of Epidemiology and Public Health, University College London, London, United Kingdom (Hugo Westerlund, Archana Singh-Manoux, Séverine Sabia, Michael G. Marmot, Meena Kumari, Jane E. Ferrie, Mark Hamer, G. David Batty, Mika Kivimäki); Department of Occupational and Environmental Medicine, Uppsala University, Uppsala, Sweden (Peter Westerholm); Department of Public Health, University of Turku, Turku, Finland (Jussi Vahtera, Sakari Suominen); Turku University Hospital, Turku, Finland (Jussi Vahtera); Folkhälsan Research Center, Helsinki, Finland (Sakari Suominen); Department of Medical Sociology, University of Düsseldorf, Düsseldorf, Germany (Johannes Siegrist); National Research Center for the Working Environment, Copenhagen, Denmark (Reiner Rugulies, Ida E. H. Madsen, Jakob B. Bjorner); Department of Public Health and Department of Psychology, University of Copenhagen, Copenhagen, Denmark (Reiner Rugulies); Department of Public Health and Clinical Medicine, Occupational and Environmental Medicine, Umeå University, Umeå, Sweden (Maria Nordin); Department of Occupational and Environmental Medicine, Bispebjerg University Hospital, Copenhagen, Denmark (Martin L. Nielsen, Marianne Borritz); Institute for Medical Informatics, Biometry, and Epidemiology, University Duisburg-Essen, Essen, Germany (Thorsten Lunau,

Karl-Heinz Jöckel, Nico Dragano); School of Sociology, Social Policy, & Social Work, Queen's University Belfast, Belfast, United Kingdom (Anne Kouvonen); Warsaw School of Social Sciences and Humanities, Wrocław, Poland (Anne Kouvonen); Department of Public Health, University of Helsinki, Helsinki, Finland (Markku Koskenvuo); Department of Health Sciences, Mid Sweden University, Sundsvall, Sweden (Anders Knutsson); School of Public Health, Université libre de Bruxelles, Brussels, Belgium (France Kittel, Annalisa Casini); Netherlands Organisation for Applied Scientific Research TNO, Hoofddorp, the Netherlands (Irene L. Houtman, Wendela E. Hoofman, Goedele A. Geuskens); School of Social and Community Medicine, University of Bristol, Bristol, United Kingdom (Jane E. Ferrie); Department of Cardiology, West-German Heart Center Essen, University Duisburg-Essen, Essen, Germany (Raimund Erbel); Department of Public Health, Ghent University, Ghent, Belgium (Dirk De Bacquer, Els Clays); and Federal Institute for Occupational Safety and Health, Berlin, Germany (Hermann Burr).

Authors' contribution: All authors participated in designing the study, generating hypotheses, interpreting the data, and writing and critically reviewing the paper. Eleonor I. Fransson and Mika Kivimäki analyzed the data and wrote the first draft of the paper.

The IPD-Work Consortium is supported by the European Union-based New and Emerging Risks in Occupational Safety and Health, anticipating and dealing with change in the workplace through coordination of the occupational safety and health risk research (NEW OSH ERA) research program (funded by the Finnish Work Environment Fund, the Swedish Research Council for Working Life and Social Research, the German Social Accident Insurance, the Danish National Research Centre for the Working Environment); the Academy of Finland (grant 132944); the BUPA Foundation (grant 22094477); and the Ministry of Social Affairs and Employment, the Netherlands. The Heinz Nixdorf Recall Study is funded by the Heinz Nixdorf Foundation, Germany, the German Ministry of Education and Science (BMBF), and the German Research Foundation (DFG). German National Accident Insurance (DGUV) supports analyses in the frame of the NEW OSH ERA project. The Dutch contribution (Permanent Onderzoek LeefSituatie) was funded by the Ministry of Social Affairs and Employment, the Netherlands. Eleonor I. Fransson was supported by the Swedish Council for Working Life and Social Research (grants 2009-2126, 2010-1823). David G. Batty is a Wellcome Trust Fellow.

Consortium funders had no role in the study design, data collection and analysis, or writing of the manuscript.

Preliminary findings from this study were presented at the PARG Symposium: Research on Physical Activity and Health in London: from Laboratory to Environmental and Population Studies, London, United Kingdom, November 28, 2011.

Conflict of interest: none declared.

## REFERENCES

- Hu G, Tuomilehto J, Silventoinen K, et al. The effects of physical activity and body mass index on cardiovascular, cancer and all-cause mortality among 47 212 middle-aged Finnish men and women. *Int J Obes (Lond)*. 2005;29(8):894–902.
- Trolle-Lagerros Y, Mucci LA, Kumle M, et al. Physical activity as a determinant of mortality in women. *Epidemiology*. 2005;16(6):780–785.
- Sofi F, Capalbo A, Cesari F, et al. Physical activity during leisure time and primary prevention of coronary heart disease: an updated meta-analysis of cohort studies. *Eur J Cardiovasc Prev Rehabil*. 2008;15(3):247–257.
- Nocon M, Hiemann T, Muller-Riemenschneider F, et al. Association of physical activity with all-cause and cardiovascular mortality: a systematic review and meta-analysis. *Eur J Cardiovasc Prev Rehabil*. 2008;15(3):239–246.
- Gill JM, Cooper AR. Physical activity and prevention of type 2 diabetes mellitus. *Sports Med*. 2008;38(10):807–824.
- Monninkhof EM, Elias SG, Vlems FA, et al. Physical activity and breast cancer: a systematic review. *Epidemiology*. 2007;18(1):137–157.
- Voskuil DW, Monninkhof EM, Elias SG, et al. Physical activity and endometrial cancer risk, a systematic review of current evidence. *Cancer Epidemiol Biomarkers Prev*. 2007;16(4):639–648.
- Harriss DJ, Atkinson G, Batterham A, et al. Lifestyle factors and colorectal cancer risk (2): a systematic review and meta-analysis of associations with leisure-time physical activity. *Colorectal Dis*. 2009;11(7):689–701.
- World Health Organization. *The World Health Report: 2002: Reducing the Risks, Promoting Healthy Life*. Geneva, Switzerland: World Health Organization; 2002.
- Centers for Disease Control and Prevention. Physical activity statistics: 1988–2008 no leisure-time physical activity trend chart. Atlanta, GA: Centers for Disease Control and Prevention; 2010. ([http://www.cdc.gov/nccdphp/dnpa/physical/stats/leisure\\_time.htm](http://www.cdc.gov/nccdphp/dnpa/physical/stats/leisure_time.htm)). (Accessed September 13, 2011).
- Karasek R, Theorell T. *Healthy Work: Stress, Productivity and the Reconstruction of Working Life*. New York, NY: Basic Books, Inc; 1990.
- Landsbergis PA, Schnall PL, Deitz DK, et al. Job strain and health behaviors: results of a prospective study. *Am J Health Promot*. 1998;12(4):237–245.
- Kouvonen A, Kivimäki M, Elovainio M, et al. Job strain and leisure-time physical activity in female and male public sector employees. *Prev Med*. 2005;41(2):532–539.
- Lallukka T, Sarlio-Lahteenkorva S, Roos E, et al. Working conditions and health behaviours among employed women and men: the Helsinki Health Study. *Prev Med*. 2004;38(1):48–56.
- Yang X, Telama R, Hirvensalo M, et al. The benefits of sustained leisure-time physical activity on job strain. *Occup Med (Lond)*. 2010;60(5):369–375.
- Tsutsumi A, Kayaba K, Yoshimura M, et al. Association between job characteristics and health behaviors in Japanese rural workers. *Int J Behav Med*. 2003;10(2):125–142.
- Bennett GG, Wolin KY, Avrunin JS, et al. Does race/ethnicity moderate the association between job strain and leisure time physical activity? *Ann Behav Med*. 2006;32(1):60–67.
- Choi B, Schnall PL, Yang H, et al. Psychosocial working conditions and active leisure-time physical activity in middle-aged US workers. *Int J Occup Med Environ Health*. 2010;23(3):239–253.
- Brisson C, Larocque B, Moisan J, et al. Psychosocial factors at work, smoking, sedentary behavior, and body mass index:



- a prevalence study among 6995 white collar workers. *J Occup Environ Med.* 2000;42(1):40–46.
20. Payne N, Jones F, Harris PR. The impact of job strain on the predictive validity of the theory of planned behaviour: an investigation of exercise and healthy eating. *Br J Health Psychol.* 2005;10(pt 1):115–131.
  21. Wemme KM, Rosvall M. Work related and non-work related stress in relation to low leisure time physical activity in a Swedish population. *J Epidemiol Community Health.* 2005; 59(5):377–379.
  22. Gimeno D, Elovainio M, Jokela M, et al. Association between passive jobs and low levels of leisure-time physical activity: the Whitehall II cohort study. *Occup Environ Med.* 2009;66(11):772–776.
  23. van Loon AJ, Tijhuis M, Surtees PG, et al. Lifestyle risk factors for cancer: the relationship with psychosocial work environment. *Int J Epidemiol.* 2000;29(5):785–792.
  24. Coetsier P, De Backer G, De Corte W, et al. Etude belge du stress au travail: aperçu du modèle de recherche et des outils d'investigation. (In French). *Rev Psychol Psychom.* 1996; 17(4):17–35.
  25. Clays E, De Bacquer D, Leynen F, et al. Long-term changes in the perception of job characteristics: results from the Belstress II—study. *J Occup Health.* 2006;48(5):339–346.
  26. Burr H, Bjorner JB, Kristensen TS, et al. Trends in the Danish work environment in 1990–2000 and their associations with labor-force changes. *Scand J Work Environ Health.* 2003;29(4):270–279.
  27. Nielsen ML, Kristensen TS, Smith-Hansen L. The Intervention Project on Absence and Well-being (IPAW): design and results from the baseline of a 5-year study. *Work & Stress.* 2002;16(3):191–206.
  28. Borritz M, Rugulies R, Christensen KB, et al. Burnout as a predictor of self-reported sickness absence among human service workers: prospective findings from three year follow up of the PUMA study. *Occup Environ Med.* 2006;63(2): 98–106.
  29. Kivimaki M, Lawlor DA, Davey Smith G, et al. Socioeconomic position, co-occurrence of behavior-related risk factors, and coronary heart disease: the Finnish Public Sector Study. *Am J Public Health.* 2007;97(5):874–879.
  30. Korkeila K, Suominen S, Ahvenainen J, et al. Non-response and related factors in a nation-wide health survey. *Eur J Epidemiol.* 2001;17(11):991–999.
  31. Vaananen A, Murray M, Koskinen A, et al. Engagement in cultural activities and cause-specific mortality: prospective cohort study. *Prev Med.* 2009;49(2-3):142–147.
  32. Goldberg M, Leclerc A, Bonenfant S, et al. Cohort profile: the GAZEL Cohort Study. *Int J Epidemiol.* 2007;36(1):32–39.
  33. Stang A, Moebus S, Dragano N, et al. Baseline recruitment and analyses of nonresponse of the Heinz Nixdorf Recall Study: identifiability of phone numbers as the major determinant of response. *Eur J Epidemiol.* 2005;20(6):489–496.
  34. de Groot W, Dekker R. *The Dutch System of Official Social Surveys.* EuReporting Working Paper No. 30. Mannheim, Germany: Mannheim Centre for European Social Research; 2001.
  35. Hasson D, Theorell T, Westerlund H, et al. Prevalence and characteristics of hearing problems in a working and non-working Swedish population. *J Epidemiol Community Health.* 2010;64(5):453–460.
  36. Magnusson Hanson LL, Theorell T, Oxenstierna G, et al. Demand, control and social climate as predictors of emotional exhaustion symptoms in working Swedish men and women. *Scand J Public Health.* 2008;36(7):737–743.
  37. Alfredsson L, Hammar N, Fransson E, et al. Job strain and major risk factors for coronary heart disease among employed males and females in a Swedish study on work, lipids and fibrinogen. *Scand J Work Environ Health.* 2002;28(4):238–248.
  38. Åkerstedt T, Nordin M, Alfredsson L, et al. Predicting changes in sleep complaints from baseline values and changes in work demands, work control, and work preoccupation—The WOLF-Project. *Sleep Med.* 2012;13(1):73–80.
  39. Peter R, Alfredsson L, Hammar N, et al. High effort, low reward, and cardiovascular risk factors in employed Swedish men and women: baseline results from the WOLF Study. *J Epidemiol Community Health.* 1998;52(9):540–547.
  40. Marmot MG, Smith GD, Stansfeld S, et al. Health inequalities among British civil servants: the Whitehall II Study. *Lancet.* 1991;337(8754):1387–1393.
  41. Marmot M, Brunner E. Cohort profile: the Whitehall II Study. *Int J Epidemiol.* 2005;34(2):251–256.
  42. Karasek R. Job demands, job decision latitude, and mental strain: implications for job redesign. *Adm Sci Q.* 1979; 24(2):285–308.
  43. Fransson EI, Nyberg ST, Heikkilä K, et al. Comparison of alternative versions of the job demand-control scales in 17 European cohort studies: the IPD-Work Consortium. *BMC Public Health.* 2012;12(1):62. (doi:10.1186/1471-2458-12-62).
  44. Heikkilä K, Nyberg ST, Fransson EI, et al. Job strain and tobacco smoking: an individual-participant data meta-analysis of 166 130 adults in 15 European studies. *PLoS One.* 2012; 7(7):e35463. (doi:10.1371/journal.pone.0035463).
  45. Riley RD, Lambert PC, Abo-Zaid G. Meta-analysis of individual participant data: rationale, conduct, and reporting. *BMJ.* 2010;340:c221. (doi:10.1136/bmj.c221).
  46. Simmonds MC, Higgins JP, Stewart LA, et al. Meta-analysis of individual patient data from randomized trials: a review of methods used in practice. *Clin Trials.* 2005;2(3):209–217.
  47. Stewart L, Tierney J, Clarke M. Chapter 18: reviews of individual patient data. In: Higgins J, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions*, version 5.0.1 (updated September 2008). Chichester, United Kingdom: The Cochrane Collaboration; 2008:18.1–18.7.
  48. Altman DG, Egger M, Davey Smith G. *Systematic Reviews in Health Care Meta-Analysis in Context.* London, United Kingdom: BMJ Books; 2001.
  49. Higgins JP, Thompson SG, Deeks JJ, et al. Measuring inconsistency in meta-analyses. *BMJ.* 2003;327(7414): 557–560.
  50. Spiegelman D, Hertzmark E. Easy SAS calculations for risk or prevalence ratios and differences. *Am J Epidemiol.* 2005;162(3):199–200.
  51. Thompson ML, Myers JE, Kriebel D. Prevalence odds ratio or prevalence ratio in the analysis of cross sectional data: what is to be done? *Occup Environ Med.* 1998;55(4):272–277.
  52. Stewart LA, Parmar MK. Meta-analysis of the literature or of individual patient data: is there a difference? *Lancet.* 1993;341(8842):418–422.
  53. Hellerstedt WL, Jeffery RW. The association of job strain and health behaviours in men and women. *Int J Epidemiol.* 1997;26(3):575–583.
  54. Smith PM, Frank JW, Mustard CA, et al. Examining the relationships between job control and health status: a path analysis approach. *J Epidemiol Community Health.* 2008; 62(1):54–61.
  55. Yang X, Telama R, Hirvensalo M, et al. Sustained involvement in youth sports activities predicts reduced chronic job strain in early midlife. *J Occup Environ Med.* 2010;52(12):1154–1159.