

## Modeling intracellular silent oscillations and rhythmic discharge in olfactory bulb mitral cells

Nicolas Fourcaud-Trocmé, Virginie Briffaud, Corine Amat

► **To cite this version:**

Nicolas Fourcaud-Trocmé, Virginie Briffaud, Corine Amat. Modeling intracellular silent oscillations and rhythmic discharge in olfactory bulb mitral cells. Gennady S Cymbalyuk and Astrid A Prinz. the Twenty Second Annual Computational Neuroscience Meeting: CNS\*2013, Jul 2013, paris, France. Biomed Central, 14 (Suppl 1), pp.P8, 2013. <inserm-00842316>

**HAL Id: inserm-00842316**

**<https://www.hal.inserm.fr/inserm-00842316>**

Submitted on 8 Jul 2013

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POSTER PRESENTATION

Open Access

# Modeling intracellular silent oscillations and rhythmic discharge in olfactory bulb mitral cells

Nicolas Fourcaud-Trocmé<sup>1,2\*</sup>, Virginie Briffaud<sup>1,2</sup>, Corine Amat<sup>1,2</sup>

From Twenty Second Annual Computational Neuroscience Meeting: CNS\*2013  
Paris, France. 13-18 July 2013

In mammals, the main olfactory bulb (MOB) is driven by air flow respiratory input into the nose. This slow (2-10 Hz) rhythmic input can be observed both at the local field potential (LFP) level, single cell discharge pattern level and intracellular slow oscillations of mitral cells (principal cells of the MOB). Using intracellular recordings in freely breathing anesthetized rats, we examined membrane potential slow oscillations (MPSO) in mitral cells and their relationship with their discharge patterns [1]. Interestingly, we found cells which showed no MPSO at resting membrane potential but had nevertheless a respiratory-synchronous discharge. Changing the excitability level of these cells with hyperpolarizing current injection favored the apparition of an MPSO while decreasing the cell discharge. We called such a dynamic *silent MPSO*. In this study, we use a modeling approach to show how such a silent MPSO can exist in mitral cells and favor the synchronization of cell discharge to the MOB slow rhythmic input.

First, using a simple integrate-and-fire (IF) model, we showed that the presence of strong and balanced excitatory and inhibitory inputs with similar oscillatory dynamics led the model to display a silent MPSO. This silent MPSO is characterized by a flat membrane potential in spite of strong synaptic conductance oscillations. Second, assuming that the slow oscillatory input could be decomposed as a mean input and stochastic fluctuation with oscillating amplitudes, we showed that the model could display a slow rhythmic discharge similar to our experimental observations.

However, MOB mitral cells have a complex anatomy with (i) an apical dendrite and a tuft receiving both excitatory nose input and local inhibitory regulation, and

(ii) long lateral dendrites which receive a large recurrent inhibitory input from the MOB network. We thus wondered how our previous modeling results could generalize in this case. To this aim, we used a previously developed detailed mitral cell model and showed with numerical simulations that silent MPSO could be generated in this model. However, the precise dynamics of the oscillations depended on the main inhibitory input of the mitral cell. If most of the inhibition comes from the tuft, then the silent MPSO can quite easily be generated in the cell, and partial imprecision in the excitation-inhibition balance can be hindered by the apical dendrite filtering. However if a fraction of the inhibition is coming from the lateral dendrites, the silent MPSOs are more difficult to stabilize and the resulting MPSOs have complex shapes resembling some of our intracellular recordings.

Overall, we found that mitral cell silent MPSOs with respiratory-patterned discharge, that we observed experimentally, can indeed be a consequence of the precise balance of their slow oscillatory excitatory and inhibitory inputs.

#### Author details

<sup>1</sup>INSERM, U1028; CNRS, UMR5292; Lyon Neuroscience Research Center, "Olfaction: from coding to memory" Team, Lyon, F-69000, France. <sup>2</sup>University Lyon 1, Villeurbanne, F-69000, France.

Published: 8 July 2013

#### References

1. Briffaud V, Fourcaud-Trocmé N, Messaoudi B, Buonviso N, Amat C: The relationship between respiration-related membrane potential slow oscillations and discharge patterns in mitral/tufted cells: what are the rules? *PLoS One* 2012, **7**(8):e43964.
2. Migliore M, Shepherd GM: Dendritic action potentials connect distributed dendrodendritic microcircuits. *J Comput Neurosci* 2008, **24**:207-221.

doi:10.1186/1471-2202-14-S1-P8

Cite this article as: Fourcaud-Trocmé et al.: Modeling intracellular silent oscillations and rhythmic discharge in olfactory bulb mitral cells. *BMC Neuroscience* 2013 **14**(Suppl 1):P8.

\* Correspondence: nfourcau@olfac.univ-lyon1.fr

<sup>1</sup>INSERM, U1028; CNRS, UMR5292; Lyon Neuroscience Research Center, "Olfaction: from coding to memory" Team, Lyon, F-69000, France  
Full list of author information is available at the end of the article