

**Dynamic changes in myocardial architecture after  
reperfused acute myocardial infarction (AMI): insights  
from the prospective REMI study (REmodeling after  
Myocardial Infarction)**

Lin Zhang, Olivier Huttin, Frédéric Moulin, Pierre-Yves Marie, Damien  
Mandry

► **To cite this version:**

Lin Zhang, Olivier Huttin, Frédéric Moulin, Pierre-Yves Marie, Damien Mandry. Dynamic changes in myocardial architecture after reperfused acute myocardial infarction (AMI): insights from the prospective REMI study (REmodeling after Myocardial Infarction). *Journal of Cardiovascular Magnetic Resonance*, BioMed Central, 2014, 16 (Suppl 1), pp.P223. <inserm-00932135>

**HAL Id: inserm-00932135**

**<http://www.hal.inserm.fr/inserm-00932135>**

Submitted on 16 Jan 2014

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POSTER PRESENTATION

Open Access

# Dynamic changes in myocardial architecture after reperfused acute myocardial infarction (AMI): insights from the prospective REMI study (REmodeling after Myocardial Infarction)

Lin Zhang<sup>2\*</sup>, Olivier Huttin<sup>1,2</sup>, Frédéric Moulin<sup>1</sup>, Pierre-Yves Marie<sup>1,3</sup>, Damien Mandry<sup>1,3</sup>

From 17th Annual SCMR Scientific Sessions  
New Orleans, LA, USA. 16-19 January 2014

## Background

Following an acute infarction, a dynamic healing process initiates via extracellular substrate deposition and turnover, resulting in ongoing remodeling of infarct territory and global function change of left ventricle. Our prospective study was designed to investigate by CMR the dynamic changes in myocardial architecture at 2-4 days and 6 months following a reperfused AMI.

## Methods

66 consecutive patients who fulfilled inclusion criteria underwent two CMRs: 2-4 days (baseline) and 6 months (follow-up) after the AMI, comprising SSFP images for analysis of left ventricular functions and late gadolinium enhancement (LGE) MRI (0.1 mmol/Kg; DOTAREM<sup>®</sup>, GUERBET, Roissy, France) for infarct analysis using a threshold method. Of 132 CMRs, 6 (4 from baseline) were non-diagnostic owing to poor EKG triggering and/or inability of the patients to hold their breath; 5 patients without evidence of infarct scar by LGE were also excluded. Hence, 57 patients (50 male; 56.7 ± 13.2 yo) were eventually analyzed. Total infarct was determined as area with signal intensity (SI) above mean+5SD of remote myocardium and was further divided into peri-infarct zone (PIZ) and infarct core by introducing a 7SD threshold; microvascular obstruction was manually included into the core; a core to PIZ ratio(C/PIZ) was calculated and a transmural score was computed using a 5-point scale on the 16-segment model derived from the AHA model. Paired t-test was used for comparison between the two

MRI and Pearson's correlation coefficients were calculated to assess prediction of LV parameters.

## Results

14 patients (25%) had an AMI in the LAD territory; 18 patients (31.6%) presented microvascular obstruction on LGE images at baseline. Dynamic changes of infarct indices and LV functional parameters are detailed in table 1; LVEF improved significantly ( $p < 0.0001$ ), owing to a decrease of LVESV ( $p < 0.05$ ), possibly related to stunning at subacute phase; the increase of LVEDV remained mild, possibly indicative that most of LV remodeling occurred prior to the baseline MRI. Total infarct, core and PIZ mass all diminished significantly ( $p < 0.0001$  for all). Both C/PIZ and transmural score remained identical. Both total scar and infarct core masses at baseline were predictive of LVEF, LVEDV and LVESV at follow-up (table 2), but not of the change in LV functional parameters.

## Conclusions

Following reperfused AMI, our study demonstrated a significant improvement in LVEF, related to an improved contractility due to stunning in the subacute phase. Myocardial damage characteristics by LGE showed a decrease of infarct size, owing to resorption of myocardial edema, in the same amount between its two components, core and PIZ.

## Funding

Grant from the French Ministry of Health Support from Laboratoire Guerbet, Roissy, France: supply of contrast medium.

<sup>2</sup>IADI U947, INSERM, Nancy, France

Full list of author information is available at the end of the article

**Table 1 Infarct index and LV functional parameters at baseline and follow-up.**

	Baseline (N = 57)	Follow-up (N = 57)	Change	%Change	P value
LV mass, g	97.2 ± 19.1	92.9 ± 20.4	-4.2 ± 15.2	-3.7 ± 15.8	<0.05
LVEF,%	43.0 ± 6.7	49.4 ± 7.7	6.4 ± 5.8	15.7 ± 15.3	<0.0001
LVEDV, mL	177.4 ± 30.4	183.0 ± 36.7	5.6 ± 22.1	3.4 ± 12.7	NS (0.061)
LVESV, mL	100.3 ± 22.7	94.4 ± 30.0	-5.9 ± 19.5	-6.2 ± 18.5	<0.05
Infarct mass, g	25.4 ± 11.8	17.0 ± 7.9	-8.5 ± 8.0	-30.6 ± 20.1	<0.0001
Infarct mass, % of LV mass	25.4 ± 8.8	17.8 ± 6.3	-7.6 ± 6.3	-27.5 ± 20.0	<0.0001
Core mass, g	12.3 ± 6.0	10.5 ± 5.7	-2.8 ± 5.3	-17.2 ± 41.3	<0.0001
Core mass, % of LV mass	13.3 ± 4.6	10.8 ± 4.8	-2.4 ± 4.6	-14.6 ± 37.4	<0.0001
PIZ mass, g	9.6 ± 4.5	6.5 ± 2.9	-3.2 ± 3.5	-26.6 ± 26.3	<0.0001
PIZ mass, % of LV mass	9.7 ± 3.6	6.9 ± 2.7	-2.7 ± 3.2	-22.8 ± 28.2	<0.0001
Core/PIZ	1.6 ± 0.8	1.7 ± 0.8	0.1 ± 1.0	28.4 ± 80.5	NS(0.405)
Transmurality score	1.30 ± 0.31	1.21 ± 0.43	-0.1 ± 0.45	-3.53 ± 35.68	NS(0.118)

LV indicates left ventricular; LVEF, LV ejection fraction; LVEDV, LV end-diastolic volume; LVESV, LV end-systolic volume; PIZ, peri-infarct zone; NS, no significant

**Table 2 Pearson's correlation coefficients between infarct indices and LV remodeling.**

	LVEF <sub>2</sub> ,%	ΔLVEF,%	LVEDV <sub>2</sub> , mL	ΔLVEDV, mL	LVESV <sub>2</sub> , mL	ΔLVESV, mL
Infarct mass <sub>1</sub> , g	-0.426**	-0.096	0.598**	0.228	0.601**	0.207
Infarct mass <sub>2</sub> , g	-0.486**	-0.198	0.615**	0.420**	0.625**	0.358**
Core mass <sub>1</sub> , g	-0.408**	-0.157	0.500**	0.137	0.518**	0.189
Core mass <sub>2</sub> , g	-0.497**	-0.248	0.561**	0.491**	0.598**	0.440**
PIZ mass <sub>1</sub> , g	-0.144	0.08	0.519**	0.111	0.436**	0.100
PIZ mass <sub>2</sub> , g	-0.341**	-0.05	0.561**	0.175	0.518**	0.108
Core/PIZ <sub>1</sub>	-0.23	-0.167	-0.109	-0.008	0.012	0.053
Core/PIZ <sub>2</sub>	-0.291*	-0.191	0.129	0.415**	0.209	0.359**
%change Core/PIZ	-0.198	-0.04	0.281*	0.409**	0.299*	0.319*
Transmurality score <sub>1</sub>	-0.274	0.011	0.362**	0.123	0.331*	0.007
Transmurality score <sub>2</sub>	-0.160	0.089	0.238	0.205	0.204	0.000

\*p < 0.05; \*\*p < 0.01; subscript 1 and 2 indicate baseline and follow-up data, respectively.

#### Authors' details

<sup>1</sup>CHU Nancy, Nancy, France. <sup>2</sup>IADI U947, INSERM, Nancy, France. <sup>3</sup>Universite de Lorraine, Nancy, France.

Published: 16 January 2014

doi:10.1186/1532-429X-16-S1-P223

**Cite this article as:** Zhang *et al.*: Dynamic changes in myocardial architecture after reperfused acute myocardial infarction (AMI): insights from the prospective REMI study (REmodeling after Myocardial Infarction). *Journal of Cardiovascular Magnetic Resonance* 2014 **16**(Suppl 1): P223.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

