

Detailed clinical characteristics for each patient

Patients (n = 20)	Gender (F/M)	Age (yrs)	Side of Hemiparesis	Time after stroke (mo)	Fugl-Meyer (/34)	PASS (/36)	FIM (/126)
S1	M	58	L	3	29	32	116
S2	M	46	R	3	28	34	114
S3	M	50	L	3	8	33	92
S4	M	58	L	2	22	28	99
S5	F	50	L	2	26	34	102
S6	M	52	R	16	25	34	110
S7	M	27	R	7	11	23	67
S8	M	51	L	2	22	33	94
S9	M	48	L	2	21	33	102
S10	F	25	L	15	33	35	115
S11	M	55	R	8	33	36	107
S12	F	66	L	1	32	36	116
S13	F	49	L	3	8	33	114
S14	F	79	R	1	27	34	123
S15	M	55	R	36	13	33	100
S16	M	19	R	31	30	36	126
S17	M	48	L	22	19	33	115
S18	F	29	R	10	30	36	118
S19	M	52	L	2	21	33	112
S20	M	76	R	37	22	34	106
Mean	14M/6 F	49.7	9R/11 L	10.3	23.0	33.2	107.4
(SD)		(15.5)		(12.1)	(7.9)	(3.0)	(13.2)

F, female; L, left; M, male; R, right; PASS, Postural Assessment Scale for Stroke Patients;

FIM, Functional Independence Measure.