

PPACK-Desmodus rotundus salivary plasminogen activator (cDSPAalpha1) prevents the passage of tissue-type plasminogen activator (rt-PA) across the blood-brain barrier and neurotoxicity.

Benoit Roussel, Yannick Hommet, Richard Macrez, Torsten Schulz, Karl-Uwe Petersen, Vincent Berezowski, Roméo Cecchelli, Carine Ali, Denis Vivien

► **To cite this version:**

Benoit Roussel, Yannick Hommet, Richard Macrez, Torsten Schulz, Karl-Uwe Petersen, et al.. PPACK-Desmodus rotundus salivary plasminogen activator (cDSPAalpha1) prevents the passage of tissue-type plasminogen activator (rt-PA) across the blood-brain barrier and neurotoxicity.. Thrombosis and Haemostasis, Schattauer, 2009, 102 (3), pp.606-8. 10.1160/TH09-02-0114 . inserm-01296793

HAL Id: inserm-01296793

<https://www.hal.inserm.fr/inserm-01296793>

Submitted on 1 Apr 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



PPACK-Desmodus rotundus salivary plasminogen activator (cDSPA α 1) prevents the passage of tissue-type plasminogen activator (rt-PA) across the blood-brain barrier and neurotoxicity

Journal:	<i>Thrombosis and Haemostasis</i>
Manuscript ID:	draft
Manuscript Type:	Letters to the Editor
Category:	Basic Science
Date Submitted by the Author:	
Complete List of Authors:	Roussel, Benoit; INSERM U919, Serine Proteases and Pathophysiology of the Neurovascular Unit, UMR-CNRS 6232 CINAPS, Cyceron, Universite de Caen Basse Normandie Hommet, Yannick; INSERM U919, Serine Proteases and Pathophysiology of the Neurovascular Unit, UMR-CNRS 6232 CINAPS, Cyceron, Universite de Caen Basse Normandie Macrez, Richard; INSERM U919, Serine Proteases and Pathophysiology of the Neurovascular Unit, UMR-CNRS 6232 CINAPS, Cyceron, Universite de Caen Basse Normandie Schulz, Torsten; PAION Deutschland GmbH Petersen, Karl-Uwe; PAION Deutschland GmbH Berezowski, Vincent; Laboratoire de Physiopathologie de la Barrière Hémato-encéphalique, Université d'Artois, EA 2465 - IMPRT 114 Cecchelli, Romeo; Laboratoire de Physiopathologie de la Barrière Hémato-encéphalique, Université d'Artois, EA 2465 - IMPRT 114 Ali, Carine; INSERM U919, Serine Proteases and Pathophysiology of the Neurovascular Unit, UMR-CNRS 6232 CINAPS, Cyceron, Universite de Caen Basse Normandie Vivien, Denis; INSERM U919, Serine Proteases and Pathophysiology of the Neurovascular Unit, UMR-CNRS 6232 CINAPS, Cyceron, Universite de Caen Basse Normandie
Keywords:	desmotepase, blood brain barrier, tissue-type plasminogen activator, excitotoxicity

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

1
2
3 **PPACK-Desmodus rotundus salivary plasminogen activator (cDSPA α 1) prevents the**
4 **passage of tissue-type plasminogen activator (rt-PA) across the blood-brain barrier and**
5 **neurotoxicity**
6
7
8
9

10
11
12 Benoit D. Roussel^{1,4}, PhD; Yannick Hommet^{1,4}, BSc; Richard Macrez¹, MSc; Torsten
13 Schulz², PhD; Karl-Uwe Petersen², MD; Vincent Berezowski³, PhD; Roméo Cecchelli³, PhD;
14 Carine Ali¹, PhD; Denis Vivien^{1*}, PhD
15
16
17
18
19

20
21
22 ¹ INSERM, INSERM U919, Serine Proteases and Pathophysiology of the neurovascular Unit,
23 CNRS, UMR CNRS 6232 CINAPs “Center for Imaging Neurosciences and Applications to
24 Pathologies”, Cyceron, Caen Cedex, F-14074 France; University of Caen Basse-Normandie,
25 Caen Cedex, F-14074 France.
26
27
28
29

30
31
32 ² PAION Deutschland GmbH, Aachen, Germany
33

34
35 ³ Laboratoire de Physiopathologie de la Barrière Hémato-Encéphalique, EA 2465, IFR114-
36 IMPRT, Université d’Artois, Faculté des Sciences Jean Perrin, France
37
38

39
40 ⁴ These authors contributed equally to this work.
41

42 * **Corresponding author:** Denis Vivien, PhD; INSERM U919, Serine Proteases and
43 Pathophysiology of the neurovascular Unit; UMR CNRS 6232 CINAPs “Center for Imaging
44 Neurosciences and Applications to Pathologies”, Cyceron, Bd H. Becquerel, BP 5229, Caen
45 Cedex, F-14074 France. Tel: +33-2-31-47-01-66 / Fax: +33-2-31-47-02-22 / e-mail:
46 vivien@cyceron.fr
47
48
49
50
51
52

53
54
55 **Acknowledgments:** This work was supported by grants from the INSERM, Université de
56 Caen Basse-Normandie, European Council (FEDER), FP6-project DiMI-LSHB-CT-2005-
57 512146-, Regional Council of Lower Normandy.
58
59
60

1
2
3
4
5
6 The treatment of ischemic stroke remains one of the most challenging areas in
7
8 medicine. To date, the only treatment approved by health authorities is early reperfusion by
9
10 the thrombolytic agent, recombinant tissue-type plasminogen activator (rt-PA)¹. Nevertheless,
11
12 despite benefit from fibrinolysis, several limitations are linked to the use of rt-PA, including a
13
14 clinically relevant risk of haemorrhage¹ and a highly suspected risk of neurotoxicity^{2,3}.
15
16 Desmoteplase (DSPA) is a highly fibrin-specific recombinant plasminogen activator (PA)
17
18 isolated from the saliva of the vampire bat *Desmodus rotundus*⁴. Experimental data suggest
19
20 that DSPA could display several advantages: i) in contrast to rt-PA, DSPA is devoid of pro-
21
22 excitotoxic effects both *in vitro* and *in vivo*^{5,6}; ii) intravenous DSPA significantly reduces the
23
24 passage of co-administered rt-PA across the intact BBB and the attendant aggravation of
25
26 excitotoxic damage⁶. In this context, the aim of the present study was to investigate, using a
27
28 predictive *in vitro* model of BBB⁷ (Fig. 1A), and an *in vivo* model of excitotoxicity, whether
29
30 an inactive form of DSPA, clogged DSPA (cDSPA), could prevent the ability of rt-PA to
31
32 cross the BBB and promote excitotoxic injuries. As previously demonstrated for native
33
34 DSPA⁶, we found that cDSPA did not alter the integrity of the BBB (passage of sucrose)
35
36 alone or in combination with rt-PA (data not shown) and can cross the BBB, despite being
37
38 catalytically inactive (Fig. 1B). Then, the passage of rt-PA alone or in combination with either
39
40 DSPA or cDSPA was analyzed using both fluorogenic proteolytic (spectrozyme®) and
41
42 zymography assays (Fig. 1C and D). Both DSPA and cDSPA reduced the passage of rt-PA by
43
44 around 30% in our *in vitro* BBB model. The effects of intravenous injection of rt-PA and
45
46 cDSPA were studied in a mouse model of excitotoxicity induced by a striatal stereotaxic
47
48 injection of NMDA (Figure 1E). Neither rt-PA nor cDSPA vehicles injected intravenously
49
50 altered the extent of NMDA-induced striatal injury. As previously demonstrated⁶, the
51
52 systemic injection of rt-PA increased the striatal lesion volume seen with NMDA by almost
53
54
55
56
57
58
59
60

1
2
3 60% (from ~10 to ~16 mm³). Interestingly, intravenous co-administration of cDSPA with rt-
4 PA, both at 1 mg/kg, suppressed the ability of exogenous rt-PA to enhance NMDA-induced
5 lesion. No micro-haemorrhages were detected in animals treated with rt-PA and/or clogged-
6 desmoteplase.
7
8
9
10
11

12 Endogenous tPA, released by vascular endothelial cells, is well-known to play a
13 critical role as a thrombolytic enzyme that activates plasminogen to plasmin. Based on this
14 activity, exogenous rt-PA-induced thrombolysis remains the only FDA-approved drug for the
15 treatment of patients in acute ischemic stroke¹. However, t-PA favors intracerebral
16 hemorrhagic transformation, limiting the licensed use of this drug to a 3 hours post-ictus
17 therapeutic window¹. Accordingly, the balance between the beneficial effects of thrombolysis
18 and the injurious cerebral effects of rt-PA may be improved by restricting the access of
19 vascular rt-PA to the brain parenchyma. Previously, we and others have demonstrated both *in*
20 *vitro* and *in vivo* that despite its common ability to cross the intact BBB through a common
21 transport system, in contrast to rt-PA, DSPA, is not pro-neurotoxic⁶. As a further
22 characterization of the transport mechanism, our results show that a proteolytically inactive
23 form of DSPA can restrict rt-PA trans-BBB passage. This is in agreement with the previous
24 demonstration that PAs can cross the BBB independently of their proteolytic
25 activity/domain⁸. We also demonstrate that clogged-desmoteplase prevents exogenous rt-PA-
26 promoted excitotoxicity *in vivo*. This is in agreement with previous demonstrations that DSPA
27 can prevent cerebral damage by rt-PA *in vivo*^{5,6} a feature likely to contribute to the increased
28 tolerance and safety of the clinical use of DSPA in the 3-9 hours time window post-onset of
29 stroke symptoms⁹. Thus, while rt-PA remains the only approved thrombolytic drug to treat
30 stroke patients, the interest in alternatives like DSPA is sustained. As intrinsic fibrinolytic
31 activity of such an antagonist would not be required or might even be counterproductive,
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 generation of a proteolytically inactive version of DSPA able to compete with rt-PA for
4 transport at the BBB might open the way to a safer use of rt-PA as a thrombolytic in stroke.
5
6
7
8
9

10 **Figure 1:** (A) Schematic representation of the in vitro BBB model (B) biot-rt-PA or biot-
11 cDSPA α 1 were applied to the upper endothelial compartment. Two hours later, their activities
12 or presence in the lower side medium were analyzed either by zymography assay or
13 immunoblotting (n=3). (C,D) rt-PA alone or with DSPA or cDSPA was applied to the upper
14 endothelial compartment. Two hours later, rt-PA activity was analyzed in the lower side
15 medium, either by fluorogenic assay (spectrozyme®) (C) or zymography assay (D). Results
16 are normalized to rt-PA (mean values \pm SEM of 3 experiments per group; * $p < 0.05$ Mann-
17 Whitney test). (E) Effects of intravenous injection of tPA alone or in combination with
18 cDSPA (1 mg/kg each) on the extent of neuronal death induced by the striatal administration
19 of NMDA (10 nmol) in mice (mean \pm SD; n=10 per group; *: $p < 0.01$ compared to NMDA
20 alone, \$: $p < 0.01$ compared to NMDA + tPA, Mann-Whitney test).
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

39 **Reagents,** Human recombinant rt-PA (Actilyse®) from Boehringer Ingelheim (Paris, France).
40 DSPA was provided by PAION Deutschland GmbH (Aachen, Germany).
41

42 **Production of cDSPA:** cDSPA was generated by coupling the Phe-Pro-Arg-
43 chloromethylketone modified tripeptide to DSPA.
44
45
46
47

48 **Biotinylation:** Since cDSPA α 1 has no intrinsic enzymatic activity, we had to develop a
49 method of detection for transport studies, based on biotinylation of the compound, allowing
50 performing SDS-PAGE/immunoblotting against biotin (with a streptavidin secondary
51 antibody). We used the FluoReporter®Biotin-XX Protein Labeling kit (Molecular Probes)
52 according to the manufacturer's instructions.
53
54
55
56
57
58
59
60

1
2
3 **Blood-brain barrier (BBB) in vitro experiments:** Transport across the BBB was studied
4 using a previously characterized in vitro model⁷, consisting of a co-culture of endothelial and
5 glial cells and shown to closely mimic the in vivo BBB. rt-PA and/or desmoteplase were
6 added to the upper side of the endothelial cells. Abluminal and luminal media were harvested
7 after the 2 hours transport experiments.
8
9

10
11
12
13
14 **Spectrozyme[®] assays** were performed using a fluorogenic substrate (Spectrozyme XF444,
15 American Diagnostica, Stamford, NJ, USA).
16
17

18
19
20 **Zymography assay** was performed by adding plasminogen (22.5µg) and casein (5mg) to a
21 12.6% SDS polyacrylamide gel.
22
23

24
25 **Striatal excitotoxic lesions:** Mice were anaesthetised with isoflurane and placed in a
26 stereotaxic frame. Body temperature was maintained at $37 \pm 0.5^\circ\text{C}$. The skull was exposed
27 and an injection pipette was stereotaxically implanted in the right striatum according to the
28 atlas of Paxinos & Watson (coordinates 0.5 mm posterior, +/- 2 mm lateral and -3 mm ventral
29 to the bregma). NMDA (10 nmol) was injected in a volume of 0.3 µl. Excitotoxic treatment
30 was followed after 10 minutes by intravenous injection of rt-PA (1 mg/kg), cDSPA (1 mg/kg),
31 rt-PA + cDSPA (1 mg/kg each), tPA vehicle (L-Arg 35 mg/kg, phosphoric acid 10 mg/kg and
32 polysorbate 80 0.2%) or cDSPA vehicle (glycine 4 mg/kg and mannitol 10.64 mg/kg).
33
34
35
36
37
38
39
40
41
42
43

44 **Histological analysis:** For volume analysis, one coronal section (20 µm) out of every 10 was
45 stained with thionine and analysed with an image analyser (Scion Image, Scion Corporation,
46 Frederick, Maryland, USA).
47
48
49

50
51 **Statistical analysis:** For *in vitro* and *in vivo* studies, data were expressed as mean \pm SEM, and
52 statistical analyses were performed using the Kruskal and Wallis test followed by a Mann-
53 Whitney test for inter-group comparisons.
54
55
56
57
58
59
60


1
2
3 **Declaration of commercial interest:** This study was supported in part by research funding
4
5 from PAION Deutschland GmbH.
6
7
8
9

10 **References**

- 11
12
13
14
15 1. NINDS. The National Institute of Neurological Disorders and Stroke rt-PA Stroke Study
16 Group. Tissue plasminogen activator for acute ischemic stroke (1995) The National Institute
17 of Neurological Disorders and Stroke tPA Stroke Study Group. N Engl J Med 333:1581-1587.
18
19
20
21 2. Tsirka SE, Gualandris A, Amaral DG, Strickland S (1995) Excitotoxin-induced neuronal
22 degeneration and seizure are mediated by tissue plasminogen activator. Nature 377:340-344.
23
24
25 3. Nicole O, Docagne F, Ali C, Margail I, Carmeliet P, MacKenzie ET, Vivien D, Buisson A
26 (2001) The proteolytic activity of tissue-plasminogen activator enhances NMDA receptor-
27 mediated signaling. Nat Med 7:59-64.
28
29
30 4. Schleuning WD, Alagon A, Boidol W, Bringmann P, Petri T, Krätzschar J, Haendler B,
31 Langer G, Baldus B, Witt W, Donner P (1992) Plasminogen activators from the saliva of
32 *Desmodus rotundus* (common vampire bat): unique fibrin specificity. Ann N Y Acad Sci
33 4:395-403.
34
35 5. Reddrop C, Moldrich RX, Beart PM, Farso M, Liberatore GT, Howells DW, Petersen KU,
36 Schleuning WD, Medcalf RL (2005) Vampire bat salivary plasminogen activator
37 (desmoteplase) inhibits tissue-type plasminogen activator-induced potentiation of excitotoxic
38 injury. Stroke 36:1241-1246.
39
40
41
42
43 6. López-Atalaya JP, Roussel BD, Ali C, Maubert E, Petersen KU, Berezowski V, Cecchelli
44 R, Orset C, Vivien D (2007) Recombinant *Desmodus rotundus* salivary plasminogen activator
45 crosses the blood-brain barrier through a low-density lipoprotein receptor-related protein-
46 dependent mechanism without exerting neurotoxic effects. Stroke 38:1036-1043.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
7. Cecchelli R, Dehouck B, Descamps L, Fenart L, Buée-Scherrer VV, Duhem C, Lundquist S, Rentfel M, Torpier G, Dehouck MP (1999) In vitro model for evaluating drug transport across the blood-brain barrier. *Adv Drug Deliv Rev* 36:165-178.
8. Benchenane K, Berezowski V, Ali C, Fernández-Monreal M, López-Atalaya JP, Brillault J, Chuquet J, Nouvelot A, MacKenzie ET, Bu G, Cecchelli R, Touzani O, Vivien D (2005) Tissue-type plasminogen activator crosses the intact blood-brain barrier by low-density lipoprotein receptor-related protein-mediated transcytosis. *Circulation* 111:2241-2249.
9. Hacke W, Albers G, Al-Rawi Y, Bogousslavsky J, Davalos A, Eliasziw M, Fischer M, Furlan A, Kaste M, Lees KR, Soehngen M, Warach S; DIAS Study Group (2005) The Desmoteplase in Acute Ischemic Stroke Trial (DIAS): a phase II MRI-based 9-hour window acute stroke thrombolysis trial with intravenous desmoteplase. *Stroke* 36:66-73.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49



- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Based on its fibrinolytic activity, tissue type plasminogen activator (tPA) is the only treatment approved so far by the authorities, to treat ischemic stroke patients (NINDS, 1995).
 - However, exogenous tPA can display deleterious effects on components of the neurovascular unit, including promotion of edema and blood-brain barrier (BBB) leakage (Aoki et al., Stroke, 2002, Yepes et al., J.Clin. Invest., 2003), which probably minimise its overall benefit.
 - Moreover, tPA was shown to have the ability to cross both the healthy and injured BBB (Benchenane et al., Circulation, 2005), thus adding to the pro-neurotoxic effect of endogenously produced parenchymal tPA (Tsirka et al., Nature, 1995; Nicole et al., Nat. Med., 2001).
 - Desmoteplase (DSPA) is a new thrombolytic agent derived from bat salivary glands (Hacke et al., Stroke, 2005), very close to tPA in terms of structural determinants, but previously demonstrated to :
 - cross the BBB as tPA does (Lopez-Atalaya et al., Stroke, 2007)
 - be devoid of pro-neurotoxic effect (Reddrop et al., Stroke, 2005; Lopez-Atalaya et al., J Cereb Blood Flow Metab, 2008)
 - Although the use of DSPA to treat stroke patients is debated (see DIAS 2 data), we provide here the demonstration that :
 - a proteolytically inactive form of DSPA can prevent the passage of exogenous tPA across the BBB and its subsequent neurotoxic activity, thus demonstrating the therapeutic potential of competitive inhibitors of tPA's transport to the brain for stroke patients.