



HAL
open science

Argan oil prevents down-regulation induced by endotoxin on liver fatty acid oxidation and gluconeogenesis and on peroxisome proliferator-activated receptor gamma coactivator-1 alpha, (PGC-1alpha), peroxisome proliferator-activated receptor gamma (PPARgamma) and estrogen related receptor alpha (ERRalpha)

Riad El Kebbaj, Pierre Andreoletti, Hammam L El Hajj, Youssef El Kharrassi, Joseph Vamecq, Stéphane I Mandard, Fatima-Ezzahra I Saih, Norbert I Latruffe, M'hammed Saïd El Kebbaj, Gérard I Lizard, et al.

► **To cite this version:**

Riad El Kebbaj, Pierre Andreoletti, Hammam L El Hajj, Youssef El Kharrassi, Joseph Vamecq, et al.. Argan oil prevents down-regulation induced by endotoxin on liver fatty acid oxidation and gluconeogenesis and on peroxisome proliferator-activated receptor gamma coactivator-1 alpha, (PGC-1alpha), peroxisome proliferator-activated receptor gamma (PPARgamma) and estrogen related receptor alpha (ERRalpha). *Biochimie Open*, 2015, 1, pp.51-59. 10.1016/j.biopen.2015.10.002 . inserm-01534672

HAL Id: inserm-01534672

<https://inserm.hal.science/inserm-01534672>

Submitted on 7 Jun 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Research paper

Argan oil prevents down-regulation induced by endotoxin on liver fatty acid oxidation and gluconeogenesis and on peroxisome proliferator-activated receptor gamma coactivator-1 α , (PGC-1 α), peroxisome proliferator-activated receptor α (PPAR α) and estrogen related receptor α (ERR α)

Riad El Kebbjaj^{a,b,c}, Pierre Andreoletti^a, Hammam I. El Hajj^a, Youssef El Kharrassi^{a,b}, Joseph Vamecq^d, Stéphane Mandard^e, Fatima-Ezzahra Saih^{a,b}, Norbert Latruffe^a, M'Hammed Saïd El Kebbjaj^f, Gérard Lizard^a, Boubker Nasser^b, Mustapha Cherkaoui-Malki^{a,*}

^a Univ. Bourgogne-Franche Comté, Laboratoire BioPeroXIL (Biochimie du Peroxysome, Inflammation et Métabolisme Lipidique), EA 7270, 21000 Dijon, France

^b Laboratoire de Biochimie et Neurosciences, Faculté des Sciences et Techniques, Université Hassan I, BP 577, 26 000 Serrat, Morocco

^c Laboratoire des Sciences et Technologies de la Santé, Institut supérieur des sciences de la santé Université Hassan I, Route de Casablanca, 14 BP 539, 26 000 Serrat, Morocco

^d INSERM and HMNO, CBP, CHRU Lille, 59037 Lille and RADEME EA 7364, Faculté de Médecine, Université de Lille 2, 59045 Lille, France

^e Lipness Team, INSERM, Research Center UMR866 and LabEx LipSTIC, Université de Bourgogne-Franche Comté, Dijon, France

^f Laboratoire de recherche sur les lipoprotéines et l'Athérosclérose, Faculté des Sciences Ben M'sik, Avenue Cdt Driss El Harti, BP 7955, Université Hassan II-Mohammedia-Casablanca, Morocco

Received 15 May 2015; accepted 20 October 2015

Available online 31 October 2015

Abstract

In patients with sepsis, liver metabolism and its capacity to provide other organs with energetic substrates are impaired. This and many other pathophysiological changes seen in human patients are reproduced in mice injected with purified endotoxin (lipopolysaccharide, LPS). In the present study, down-regulation of genes involved in hepatic fatty acid oxidation (FAOx) and gluconeogenesis in mice exposed to LPS was challenged by nutritional intervention with Argan oil. Mice given a standard chow supplemented or not with either 6% (w/w) Argan oil (AO) or 6% (w/w) olive oil (OO) prior to exposure to LPS were explored for liver gene expressions assessed by mRNA transcript levels and/or enzyme activities. AO (or OO) food supplementation reveals that, in LPS-treated mice, hepatic expression of genes involved in FAOx and gluconeogenesis was preserved. This preventive protection might be related to the recovery of the gene expressions of nuclear receptors peroxisome proliferator-activated receptor α (PPAR α) and estrogen related receptor α (ERR α) and their coactivator peroxisome proliferator-activated receptor gamma coactivator-1 α , (PGC-1 α). These preventive mechanisms conveyed by AO against LPS-induced metabolic dysregulation might add new therapeutic potentialities in the management of human sepsis.

© 2015 The Authors. Published by Elsevier B.V. on behalf of Société Française de Biochimie et Biologie Moléculaire (SFBBM).

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: Argan oil; Beta-oxidation; Coactivator; Gluconeogenesis; Nuclear receptor.

Abbreviations: ACADS, acyl CoA dehydrogenase short-chain; ACADM, acyl CoA dehydrogenase medium-chain; ACADL, acyl CoA dehydrogenase long-chain; AO, Argan oil; ACOX1, acyl-CoA oxidase 1; ERR α , estrogen related receptor α ; G6PH, glucose-6-phosphatase; Glut2, glucose transporter 2; Glut4, glucose transporter 4; HNF-4 α , hepatic nuclear factor-4 α ; OO, olive oil; LPS, lipopolysaccharide; PGC-1 α , peroxisome proliferator-

activated receptor γ coactivator-1 α ; PEPCK, phosphoenolpyruvate carboxykinase; PPAR α , peroxisome proliferator-activated receptor α .

* Corresponding author. Laboratoire BioPeroXIL EA7270, Univ. Bourgogne-Franche Comté, Faculté des Sciences Gabriel, 6 Bd Gabriel, 21000 Dijon, France. Tel.: +33 380396205.

E-mail address: malki@u-bourgogne.fr (M. Cherkaoui-Malki).

<http://dx.doi.org/10.1016/j.biopen.2015.10.002>

2214-0085/© 2015 The Authors. Published by Elsevier B.V. on behalf of Société Française de Biochimie et Biologie Moléculaire (SFBBM). This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Bacterial infection is a common cause of sepsis, a pathological state inducing a severe organ dysfunction and a high mortality rate, and requiring intensive care [1–3]. This acute syndrome is associated with systemic inflammation and disturbed metabolism [4,5]. During bacterial infection, release in the host of endotoxins (lipopolysaccharides, LPS) from gram-negative bacteria membrane generates a potent inflammatory cytokine response and severely impairs lipid metabolism, inducing reduced serum high density lipoprotein (HDL), increased plasma free fatty acids and triglycerides levels [3]. These metabolic changes are mainly accounted for by enhanced hepatic triglyceride synthesis and adipose tissue lipolysis combined with a drop in fatty acid oxidation (FAOx) in several tissues including heart, kidney, liver and skeletal muscle [3,6–10]. The downregulation of FAOx by LPS is correlated with decreased expressions of the nuclear receptor Peroxisome Proliferator-Activated Receptor (PPAR) α and its coactivator PPAR γ Coactivator (PGC)-1 α , which physiologically work in concert to regulate FAOx-related gene expressions [11,12]. In this respect, ligand-dependent activation of the nuclear receptor PPAR α prompts its heterodimerization with Retinoid X Receptor (RXR) α [13,14]. The PPAR α /RXR α complex binds to PPAR α -response elements (PPRE) of target genes which may code for mitochondrial and peroxisomal enzymes involved in fatty acid β -oxidation pathways such as carnitine palmitoyl transferase 1 (CPT1a and CPT1b), short-, medium-, long- and very long-chain acyl CoA dehydrogenases (ACADS, ACADM, ACADL and ACADVL) [15–19], acyl-CoA oxidase 1 (ACOX1) [17,20,21] and other proteins [12,22]. On the other hand, Lipin-1, a phosphatidate phosphatase, has arisen as an additional transcriptional co-regulator of PPAR α -PGC-1 α -directed gene expression [23]. Its interaction with PPAR α -PGC-1 α complex promotes the induction of FAOx genes [24]. Beside PPAR α , estrogen related receptor (ERR) α or (ESRR α), an orphan nuclear receptor, has been also shown to regulate energy metabolism gene expression [25,26], particularly genes involved in FAOx [27,28]. This transcriptional regulation involves interaction with PGC-1 α coactivator through a protein motif specifically dedicated to ERR α [29,30]. In liver, another interaction of PGC-1 α is also observed for hepatic nuclear factor-4 α (HNF-4 α) to control genes coding gluconeogenesis proteins (phosphoenolpyruvate carboxykinase (PEPCK), glucose-6-phosphatase (G6PH)) and glucose transporter 4 (Glut4) [31].

More than 20% of patients with sepsis develop liver dysfunction [1] and hence dysregulation of hepatic metabolism and reduced energy supply for other organs. In mouse models of sepsis, injection of purified LPS triggers many pathophysiological changes resembling those described in human patients [32]. Though down-regulatory mechanisms by which LPS impacts FAOx have been extensively studied, little attention has been actually paid to mechanisms capable of preserving normal FAOx and inflammation status. Interestingly, supplementation of parenteral nutrition with fish oil to patients, during the postoperative period, revealed lowest levels of circulating inflammatory mediators [33–35]. Accordingly, polyunsaturated fatty acid-rich diet

has been reported to reduce acute inflammation and to promote anti-inflammatory process in mice [36]. Therefore, lipid nutritional support might help the prevention of not only inflammatory damages but also disrupted lipid homeostasis.

Argan edible oil (AO) is obtained by cold-pressure of roasted kernels from *Argania spinosa* [L.] Skeels, a singular Mediterranean species growing in the southwestern region of Morocco. Argan oil is used as a traditional food ingredient in the ‘Amazigh diet’, bringing almost 25% of total diet fat intake to indigenous consumers [37]. Accordingly, early clinical studies on Argan oil reported a decrease in plasma low density lipoprotein-cholesterol (LDL-cholesterol) and lipid hydroperoxides along with a rise in plasma tocopherol concentration [38]. Health benefits of this delectable virgin oil have been highlighted by several studies documenting its cardiovascular protective potential including hypocholesterolemic and hypotriglyceridemic properties in consumer populations [39–41]. AO has been also shown to reduce circulating LDL-cholesterol and ApoB and, in AO consumers, to increase HDL and ApoAI [40,41] whereas in human macrophages it increases HDL-mediated cholesterol efflux and reduces LDL-lipid peroxidation [38,39].

Therefore, in an attempt to test our hypothesis regarding the preventive effects of Argan oil against LPS-induced FAOx downregulation, mice pretreated with AO were subsequently injured by LPS to determine whether an experimental support may be or not given to this working hypothesis. The effects of AO against sepsis-associated liver hyperlipidemia are compared to those of olive oil (OO), a more usual ingredient in Mediterranean diets. We report here that, in fact, AO-enriched diet prevents LPS-associated hyperlipidemic effect through the induction of the hepatic expressions of PPAR α , ERR α and their coactivator PGC-1 α along with the up-regulation of their mitochondrial (ACADS, ACADM, ACADVL) and peroxisomal (ACOX1) target genes.

2. Material and methods

2.1. Argan oil treatment

Swiss OF1 mice (12–16 week-old) were obtained from IFFA CREDO (Casablanca). They were acclimatized in the laboratory for 10 days at 22 ± 2 °C with standard chow and water *ad libitum*. Animal studies were conducted in accordance with the protocols of Animal Use and Care of the University of Hassan 1st, Settat, Morocco. The virgin Argan oil used in this work was obtained from the Aklim area in the northeast of Morocco. Six groups of mice (5 mice/group) received during 25 days: a standard chow (2 groups, control); a standard chow supplemented with 6% (w/w) of Argan oil (2 groups, AO) or a standard chow supplemented with 6% (w/w) of olive oil (2 groups, OO). Oils were included in the diets by direct mixing with the standard animal chow. Sixteen hours before euthanasia and during the fed state, one group from control (+LPS), AO (AO + LPS) and OO (OO + LPS) respectively received (5 mg/kg) intraperitoneal injections of 100 μ g of *Escherichia coli* 0111:B4 LPS (Sigma) resuspended in phosphate-buffered saline (PBS) or an equal volume of PBS alone.

2.2. Composition of oils

Both Argan and olive oils (AO and OO) contain mono and polyunsaturated fatty acids. However, Argan oil has 35% of C18:2n-6 and 45% of C18:1n-9 while olive oil shows only 6% of C18:2n-6 and more than 75% of C18:1n-9, leading to a higher unsaturation index of AO (120.4) versus OO (108.3) [42].

2.3. Quantitative PCR analysis

Total RNA from liver was extracted using the RNeasy Mini kit (Qiagen) following the manufacturer's instructions. cDNA was generated by reverse transcription using Moloney Murine Leukemia Virus Reverse Transcriptase (Promega) according to the manufacturer's protocol and analyzed by quantitative PCR using the GoTaq[®] qPCR Master Mix (Promega), and a StepOnePlus Real-Time PCR System (Applied Biosystem). The primer sequences were chosen using the Beacon Designer Software (Bio-Rad). Oligonucleotide sequences are shown in the [Supplementary Table 1](#). PCR reactions were carried out in duplicate in a final volume of 12.5 μ L containing 6.25 μ L of MESA Green qPCR Master mix (Eurogentec), 2.5 μ L of cDNA and forward and reverse primers at 300 nM. The PCR enzyme (*Taq* DNA polymerase) was heat-activated at 95 °C for 10 min, and the DNA was amplified for 40 cycles at 95 °C for 15 s, 60 °C for 30 s, and 72 °C for 30 s, followed by a melting curve analysis to control the absence of nonspecific products. For each transcript, the amplification efficiency was determined by the slope of the standard curve generated from two fold serial dilutions of cDNA. Gene expression was quantified using cycle to threshold (Ct) values and normalized by the reference gene, *36B4* encoding the acidic ribosomal phosphoprotein P0. To this end, the quantitative gene expression was determined according to $2^{-\Delta\Delta C_t}$ with $\Delta C_t = (C_t \text{ of the gene studied}) - (C_t \text{ of the } 36B4 \text{ gene})$.

2.4. Enzymatic activity measurements

One hundred mg of liver tissue were homogenized by a Potter-Elvehjem homogenizer in 0.2 ml of a buffer containing 250 mmol/L sucrose, 20 mmol/L Tris-HCl pH 7.5 and 2 mmol/L EDTA. After centrifugation at 600 g for 5 min at 4 °C, the supernatant was collected and stored at -80 °C until use. Peroxisomal acyl-CoA oxidase (ACOX1) activity was measured by the fluorometric assay using palmitoyl-CoA as a substrate as described previously [43]. Catalase activity was monitored at 240 nm as described elsewhere [44]. Mitochondrial acyl-CoA dehydrogenases activities were followed at 600 nm on acyl-CoAs of different chain lengths (for experimental details, see legends to figures) [45].

2.5. Statistical analysis

Statistical analyses to compare two experimental groups were performed with, an unpaired, two-tailed, Student-t test (Excel software) for calculating the probability values and data were considered statistically different at a P-value of 0.05 or less.

3. Results

3.1. Body weight under experimental diets

[Fig. S1](#) shows the time-course of body weights evolution during the nutritional intervention. Each group of mice was weighted at four different time points during the 3 weeks that precede the LPS administration; no significant differences in body weight have been found between the different experimental groups.

3.2. Argan oil modulates the expression of PPAR α , PGC-1 α and related target genes

Interestingly, polyunsaturated fatty acids are known to be activators of PPAR α , a nuclear receptor which governs lipid metabolism and fatty acid oxidation [46]. Here we report ([Fig. 1A](#)) that mice treated with AO or OO showed a significant increase in hepatic PPAR α mRNA, while the mRNA expression of ERR α displayed no significant variation ([Fig. 1B](#)). Intriguingly, the expression of PGC-1 α mRNA was specifically down regulated by AO ([Fig. 1C](#)). However, the expression of PPAR α target genes (i.e. *ACOX1* and *ACADM*) was clearly induced by both AO and OO, while *ACADS* mRNA level was only increased by AO ([Fig. 1E, F](#)). Regarding the mitochondrial fatty acid beta-oxidation activities, only short (C4:0)-, long (C12:0)- and very long (C16:0)-acyl-CoA mitochondrial dehydrogenases were increased but not the medium (C:8) acyl-CoA dehydrogenase activity ([Fig. 2C, D](#)). By contrast to mitochondria, peroxisomal palmitoyl-CoA oxidase was not changed and only peroxisomal catalase activity was induced by AO or OO ([Fig. 2A](#)). These results support experimentally that AO and OO up-regulate specifically mitochondrial fatty acid β -oxidation (except for the medium chain) and peroxisomal catalase activities ([Fig. 2B](#)).

3.3. LPS induces selective changes in mitochondrial and peroxisomal FAOx gene expression

Consistent with previous studies [47,48], treatment with LPS strongly decreased the expression of nuclear receptors, PPAR α and ERR α , mRNA levels in mouse liver ([Fig. 1A, B](#)). LPS also decreased the hepatic expression of mRNA level of PGC-1 α . However in these conditions, lipin-1, which may interact with PGC-1 α [24], showed enhanced mRNA levels ([Fig. 1D](#)). Measurements of the expression of mitochondrial and peroxisomal FAOx gene products at two stages (mRNA levels and enzyme activities) and 16 h after LPS injection showed a selective decrease of *ACADS* without changes in *ACOX1*, *ACADM* and *ACADVL* mRNA expressions ([Fig. 1E, F](#)). In addition, LPS treatment led to a reduction of mitochondrial *ACADS* enzyme activity ([Fig. 2C](#)) and also, as further discussed, of peroxisomal *ACOX1* activity ([Fig. 2A](#)), while the activity of peroxisomal catalase was enhanced ([Fig. 2B](#)).




Fig. 1. Argan oil preserves hepatic mRNA expressions of nuclear receptors PPAR α and ERR α , coactivator PGC-1 α and target genes during exposure of mice to LPS. Real-time PCR was used to quantify the hepatic mRNA levels of PPAR α (A) and ERR α (B), coactivators PGC-1 α (C) and lipin-1 (D) their target genes: *Acox1* (E), *Acads*, *Acadm* and *Acadvl* (F). All real-time PCR reactions were performed in duplicate. All values are means \pm SEM (n = 5/group) and are normalized to control mice. Symbols (*, # and +) correspond to a statistical significance of higher mean signal intensity, (p < 0.01 for *** and +++, p < 0.02 for ** and ++, p < 0.05 for * and #), compared with the control (*) or with the LPS-treated mice (+). Mice received for 25 days a standard chow (control); a standard chow supplemented with 6% (w/w) of Argan oil (AO) or a standard chow supplemented with 6% (w/w) of olive oil (OO). Sixteen hours before euthanasia, one group from control (+LPS), AO (AO + LPS) and OO (OO + LPS) respectively received intraperitoneal injection of 100 μ g LPS.

3.4. Argan oil protects against the drops induced by LPS in hepatic expression of PPAR α , ERR α and coactivator PGC-1 α

To evaluate the direct therapeutic benefit of Argan oil (AO), mice were pretreated with AO for 24 days before LPS injection taking place 16 h prior to euthanasia. The effects of AO were compared to those of OO, and as illustrated by Fig. 1A, preserved levels of PPAR α mRNA were observed in AO + LPS-treated mouse livers, being comparable to protection observed in livers from OO + LPS-treated mice (Fig. 1A). By contrast, only AO + LPS induced substantial significant increases in liver expression of ERR α and PGC-1 α mRNA when compared to LPS alone, while OO + LPS induced only a modest significant increase in PGC-1 α mRNA level (Fig. 1B and C). At the opposite, pretreatment with AO + LPS showed an equal increase in levels of Lipin-1 mRNA as in LPS group, revealing no effect of AO. However, combined OO pretreatment and LPS injury reduced significantly the expression of Lipin-1 mRNA (Fig. 1D). Evaluation of mRNA levels of PPAR α target genes involved in mitochondrial and peroxisomal FAOx (Fig. 1E, F) showed that

LPS provoked only a decrease in ACADS mRNA, being without significant effects on mRNA levels of ACADM, ACADVL and ACOX1. Measurements of FAOx enzyme activities revealed that the selective reduction by LPS treatment of ACOX1 and ACADS (C4:0) activities were prevented by AO or OO (AO + LPS or OO + LPS), while for the other dehydrogenase activities, no effects of the oils were observed except for C16:0 with AO + LPS versus LPS (Fig. 2A, C and D). The enhanced activity of peroxisomal catalase, after LPS treatment, was preserved by OO pretreatment (OO + LPS) and to a lesser extent by AO (AO + LPS) (Fig. 2B).

3.5. Argan oil preserves hepatic gluconeogenesis gene expressions during LPS-induced liver dysfunction

HNF4 α is a critical nuclear receptor of PGC-1 α -mediated gluconeogenesis and controls the expression of gluconeogenic genes (PEPCK, G6PH) [31]. Here we showed that AO and OO treatments had no effect on the mRNA level of HNF4 α , and 16 h after LPS administration, there was still no change in the expression of HNF4 α (Fig. 3A), though the expression of its known




Fig. 2. Argan oil protects hepatic mitochondrial and peroxisomal fatty acid oxidation during exposure of mice to LPS. The specific activities of mitochondrial acyl-CoA dehydrogenases (SCAD, MCAD, LCAD and VLCAD) and peroxisomal enzymes (ACOX1 and Catalase) were measured in liver homogenates as described in “Material and Methods” section. All values are means \pm SEM ($n = 5$ /group). Symbols (*, # and +) correspond to a statistical significance of higher mean signal intensity ($p < 0.01$ for ***, □□□ and ###; $p < 0.02$ for **, ## and ++; $p < 0.05$ for *, □ and #), compared with the control mice (*) or with the LPS-treated mice (+). Mice received for 25 days a standard chow (control); a standard chow supplemented with 6% (w/w) of Argan oil (AO) or a standard chow supplemented with 6% (w/w) of olive oil (OO). Sixteen hours before euthanasia, one group from control (+LPS), AO (AO + LPS) and OO (OO + LPS) respectively received intraperitoneal injection of 100 μ g LPS.

target genes were induced to different extents by AO and OO treatments. Indeed, AO up-regulated the expression of *PEPCK*, *G6PH* and *Glut2* mRNA, while OO, *PEPCK* and *Glut4* mRNA levels (Fig. 3B). Finally, administration of LPS deeply reduced the mRNA expression of these four target genes (Fig. 3B). Interestingly, AO + LPS-treated mice exhibited preservation of liver gluconeogenic gene expressions, particularly *PEPCK*, *G6PH* and *Glut4*, which were kept quasi-normal. However, the expression of *Glut2*, the highly expressed glucose transporter in liver, was more decreased in AO + LPS-treated compared to LPS-treated mice (Fig. 3B). OO + LPS-treated mice exhibited also prevention towards LPS-downregulation of liver gluconeogenic genes and no protection for *Glut4* was induced by OO (Fig. 3B).

4. Discussion

The present work provides evidence that mice fed with AO have enhanced expressions of several hepatic FAOx and gluconeogenesis transcripts and this AO-mediated upregulation persists during endotoxic LPS shock. This protective effect appears to associate coregulations of hepatic nuclear receptors PPAR α , ERR α and HNF-4 α and their coactivator PGC-1 α [31]. In addition, AO seems to have specific effects on the activities of mitochondrial acyl-CoA dehydrogenases and peroxisomal catalase.

4.1. AO and OO vs control diets

The body weight of mice fed dietary Argan oil or olive oil did not show any significant difference in comparison to the

body weight found in mice fed the control diet. Although data on mice fed Argan oil are absent in the literature, 10% diet supplementation by AO or OO in rat during 4 weeks also showed no significant differences [49].

In the absence of LPS, AO increases mRNA expression of PPAR α and of its mitochondrial (*ACADS*, *ACADM*) and peroxisomal (*ACOX1*) target genes. Regarding OO treatment, we obtained similar results as with AO for *ACADM* and *ACOX1* mRNA. Even if these oils have different fatty acid compositions, AO and OO induced almost a similar induction of PPAR α and its target genes. As mentioned above, OO contains mainly about 70% of oleic acid and only 6–9% of linoleic acid, while AO harbors 35% of linoleic acid and 45% of oleic acid, indicating that AO is richer in polyunsaturated fatty acids [42]. FAOx induction by OO has been shown to be dependent on PPAR α , since induction by OO or fish oil of hepatic *ACOX1* mRNA is abrogated in *Ppar α* null mice [50]. In this respect, several fatty acids and their polyunsaturated derivatives have been shown to activate responsive element of PPAR α target genes and the generation of *Ppar α* null mice established that PPAR α coordinates transcriptional activation of the genes coding for proteins catalyzing FAOx pathways [14,51–53]. Furthermore, at the energetic level fatty acids are more essential than glucose to the adaptation-phase responses in acute or chronic systemic inflammatory diseases [54]. Thus, it will be of interest to compare in the future the potential preventive effect of supplementing AO to curative properties of its parenteral administration during a septic shock.

On the other hand, though AO or OO increases mRNA expression of mitochondrial acyl-CoA dehydrogenases and




Fig. 3. Argan oil maintains hepatic gluconeogenesis during exposure of mice to LPS- Real-time PCR was used to quantify the hepatic mRNA levels of HNF-4 α (A) and PEPCK, G6PH and Glut4 (B). All real-time PCR reactions were performed in duplicate. All values are means \pm SEM ($n = 5$ /group) and are normalized to control mice. Symbols (*, §, £ and a) correspond to a statistical significance of higher mean signal intensity, ($p < 0.01$ for ***, £££ and §§§; $p < 0.02$ for **, §§ and b; $p < 0.05$ for *, £ and §), compared with the control mice (*, §, £) or with the LPS-treated mice (a and b). Mice received for 25 days a standard chow (control); a standard chow supplemented with 6% (w/w) of Argan oil (AO) or a standard chow supplemented with 6% (w/w) of olive oil (OO). Sixteen hours before euthanasia, one group from control (+LPS), AO (AO + LPS) and OO (OO + LPS) respectively received intraperitoneal injection of 100 μ g LPS.

peroxisomal ACOX1, only mitochondrial oxidation of C4:0, C12:0 and C16:0 acyl-CoA esters and not peroxisomal palmitoyl-CoA oxidase activity were increased. Acyl-CoA oxidase is long known to bind weakly its flavine adenine dinucleotide (FAD) [50,55] and this might contribute to intraperoxisomal dissociation of holoenzyme into FAD and apoenzyme (less stable than holoenzyme). Mitochondrial matrix contains a FAD synthetase [56] and, therefore, might better secure protein need in FAD, and hence a better stability of acyl-CoA dehydrogenases by favoring holoenzyme *vs* apoenzyme forms. In addition, absence of the induction of mitochondrial octanoyl-CoA (C8:0) dehydrogenase activity by different treatments may be related to the process of fatty acid degradations in mitochondria and peroxisomes respectively [53]. Thus, the already known incomplete chain shortening of fatty acyl-CoA in peroxisomes would participate to the octanoyl-CoA export for replenishment of the mitochondrial pool [53].

By contrast to OO, gene expressions of both PGC-1 α and Lipin-1 were down regulated by AO. Such effects might explain the observation of Berrougui et al. [57] that AO diminished both LDL and TG according to the key role of Lipin-1 in the assembly and secretion of hepatic very low-density lipoprotein and the increase of TG synthesis as well [58].

4.2. LPS + AO and LPS + OO vs control + LPS diets

In rodents, previous studies have shown that LPS (after a single bolus) induced similar cytokines profiles in either fed or 48 h-fasted rats [59]. Accordingly, given that 24 h-fasting per se significantly enhances hepatic PPAR α mRNA expression and activity [60] and due to its potent anorexigenic effect, LPS has been injected during the fed state [61].

For the first time, AO is shown to enhance gene expressions of hepatic FAOx and gluconeogenesis in a way persisting during endotoxic LPS shock. This protective mechanism appears to involve coregulation by PGC-1 α of hepatic nuclear receptors PPAR α , ERR α and HNF-4 α [27]. The present work also reports that AO and OO increase liver expression of gluconeogenic genes. Hepatic gluconeogenesis is regulated by PGC-1 α through the coactivation of HNF-4 α [31]. Prevention of the LPS-associated downregulation of gluconeogenic genes is better accounted for by PGC-1 α than by HNF-4 α without ruling out lipin-1 activation. Under Lipin-1 RNAi, PEPCK and G6PH are indeed down regulated [62]. Dysregulation of lipid metabolism by LPS injection is characterized by a dramatic decrease in mitochondrial fatty acid oxidative enzymes [63] and hence fatty acid oxidation (FAOx) in several tissues, including liver

[3,6–10]. Proposed underlying mechanisms include a reduction in PPAR α and its coactivator PGC-1 α expressions and hence in mRNA levels of FAOx genes [3,47,48]. Additionally, we showed a strong decrease of ERR α mRNA and increase in liver Lipin-1 transcripts after LPS treatment. Interestingly, both ERR α and Lipin-1 play key roles in the expression of FAOx genes [48,64]. LPS decreases Lipin-1 mRNA in mouse adipose tissue but not skeletal muscle [64] and ERR α in liver, heart, and kidney of mice markedly during the LPS-induced acute-response phase [48]. Intriguingly, AO supplementation in LPS treated mice here has no effect on the level of Lipin-1 mRNA, in contrast to OO supplementation which prevents enhancement of Lipin-1 mRNA by LPS. Such differential effect between AO and OO is similar to what we obtained for PGC-1 α transcripts. Knowing that Lipin-1 RNAi has been shown to mediate reduction of PGC-1 α mRNA [62], we could suggest that increase in Lipin-1 may participate to the preservation of PGC-1 α mRNA by AO. Our results corroborate and extend previous studies demonstrating that suppression by LPS of FAOx addresses, particularly in liver, peroxisomal ACOX1 and ACADS genes. Such LPS suppressive effect is prevented by nutritional supplementation with AO, which also preserves the gene expressions of ERR α , PPAR α and their coactivator PGC-1 α .

LPS treatment exhibited an opposite effect on peroxisomal beta-oxidation and antioxidant activities, which were decreased for ACOX1 and increased for catalase respectively. This may be related to the fact that ACOX1 is an H₂O₂-generating enzyme, while catalase is an H₂O₂-degrading enzyme [52,53]. The LPS-associated generation of reactive oxygen species is believed to play a key role during the pathogenesis of sepsis [65]. Accordingly, the inhibition of catalase was associated with the progression of LPS/D-galactosamine-induced fulminant liver injury [66]. Thus, the preservation of catalase activity by AO may have a protective effect against the exacerbation of liver injury during LPS-induced endotoxemia.

5. Conclusion

In conclusion, the present work showed that Argan oil protects against the decreased expression of genes involved in hepatic FAOx and gluconeogenesis usually observed during the acute response phase associated with LPS administration. This preventive protection might be related to the recovery of the gene expressions of nuclear receptors PPAR α and ERR α as well as of their coactivator PGC-1 α . Such recovery may explain the preservations of mitochondrial and peroxisomal enzymatic activities in parallel to an improvement of gluconeogenic gene expressions.

Conflict of interests

The authors have declared no conflict of interest.

Acknowledgments

This work was supported by the Action Intégrée of the Comité Mixte Inter-universitaire Franco-Marocain

(CMIFM, AIMA/14/310, CampusFrance) from the PHC Volubilis/Toubkal program (No30293PA), Ministère des Affaires Étrangères, The Centre National Pour la Recherche Scientifique et Technique (CNRS)- Morocco, the Conseil Régional de Bourgogne (PARI 2012:A324; PARI2013:B135; CP S0009) and the Ministère de l'enseignement et de la Recherche (crédits récurrents).

Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.biopen.2015.10.002>.

References

- [1] J.L. Vincent, Y. Sakr, C.L. Sprung, V.M. Ranieri, K. Reinhart, H. Gerlach, R. Moreno, J. Carlet, J.R. Le Gall, D. Payen, Sepsis in European intensive care units: results of the SOAP study, *Crit. Care Med.* 34 (2006) 344–353.
- [2] M. Maeder, T. Fehr, H. Rickli, P. Ammann, Sepsis-associated myocardial dysfunction: diagnostic and prognostic impact of cardiac troponins and natriuretic peptides, *Chest* 129 (2006) 1349–1366.
- [3] W. Khovidhunkit, M.S. Kim, R.A. Memon, J.K. Shigenaga, A.H. Moser, K.R. Feingold, C. Grunfeld, Effects of infection and inflammation on lipid and lipoprotein metabolism: mechanisms and consequences to the host, *J. Lipid Res.* 45 (2004) 1169–1196.
- [4] J. Cohen, The immunopathogenesis of sepsis, *Nature* 420 (2002) 885–891.
- [5] S. Sriskandan, D.M. Altmann, The immunology of sepsis, *J. Pathol.* 214 (2008) 211–223.
- [6] M.S. Liu, J.J. Spitzer, In vitro effects of *E. coli* endotoxin on fatty acid and lactate oxidation in canine myocardium, *Circ. Shock* 4 (1977) 181–190.
- [7] X. Wang, R.D. Evans, Effect of endotoxin and platelet-activating factor on lipid oxidation in the rat heart, *J. Mol. Cell. Cardiol.* 29 (1997) 1915–1926.
- [8] A.C. Johnson, A. Stahl, R.A. Zager, Triglyceride accumulation in injured renal tubular cells: alterations in both synthetic and catabolic pathways, *Kidney Int.* 67 (2005) 2196–2209.
- [9] R.A. Zager, A.C. Johnson, S.Y. Hanson, Renal tubular triglyceride accumulation following endotoxic, toxic, and ischemic injury, *Kidney Int.* 67 (2005) 111–121.
- [10] K.R. Feingold, I. Stapanian, R.A. Memon, A.H. Moser, J.K. Shigenaga, W. Doerrler, C.A. Dinarello, C. Grunfeld, Endotoxin rapidly induces changes in lipid metabolism that produce hypertriglyceridemia: low doses stimulate hepatic triglyceride production while high doses inhibit clearance, *J. Lipid Res.* 33 (1992) 1765–1776.
- [11] C. Handschin, B.M. Spiegelman, Peroxisome proliferator-activated receptor gamma coactivator 1 coactivators, energy homeostasis, and metabolism, *Endocr. Rev.* 27 (2006) 728–735.
- [12] M. Rakhshandehroo, B. Knoch, M. Muller, S. Kersten, Peroxisome proliferator-activated receptor alpha target genes, *PPAR Res.* 2010 (2010) 20–39.
- [13] A. Vluggens, P. Andreoletti, N. Viswakarma, Y. Jia, K. Matsumoto, W. Kulik, M. Khan, J. Huang, D. Guo, S. Yu, J. Sarkar, I. Singh, M.S. Rao, R.J. Wanders, J.K. Reddy, M. Cherkaoui-Malki, Reversal of mouse Acyl-CoA oxidase 1 (ACOX1) null phenotype by human ACOX1b isoform [corrected], *Lab. Invest.* 90 (2010) 696–708.
- [14] A. Vluggens, J.K. Reddy, Nuclear receptors and transcription factors in the development of fatty liver disease, *Curr. Drug Metab.* 13 (2012) 1422–1435.
- [15] J.M. Brandt, F. Djouadi, D.P. Kelly, Fatty acids activate transcription of the muscle carnitine palmitoyltransferase I gene in cardiac myocytes via the peroxisome proliferator-activated receptor alpha, *J. Biol. Chem.* 273 (1998) 23786–23792.

- [16] S. Kersten, J. Seydoux, J.M. Peters, F.J. Gonzalez, B. Desvergne, W. Wahli, Peroxisome proliferator-activated receptor alpha mediates the adaptive response to fasting, *J. Clin. Invest.* 103 (1999) 1489–1498.
- [17] T.C. Leone, C.J. Weinheimer, D.P. Kelly, A critical role for the peroxisome proliferator-activated receptor alpha (PPARalpha) in the cellular fasting response: the PPARalpha-null mouse as a model of fatty acid oxidation disorders, *Proc. Natl. Acad. Sci. U. S. A.* 96 (1999) 7473–7478.
- [18] C. Mascaró, E. Acosta, J.A. Ortiz, P.F. Marrero, F.G. Hegardt, D. Haro, Control of human muscle-type carnitine palmitoyltransferase I gene transcription by peroxisome proliferator-activated receptor, *J. Biol. Chem.* 273 (1998) 8560–8563.
- [19] M. Rakhshandehroo, G. Hooiveld, M. Muller, S. Kersten, Comparative analysis of gene regulation by the transcription factor PPARalpha between mouse and human, *PLoS One* 4 (2009) e6796.
- [20] T. Aoyama, J.M. Peters, N. Iritani, T. Nakajima, K. Furihata, T. Hashimoto, F.J. Gonzalez, Altered constitutive expression of fatty acid-metabolizing enzymes in mice lacking the peroxisome proliferator-activated receptor alpha (PPARalpha), *J. Biol. Chem.* 273 (1998) 5678–5684.
- [21] U. Varanasi, R. Chu, Q. Huang, R. Castellon, A.V. Yeldandi, J.K. Reddy, Identification of a peroxisome proliferator-responsive element upstream of the human peroxisomal fatty acyl coenzyme A oxidase gene, *J. Biol. Chem.* 271 (1996) 2147–2155.
- [22] M. Cherkaoui-Malki, K. Meyer, W.Q. Cao, N. Latruffe, A.V. Yeldandi, M.S. Rao, C.A. Bradfield, J.K. Reddy, Identification of novel peroxisome proliferator-activated receptor alpha (PPARalpha) target genes in mouse liver using cDNA microarray analysis, *Gene Expr.* 9 (2001) 291–304.
- [23] M.C. Sugden, P.W. Caton, M.J. Holness, PPAR control: it's SIRTainly as easy as PGC, *J. Endocrinol.* 204 (2010) 93–104.
- [24] B.N. Finck, M.C. Gropler, Z. Chen, T.C. Leone, M.A. Croce, T.E. Harris, J.C. Lawrence Jr., D.P. Kelly, Lipin 1 is an inducible amplifier of the hepatic PGC-1alpha/PPARalpha regulatory pathway, *Cell Metab.* 4 (2006) 199–210.
- [25] J.A. Villena, A. Kralli, ERRalpha: a metabolic function for the oldest orphan, *trends endocrinol, Metab.* 19 (2008) 269–276.
- [26] V. Giguere, Transcriptional control of energy homeostasis by the estrogen-related receptors, *Endocr. Rev.* 29 (2008) 677–696.
- [27] S.N. Schreiber, D. Knutti, K. Brogli, T. Uhlmann, A. Kralli, The transcriptional coactivator PGC-1 regulates the expression and activity of the orphan nuclear receptor estrogen-related receptor alpha (ERRalpha), *J. Biol. Chem.* 278 (2003) 9013–9018.
- [28] V.K. Mootha, C. Handschin, D. Arlow, X. Xie, J. St Pierre, S. Sihag, W. Yang, D. Altshuler, P. Puigserver, N. Patterson, P.J. Willy, I.G. Schulman, R.A. Heyman, E.S. Lander, B.M. Spiegelman, Erralpha and Gabpa/b specify PGC-1alpha-dependent oxidative phosphorylation gene expression that is altered in diabetic muscle, *Proc. Natl. Acad. Sci. U. S. A.* 101 (2004) 6570–6575.
- [29] J.M. Huss, R.P. Kopp, D.P. Kelly, Peroxisome proliferator-activated receptor coactivator-1alpha (PGC-1alpha) coactivates the cardiac-enriched nuclear receptors estrogen-related receptor-alpha and -gamma. Identification of novel leucine-rich interaction motif within PGC-1alpha, *J. Biol. Chem.* 277 (2002) 40265–40274.
- [30] S. Gaillard, M.A. Dwyer, D.P. McDonnell, Definition of the molecular basis for estrogen receptor-related receptor-alpha-cofactor interactions, *Mol. Endocrinol.* 21 (2007) 62–76.
- [31] J. Rhee, Y. Inoue, J.C. Yoon, P. Puigserver, M. Fan, F.J. Gonzalez, B.M. Spiegelman, Regulation of hepatic fasting response by PPARgamma coactivator-1alpha (PGC-1): requirement for hepatocyte nuclear factor 4alpha in gluconeogenesis, *Proc. Natl. Acad. Sci. U. S. A.* 100 (2003) 4012–4017.
- [32] W.J. Lin, W.C. Yeh, Implication of toll-like receptor and tumor necrosis factor alpha signaling in septic shock, *Shock* 24 (2005) 206–209.
- [33] P.C. Calder, Long-chain n-3 fatty acids and inflammation: potential application in surgical and trauma patients, *Braz. J. Med. Biol. Res.* 36 (2003) 433–446.
- [34] B.J. Morlion, E. Torwesten, H. Lessire, G. Sturm, B.M. Peskar, P. Furst, C. Puchstein, The effect of parenteral fish oil on leukocyte membrane fatty acid composition and leukotriene-synthesizing capacity in patients with postoperative trauma, *Metab. Clin. Exp.* 45 (1996) 1208–1213.
- [35] M. Roulet, P. Frascarolo, M. Pilet, G. Chapuis, Effects of intravenously infused fish oil on platelet fatty acid phospholipid composition and on platelet function in postoperative trauma, *JPEN J. Parenter. Enter. Nutr.* 21 (1997) 296–301.
- [36] S. Sadeghi, F.A. Wallace, P.C. Calder, Dietary lipids modify the cytokine response to bacterial lipopolysaccharide in mice, *Immunology* 96 (1999) 404–410.
- [37] I. Chafchaoui-Moussaoui, Z. Charrouf, D. Guillaume, Triterpenoids from *Argania spinosa*: 20 years of research, *Nat. Prod. Commun.* 8 (2013) 43–46.
- [38] H. Berrougui, M. Cloutier, M. Isabelle, A. Khalil, Phenolic-extract from Argan oil (*Argania spinosa* L.) inhibits human low-density lipoprotein (LDL) oxidation and enhances cholesterol efflux from human THP-1 macrophages, *Atherosclerosis* 184 (2006) 389–396.
- [39] M. Cherki, A. Derouiche, A. Drissi, M. El Messal, Y. Bamou, A. Idrissi-Oudghiri, A. Khalil, A. Adlouni, Consumption of Argan oil may have an antiatherogenic effect by improving paraoxonase activities and antioxidant status: intervention study in healthy men, *Nutr. Metab. Cardiovasc. Dis.* 15 (2005) 352–360.
- [40] A. Derouiche, M. Cherki, A. Drissi, Y. Bamou, M. El Messal, A. Idrissi-Oudghiri, J.M. Lecerf, A. Adlouni, Nutritional intervention study with Argan oil in man: effects on lipids and apolipoproteins, *Ann. Nutr. Metab.* 49 (2005) 196–201.
- [41] A. Drissi, J. Girona, M. Cherki, G. Godas, A. Derouiche, M. El Messal, R. Saile, A. Kettani, R. Sola, L. Masana, A. Adlouni, Evidence of hypolipemic and antioxidant properties of Argan oil derived from the Argan tree (*Argania spinosa*), *Clin. Nutr.* 23 (2004) 1159–1166.
- [42] R. El Kebbij, S. El Kamouni, H.I. El Hajj, P. Andreoletti, J. Gresti, N. Latruffe, M.S. El Kebbij, J. Vamecq, G. Lizard, B. Nasser, M. Cherkaoui-Malki, Modulation of peroxisomes abundance by Argan oil and lipopolysaccharides in acyl-CoA oxidase 1-deficient fibroblasts, *Health* 5 (2013) 62–69.
- [43] D. Oaxaca-Castillo, P. Andreoletti, A. Vluggens, S. Yu, P.P. van Veldhoven, J.K. Reddy, M. Cherkaoui-Malki, Biochemical characterization of two functional human liver acyl-CoA oxidase isoforms 1a and 1b encoded by a single gene, *Biochem. Biophys. Res. Commun.* 360 (2007) 314–319.
- [44] J. Ni, Y. Sasaki, S. Tokuyama, A. Sogabe, Y. Tahara, Conversion of a typical catalase from *Bacillus* sp. TE124 to a catalase-peroxidase by directed evolution, *J. Biosci. Bioeng.* 93 (2002) 31–36.
- [45] D.J. Hryb, J.F. Hogg, Chain length specificities of peroxisomal and mitochondrial beta-oxidation in rat liver, *Biochem. Biophys. Res. Commun.* 87 (1979) 1200–1206.
- [46] B. Desvergne, W. Wahli, Peroxisome proliferator-activated receptors: nuclear control of metabolism, *Endocr. Rev.* 20 (1999) 649–688.
- [47] A.P. Beigneux, A.H. Moser, J.K. Shigenaga, C. Grunfeld, K.R. Feingold, The acute phase response is associated with retinoid X receptor repression in rodent liver, *J. Biol. Chem.* 275 (2000) 16390–16399.
- [48] M.S. Kim, J.K. Shigenaga, A.H. Moser, K.R. Feingold, C. Grunfeld, Suppression of estrogen-related receptor alpha and medium-chain acyl-coenzyme A dehydrogenase in the acute-phase response, *J. Lipid Res.* 46 (2005) 2282–2288.
- [49] A. Benzaria, N. Meskini, M. Dubois, M. Croset, G. Nemoz, M. Lagarde, A.F. Prigent, Effect of dietary Argan oil on fatty acid composition, proliferation, and phospholipase D activity of rat thymocytes, *Nutrition* 22 (2006) 628–637.
- [50] B. Ren, A.P. Thelen, J.M. Peters, F.J. Gonzalez, D.B. Jump, Polyunsaturated fatty acid suppression of hepatic fatty acid synthase and S14 gene expression does not require peroxisome proliferator-activated receptor alpha, *J. Biol. Chem.* 272 (1997) 26827–26832.
- [51] B.M. Forman, J. Chen, R.M. Evans, Hypolipidemic drugs, polyunsaturated fatty acids, and eicosanoids are ligands for peroxisome proliferator-activated receptors alpha and delta, *Proc. Natl. Acad. Sci. U. S. A.* 94 (1997) 4312–4317.

- [52] J. Vamecq, M. Cherkaoui-Malki, P. Andreoletti, N. Latruffe, The human peroxisome in health and disease: the story of an oddity becoming a vital organelle, *Biochimie* 98 (2014) 4–15.
- [53] M. Cherkaoui-Malki, S. Surapureddi, H.I. El-Hajj, J. Vamecq, P. Andreoletti, Hepatic steatosis and peroxisomal fatty acid beta-oxidation, *Curr. Drug Metab.* 13 (2012) 1412–1421.
- [54] T.F. Liu, C.M. Brown, M. El Gazzar, L. McPhail, P. Millet, A. Rao, V.T. Vachharajani, B.K. Yoza, C.E. McCall, Fueling the flame: bioenergy couples metabolism and inflammation, *J. Leukoc. Biol.* 92 (2012) 499–507.
- [55] N.C. Inestrosa, M. Bronfman, F. Leighton, Properties of fatty acyl-CoA oxidase from rat liver, a peroxisomal flavoprotein, *Life Sci.* 25 (1979) 1127–1135.
- [56] M. Barile, C. Brizio, D. Valenti, C. De Virgilio, S. Passarella, The riboflavin/FAD cycle in rat liver mitochondria, *Eur. J. Biochem.* 267 (2000) 4888–4900.
- [57] H. Berrougui, A. Ettaib, M.D. Herrera Gonzalez, M. Alvarez de Sotomayor, N. Bennani-Kabchi, M. Hmamouchi, Hypolipidemic and hypocholesterolemic effect of Argan oil (*Argania spinosa* L.) in Meriones shawi rats, *J. Ethnopharmacol.* 89 (2003) 15–18.
- [58] M. Bou Khalil, M. Sundaram, H.Y. Zhang, P.H. Links, J.F. Raven, B. Manmontri, M. Sariahmetoglu, K. Tran, K. Reue, D.N. Brindley, Z. Yao, The level and compartmentalization of phosphatidate phosphatase-1 (lipin-1) control the assembly and secretion of hepatic VLDL, *J. Lipid Res.* 50 (2009) 47–58.
- [59] W. Inoue, G. Somay, S. Poole, G.N. Luheshi, Immune-to-brain signaling and central prostaglandin E2 synthesis in fasted rats with altered lipopolysaccharide-induced fever, *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 295 (2008) R133–R143.
- [60] D. Patsouris, S. Mandard, P.J. Voshol, P. Escher, N.S. Tan, L.M. Havekes, W. Koenig, W. Marz, S. Tafuri, W. Wahli, M. Muller, S. Kersten, PPARalpha governs glycerol metabolism, *J. Clin. Invest.* 114 (2004) 94–103.
- [61] H. Volkoff, R.E. Peter, Effects of lipopolysaccharide treatment on feeding of goldfish: role of appetite-regulating peptides, *Brain Res.* 998 (2004) 139–147.
- [62] D. Ryu, K.J. Oh, H.Y. Jo, S. Hedrick, Y.N. Kim, Y.J. Hwang, T.S. Park, J.S. Han, C.S. Choi, M. Montminy, S.H. Koo, TORC2 regulates hepatic insulin signaling via a mammalian phosphatidic acid phosphatase, LIPIN1, *Cell Metab.* 9 (2009) 240–251.
- [63] K.R. Feingold, Y. Wang, A. Moser, J.K. Shigenaga, C. Grunfeld, LPS decreases fatty acid oxidation and nuclear hormone receptors in the kidney, *J. Lipid Res.* 49 (2008) 2179–2187.
- [64] B. Lu, Y. Lu, A.H. Moser, J.K. Shigenaga, C. Grunfeld, K.R. Feingold, LPS and proinflammatory cytokines decrease lipin-1 in mouse adipose tissue and 3T3-L1 adipocytes, *Am. J. Physiol. Endocrinol. Metab.* 295 (2008) E1502–E1509.
- [65] N. Singh, L. Li, Reduced oxidative tissue damage during endotoxemia in IRAK-1 deficient mice, *Mol. Immunol.* 50 (2012) 244–252.
- [66] M. Jia, Y. Jing, Q. Ai, R. Jiang, J. Wan, L. Lin, D. Zhou, Q. Che, L. Li, L. Tang, Y. Shen, L. Zhang, Potential role of catalase in mice with lipopolysaccharide/D-galactosamine-induced fulminant liver injury, *Hepatol. Res.* 44 (2014) 1151–1158.