



**HAL**  
open science

## **Erasure of CpG methylation in Arabidopsis alters patterns of histone H3 methylation in heterochromatin.**

Muhammad M. Tariq, Hidetoshi Saze, Aline V. Probst, Jacek Lichota, Yoshiki Habu, Jerzy Paszkowski

### ► To cite this version:

Muhammad M. Tariq, Hidetoshi Saze, Aline V. Probst, Jacek Lichota, Yoshiki Habu, et al.. Erasure of CpG methylation in Arabidopsis alters patterns of histone H3 methylation in heterochromatin.. Proceedings of the National Academy of Sciences of the United States of America, 2003, 100 (15), pp.8823-7. 10.1073/pnas.1432939100 . inserm-01629705

**HAL Id: inserm-01629705**

**<https://inserm.hal.science/inserm-01629705>**

Submitted on 6 Nov 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



tions were performed with gene-specific primers located near the 3' end of the coding region; cDNA was PCR amplified for 40 cycles as described above.

**Immunostaining.** Leaf protoplasts were isolated and fixed according to [www.arabidopsis.org/cshl-course/7-gene-expression.html](http://www.arabidopsis.org/cshl-course/7-gene-expression.html). After rehydration in PBS, slides were blocked in 2% BSA in PBS (30 min, 37°C) and incubated overnight at 4°C in 1% BSA in PBS containing antibodies (Upstate Biotechnology) specific to lysine-9-dimethylated H3 (1:100 dilution), lysine-4-dimethylated H3 (1:500 dilution), or tetraacetylated H4 (1:100 dilution). Detection was carried out with an FITC-coupled antibody to rabbit IgG (Molecular Probes; 1:100 dilution, 37°C, 40 min) in 0.5% BSA in PBS. DNA was counterstained with 4',6-diamidino-2-phenylindole (DAPI) in Vectashield (Vector Laboratories). Images were analyzed with a Deltavision deconvolution microscope (Applied Precision, Issaquah, WA). WORX software (Applied Precision) was applied to deconvolute the images; single representative layers were chosen for Fig. 3. Fluorescent *in situ* hybridization with a 180-bp repeats probe and immunostaining analysis with a 5-methylcytosine antibody were performed as described (13, 21).


**Western-Blot Analysis.** Leaves of 4-week-old plants were homogenized in histone-extraction buffer (10 mM Tris-HCl, pH 7.5/2 mM EDTA/0.25 M HCl/5 mM 2-mercaptoethanol/0.2 mM PMSF). After centrifugation at 12,000 × *g* for 10 min, soluble proteins were precipitated with 25% trichloroacetic acid and collected by centrifugation at 17,000 × *g* for 30 min. The pellet was washed twice with acetone, resuspended in Laemmli buffer, separated electrophoretically, and Western blotted and detected by using standard procedures (22).

## Results and Discussion


In our strain (named *met1* throughout this article), we determined the methylation status of histone H3 at the nontranscribed *Ta2* retrotransposon that resides in a pericentromeric region of chromosome 1. *Ta2* has been shown to contain hypermethylated

DNA that is packaged to nucleosomes with H3K9<sup>Me</sup> (12). We compared WT and *met1* plants by using ChIP with antibodies recognizing H3K9<sup>Me</sup> or histone H3 dimethylated at lysine 4 (H3K4<sup>Me</sup>) (Fig. 1 *A* and *B*). H3K9<sup>Me</sup>, which is abundant at *Ta2* in WT, was almost undetectable in the *met1* mutant (Fig. 1 *A* and *B*). In *met1*, *Ta2* consistently gained H3K4<sup>Me</sup> (Fig. 1 *A* and *B*), a marker for transcriptionally active genes (6). Indeed, in the *met1* mutant, transcription of *Ta2* was clearly activated (Fig. 1 *C*). Depletion of CpG methylation at *Ta2*, therefore, likely changed the methylation status of histone H3. This change, however, could have been an indirect effect of transcriptional activation. To assess the generality of this observation and relate it to transcriptional reactivation, we examined levels of H3K9<sup>Me</sup> and H3K4<sup>Me</sup> in three additional target genes residing within the heterochromatic knob on chromosome 4 (At4g03760, T5L23.26, and At4g03870) (Fig. 1 *A* and *B*). Like *Ta2*, these loci are associated with H3K9<sup>Me</sup> in the WT (19). Four independent ChIP experiments were performed for each of the four target sequences and the *Actin2/7* control (20); these ChIP experiments yielded equivalent results (Fig. 1 *A* and *B*). The abundant H3K9<sup>Me</sup> found in all of the targets in the WT was drastically reduced and replaced by H3K4<sup>Me</sup> in the *met1* mutant (Fig. 1 *A* and *B*). When transcript levels originating from these loci in WT and *met1* (Fig. 1 *C*) were compared, transcripts were detected in the *met1* mutant for *Ta2*, At4g03760, and T5L23.26 (Fig. 1 *C*), but not for At4g03870 (Fig. 1 *C*). This finding suggested that the loss of H3K9<sup>Me</sup> in the *met1* mutant could occur also at silent templates; very low levels of transcription, however, cannot be excluded.

To substantiate this observation, we examined H3K9<sup>Me</sup> at 180-bp centromeric repeats considered transcriptionally inert in both WT and DNA methylation-deficient mutants (12). To examine changes in DNA methylation and histone modifications accurately and correlate them to the transcriptional status of repeats, we designed primers (180F4R5) specific to a single-copy sequence (GenBank accession no. AC083859, nucleotides 61536–62127) that flanks a particular set of 180-bp repeats residing in the pericentromeric



**Fig. 1.** Histone 3 methylation analyzed by ChIP and transcript levels of selected heterochromatic loci. (A) ChIP with antibodies recognizing H3K4<sup>Me</sup> (K4) and H3K9<sup>Me</sup> (K9) of chromatin extracts from WT and the *met1* mutant (*met1*). Controls: chromatin extract precipitated without antibody (mock), chromatin extract (input), and PCR without template (-ve). Analyzed target loci are listed to the left of the corresponding images. (B) Quantification of ChIP analysis corresponding to images presented in A. Ordinates are fold difference. (C) RT-PCR analysis of transcripts, with (+) and without (-) reverse transcriptase (RT).




**Fig. 2.** Histone 3 methylation analysis by ChIP of 180-bp centromeric repeats. (A) ChIP with antibodies recognizing H3K4<sup>Me</sup> (K4) and H3K9<sup>Me</sup> (K9) of chromatin extracts from WT and the *met1* mutant (*met1*); controls as in Fig. 1. Amplification was performed using primers to a specific subset (180F4R5) or broader range of repeats, as described in the text. (B) Quantification of ChIP analysis corresponding to images presented in A. Ordinates are fold difference.


region on chromosome 1. This primer eliminated interference by other 180-bp repeats distributed over all five chromosomes. As revealed by a bisulfite analysis assay, CpG methylation on 180F4R5 repeats is completely erased in *met1*, but reduced only to 57.6% and 73% at CpNpG and CpNpN sites, respectively (18). Analysis of other target sequences showed similar results (data not shown). The ChIP assays showed a strong enrichment in H3K9<sup>Me</sup> at these repeats in WT plants and a 17-fold depletion in the *met1* mutant (Fig. 2A and B). Importantly, multiple RT-PCR analyses did not detect RNA from this region in either WT or the *met1* mutant (data not shown). To examine whether this finding was specific to this particular subset of 180-bp repeats or could be confirmed globally,

we used unselective 180-bp repeat primers that generate a ladder because of multiple annealing sites throughout the genome (12). The *met1* mutation also caused a strong reduction in H3K9<sup>Me</sup> at these combined targets (Fig. 2A and B). In an RT-PCR assay with the same primers, however, we detected the appearance of traces of RNA in the *met1* mutant (data not shown). This appearance of RNA traces is consistent with the earlier studies (12).

Because 180-bp repeats are an integral part of the centromeric region of all *Arabidopsis* chromosomes and histone methylation changes in the *met1* mutant seem to be rather drastic, we attempted to visualize H3K9<sup>Me</sup> alterations by immunostaining WT and *met1* nuclei by using an antibody that recognizes



**Fig. 3.** Immunolocalization of H3K9<sup>Me</sup>, H3K4<sup>Me</sup>, and tetraacetylated histone H4 in WT and *met1* nuclei. (A) Detection of H3K9<sup>Me</sup>. (B) Detection of H3K4<sup>Me</sup>. (C) Detection of H4 acetylation. Shown are FITC immunostaining (green) and DAPI staining (blue). Chromocenters are visible as light blue, and densely DAPI-stained structures are marked by arrowheads in the leftmost A. (D) Visualization of WT and *met1* chromocenters with fluorescent *in situ* hybridization using 180-bp centromeric repeats probe (red) and immunostaining using 5-methylcytosine antibody ( $\alpha$ -5MeC, green). (E) Analysis of H3 and H4 methylation and acetylation, respectively, from WT and *met1* plants by Western blot probed with antibody directed against H3K9<sup>Me</sup> ( $\alpha$ -H3K9<sup>Me</sup>), H3K4<sup>Me</sup> ( $\alpha$ -H3K4<sup>Me</sup>), or tetraacetylated H4 ( $\alpha$ -H4Ac). (Bottom) Coomassie blue-stained proteins.



**Fig. 4.** Histone 4 acetylation analyzed by ChIP of selected heterochromatic loci. (A) ChIP with antibodies recognizing histone H4 acetylated at lysines 5, 8, 12, and 16 (H4Ac) of chromatin extracts from WT and the *met1* mutant; controls as in Fig. 1. Analyzed target loci are listed to the left of corresponding images. (B) Quantification of ChIP analysis corresponding to images presented in A. H4Ac was not detected at T5L23.26 and At4g03870 in either WT or *met1*. Ordinates are fold difference.

H3K9<sup>Me</sup> ( $\alpha$ -H3K9). In the WT, the dense heterochromatic chromocenters, known to be enriched in H3K9<sup>Me</sup>, were clearly stained for H3K9<sup>Me</sup> (Fig. 3A). Similar staining performed previously with a point-mutant allele in the *MET1* gene showed a reduction in H3K9<sup>Me</sup>, but part of the signal was still retained at the chromocenters (13). In *met1* nuclei, in contrast, chromocenters detectable by DAPI staining remained unlabeled by the H3K9<sup>Me</sup> antibody (Fig. 3A). Thus in *met1*, the H3K9<sup>Me</sup> signal was reduced and completely redistributed away from chromocenters (Fig. 3A). Very similar images of *met1* nuclei were obtained with antibodies directed against 5-methylcytosine (Fig. 3D), confirming a drastic, concerted change in DNA methylation and H3K9<sup>Me</sup> at the centromeric heterochromatin. Surprisingly, images obtained by fluorescent *in situ* hybridization with probes specific to centromeric 180-bp repeats appear to be very similar for the *met1* and WT (Fig. 3D), which is in contrast to the *ddm1* (decrease in DNA methylation 1) mutant, where centromeric DNA disperses (13, 23). The complete depletion of methylated CpGs and drastic reduction of H3K9<sup>Me</sup> in the *met1* strain suggest that both modifications are dispensable for maintaining the compact, heterochromatic structures enclosing centromeric DNA. This finding is in accordance with a recent cytological study of the *kyp* mutant, indicating that H3K9<sup>Me</sup> is unessential for the maintenance of heterochromatic structure of *Arabidopsis* chromocenters (24).

In contrast to immunostaining for H3K9<sup>Me</sup>,  $\alpha$ -H3K4<sup>Me</sup> showed exclusion of the signal from chromocenters (Fig. 3B) in both WT and *met1*. Thus, the cytological data were in agreement with ChIP analyses, demonstrating a severe reduction in H3K9<sup>Me</sup> at 180-bp repeats but retention of a constantly low level of H3K4<sup>Me</sup> in the *met1* strain. Because cytological data suggested redistribution of H3K9<sup>Me</sup> away from chromocenters in the *met1* mutant (rather than a general reduction in level) we compared

by Western blotting the proportion of H3K9<sup>Me</sup> in enriched histone fractions. Indeed, similar levels of H3K9<sup>Me</sup> were detected in WT and *met1*, but there appeared to be an increase in abundance of H3K4<sup>Me</sup>, which may reflect the alleviation of silencing (Fig. 3E).

A tight link between CpG methylation and histone deacetylation and transcriptional repression has been postulated in mammalian cells (25–28). We investigated, therefore, whether depletion of CpG methylation in the *met1* mutant affects levels of histone acetylation that could be related to changes in transcriptional activity. We performed ChIP with a tetraacetylated histone H4 antibody ( $\alpha$ -H4Ac) by using all of the target sequences described above. According to the ChIP analyses, acetylated H4 (H4Ac) was enriched approximately 3- to 6-fold in the *met1* mutant compared with the WT. However, this enrichment was restricted to targets that were transcriptionally reactivated in the *met1* (Fig. 4). H4Ac was not detected at the At4g03870 locus or at nontranscribed 180-bp repeats (Fig. 4 and data not shown). Notably, H4Ac was almost undetectable at the T5L23.26 locus, reactivated in the *met1* mutant (Figs. 1C and 4). Hyperacetylation, however, is not always associated with transcriptional reactivation of previously silent loci (29, 30) and the mechanism of T5L23.26 activation may be of this category. Immunostaining of WT and *met1* nuclei with  $\alpha$ -H4Ac antibody showed that chromocenters remained hypoacetylated in the mutant (Fig. 3C). Western blotting showed no change in histone-acetylation levels (Fig. 3E).

Our results clearly support the notion that methylation at H3K9 is directed by DNA methylation at CpG sites (13). They show, in addition, that this process probably occurs in a transcription-independent fashion. Obviously, very low transcriptional activity might have escaped detection. The recently discovered *in vitro* and *in vivo* interaction of human proteins

recognizing methylated cytosine in CpG (MeCP2, MBD1) with histone methyltransferase also suggests such a link (31, 32). The effect of CpG methylation loss in the *met1* is in contrast to methylation outside CpGs, which has no effect on H3K9<sup>Me</sup> when strongly reduced by *cmt3* (chromomethylase 3) mutations (12, 33). Interestingly, in *kyp* mutants defective in histone H3 lysine 9 methyltransferase DNA methylation marks are affected only at CpNpG and CpNpN sites (not at CpGs sites), suggesting that non-CpG methylation is regulated by histone methylation (10). Recently, the putative chromatin-remodeling factor DDM1 (decrease in DNA methylation 1) was shown to be required for maintenance of H3K9<sup>Me</sup> (19). In *ddm1* mutants, however, cytosine methylation in all sequence contexts including CpGs is affected, making it difficult to distinguish direct from indirect effects on histone methylation. Considering that CpG methylation determines the status of histone methylation, the loss of histone methylation in the *ddm1* mutants could be due to the depletion of CpG methylation. Both *met1* and *ddm1* mutations also reduce levels of cytosine methylation in sequences outside CpGs. Although this reduction may contribute to the loss of histone methylation (12), given that DNA methylation outside CpGs acts downstream of histone methylation (10), changes in

CpNpG and CpNpN observed in *met1* and *ddm1* are most likely secondary effects of these mutations on H3K9<sup>Me</sup>. It is notable that, in contrast to *met1*, chromocenters in *ddm1* nuclei decompose (13, 21). It is conceivable, therefore, that structural alterations of heterochromatin seen in the *ddm1* mutant are not linked directly to the methylated CpG/H3K9<sup>Me</sup> depletion but to another, unknown mechanism. Because cytological data suggested redistribution of H3K9<sup>Me</sup> in the *met1* mutant, it would be of interest to map this redistribution in more detail.

Our results clearly support the existence *in vivo* of a self-reinforcing system that contributes to the formation of silent heterochromatin in which CpG methylation plays the role of a central scaffold directing histone methylation, possibly independent of transcription. This function of CpG methylation explains the severe and immediate phenotypes of *met1* and the gametophytic effects (18) that are much less pronounced or absent in the weaker *met1* mutant allele (16) used in all previous studies.

We thank Fang Lin Sun for helpful discussions regarding ChIP, Karim Afsar and Susanne Lienhardt for technical assistance, and Thomas Hohn, Witold Filipowicz, Barbara Hohn, Anne Blonstein, and Patrick King for helpful comments on the manuscript.

1. Finnegan, E. J. & Kovac, K. A. (2000) *Plant Mol. Biol.* **43**, 189–201.
2. Bird, A. (2002) *Genes Dev.* **16**, 6–21.
3. Li, E., Bestor, T. H. & Jaenisch, R. (1992) *Cell* **69**, 915–926.
4. Finnegan, E. J., Peacock, W. J. & Dennis, E. S. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 8449–8454.
5. Ronemus, M. J., Galbiati, M., Ticknor, C., Chen, J. & Dellaporta, S. L. (1996) *Science* **273**, 654–657.
6. Jenuwein, T. & Allis, C. D. (2001) *Science* **293**, 1074–1080.
7. Tamaru, H. & Selker, E. U. (2001) *Nature* **414**, 277–283.
8. Tamaru, H., Zhang, X., McMillen, D., Singh, P. B., Nakayama, J., Grewal, S. I., Allis, C. D., Cheng, X. & Selker, E. U. (2003) *Nat. Genet.* **34**, 75–79.
9. Kouzminova, E. & Selker, E. U. (2001) *EMBO J.* **20**, 4309–4323.
10. Jackson, J. P., Lindroth, A. M., Cao, X. & Jacobsen, S. E. (2002) *Nature* **416**, 556–560.
11. Zilberman, D., Cao, X. & Jacobsen, S. E. (2003) *Science* **299**, 716–719.
12. Johnson, L., Cao, X. & Jacobsen, S. E. (2002) *Curr. Biol.* **12**, 1360–1367.
13. Soppe, W. J., Jasencakova, Z., Houben, A., Kakutani, T., Meister, A., Huang, M. S., Jacobsen, S. E., Schubert, I. & Fransz, P. F. (2002) *EMBO J.* **21**, 6549–6559.
14. Barteel, L. & Bender, J. (2001) *Nucleic Acids Res.* **29**, 2127–2134.
15. Vongs, A., Kakutani, T., Martienssen, R. A. & Richards, E. J. (1993) *Science* **260**, 1926–1928.
16. Kankel, M. W., Ramsey, D. E., Stokes, T. L., Flowers, S. K., Haag, J. R., Jeddeloh, J. A., Riddle, N. C., Verbsky, M. L. & Richards, E. J. (2003) *Genetics* **163**, 1109–1122.
17. Ahmad, K. & Henikoff, S. (2002) *Mol. Cell* **9**, 1191–1200.
18. Saze, H., Mittelsten Scheid, O. & Paszkowski, J. (2003) *Nat. Genet.* **34**, 65–69.
19. Gendrel, A. V., Lippman, Z., Yordan, C., Colot, V. & Martienssen, R. A. (2002) *Science* **297**, 1871–1873.
20. An, Y. Q., McDowell, J. M., Huang, S., McKinney, E. C., Chambliss, S. & Meagher, R. B. (1996) *Plant J.* **10**, 107–121.
21. Mittelsten Scheid, O., Probst, A. V., Afsar, K. & Paszkowski, J. (2002) *Proc. Natl. Acad. Sci. USA* **99**, 13659–13662.
22. Sambrook, J., Fritsch, E. F. & Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Lab. Press, Plainview, NY), 2nd Ed.
23. Probst, A. V., Fransz, P. F., Paszkowski, J. & Scheid, O. M. (2003) *Plant J.* **33**, 743–749.
24. Jasencakova, Z., Soppe, W. J., Meister, A., Gernand, D., Turner, B. M. & Schubert, I. (2003) *Plant J.* **33**, 471–480.
25. Ng, H. H., Zhang, Y., Hendrich, B., Johnson, C. A., Turner, B. M., Erdjument-Bromage, H., Tempst, P., Reinberg, D. & Bird, A. (1999) *Nat. Genet.* **23**, 58–61.
26. Wade, P. A., Geggion, A., Jones, P. L., Ballestar, E., Aubry, F. & Wolffe, A. P. (1999) *Nat. Genet.* **23**, 62–66.
27. Jones, P. L., Veenstra, G. J., Wade, P. A., Vermaak, D., Kass, S. U., Landsberger, N., Strouboulis, J. & Wolffe, A. P. (1998) *Nat. Genet.* **19**, 187–191.
28. Nan, X., Ng, H. H., Johnson, C. A., Laherty, C. D., Turner, B. M., Eisenman, R. N. & Bird, A. (1998) *Nature* **393**, 386–389.
29. Lorincz, M. C., Schubert, D., Goeke, S. C., Walters, M., Groudine, M. & Martin, D. I. (2000) *Mol. Cell Biol.* **20**, 842–850.
30. Benjamin, D. & Jost, J. P. (2001) *Nucleic Acids Res.* **29**, 3603–3610.
31. Fuks, F., Hurd, P. J., Wolf, D., Nan, X., Bird, A. P. & Kouzarides, T. (2003) *J. Biol. Chem.* **278**, 4035–4040.
32. Fujita, N., Watanabe, S., Ichimura, T., Tsuruzoe, S., Shinkai, Y., Tachibana, M., Chiba, T. & Nakao, M. (April 23, 2003) *J. Biol. Chem.*, 10.1074/jbc.M302283200.
33. Lindroth, A. M., Cao, X., Jackson, J. P., Zilberman, D., McCallum, C. M., Henikoff, S. & Jacobsen, S. E. (2001) *Science* **292**, 2077–2080.