



HAL
open science

Alcohol, tobacco and cannabis use are associated with job loss at follow-up: Findings from the CONSTANCES cohort

Guillaume Airagnes, Cédric Lemogne, Pierre Meneton, Marie Plessz, Marcel Goldberg, Nicolas Hoertel, Yves Roquelaure, Frédéric Limosin, Marie Zins

► To cite this version:

Guillaume Airagnes, Cédric Lemogne, Pierre Meneton, Marie Plessz, Marcel Goldberg, et al.. Alcohol, tobacco and cannabis use are associated with job loss at follow-up: Findings from the CONSTANCES cohort. PLoS ONE, 2019, 14 (9), pp.e0222361. 10.1371/journal.pone.0222361 . inserm-02316589

HAL Id: inserm-02316589

<https://inserm.hal.science/inserm-02316589>

Submitted on 15 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH ARTICLE

Alcohol, tobacco and cannabis use are associated with job loss at follow-up: Findings from the CONSTANCES cohort

Guillaume Airagnes^{1,2,3,4*}, Cédric Lemogne^{1,2,5}, Pierre Meneton⁶, Marie Plessz⁶, Marcel Goldberg^{2,3}, Nicolas Hoertel^{1,2,5}, Yves Roquelaure⁷, Frédéric Limosin^{1,2,5}, Marie Zins^{2,3,4}

1 Hôpitaux Universitaires Paris Ouest, Department of Psychiatry and Addictology, AP-HP, Paris, France, **2** Faculté de Médecine, Université Paris Descartes, Sorbonne Paris Cité, Paris, France, **3** UMS 011, Population-based Epidemiological Cohorts, Inserm, Villejuif, France, **4** UMR 1168, VIMA, Inserm, Villejuif, France, **5** U 894, Centre Psychiatrie et Neurosciences, Inserm, Paris, France, **6** UMR 1142, Inserm, Sorbonne Université, Université Paris, Paris, France, **7** UMR 1085, Ester, Irest Inserm, Université d'Angers, Angers, France

* guillaume.airagnes@aphp.fr



Abstract

OPEN ACCESS

Citation: Airagnes G, Lemogne C, Meneton P, Plessz M, Goldberg M, Hoertel N, et al. (2019) Alcohol, tobacco and cannabis use are associated with job loss at follow-up: Findings from the CONSTANCES cohort. *PLoS ONE* 14(9): e0222361. <https://doi.org/10.1371/journal.pone.0222361>

Editor: Raymond Niaura, Global Public Health, New York University, UNITED STATES

Received: March 22, 2019

Accepted: August 27, 2019

Published: September 9, 2019

Copyright: © 2019 Airagnes et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Personal health data underlying the findings of our study are not publicly available due to legal reasons related to data privacy protection. However, the data are available upon request to all interested researchers after authorization of the French "Commission nationale de l'informatique et des libertés". The persons to contact are Dr. Marie Zins ([rf. mresni@sniz.eiram](mailto:mresni@sniz.eiram)) or Pr. Marcel Goldberg ([rf. mresni@grebdlog.lecram](mailto:mresni@grebdlog.lecram)).

Background

Substance use is more prevalent among unemployed subjects compared to employed ones. However, quantifying the risk subsequent of job loss at short-term according to substance use remains underexplored as well as examining if this association persists across various sociodemographic and occupational positions previously linked to job loss. We examined this issue prospectively for alcohol, tobacco, cannabis use and their combination, among a large population-based sample of men and women, while taking into account age, gender, overall health status and depressive symptoms.

Methods

From the French population-based CONSTANCES cohort, 18,879 working participants were included between 2012 and 2016. At baseline, alcohol use disorder risk according to the Alcohol Use Disorders Identification Test (mild, dangerous, problematic or dependence), tobacco (non-smoker, former smoker, 1–9, 10–19, >19 cigarettes/day) and cannabis use (never, not in past year, less than once a month, once a month or more) were assessed. Employment status at one-year (working versus not working) was the dependent variable. Logistic regressions provided Odds Ratios (OR(95%CI)) of job loss at one-year, adjusting for age, gender, self-reported health and depressive state (measured with the Center of Epidemiologic Studies Depression scale). Stratified analyses were performed for education, occupational grade, household income, job stress (measured with the Effort-Reward Imbalance), type of job contract, type of work time and history of unemployment. In sensitivity analyses, employment status over a three-year follow-up was used as dependent variable.

Funding: The CONSTANCES cohort is supported by the Caisse Nationale d'Assurance Maladie des travailleurs salariés-CNAMTS. CONSTANCES is accredited as a "National Infrastructure for Biology and health" by the governmental Investissements d'avenir program and was funded by the Agence nationale de la recherche (ANR-11-INBS-0002 Grant). CONSTANCES also receives funding from MSD, AstraZeneca and Lundbeck managed by INSERM-Transfert. This study has received a funding from the Interministerial Mission for Combating Drugs and Addictive Behaviors ("Mission Interministérielle de Lutte contre les Drogues et les Conduites Addictives", MILDECA). None of the authors are salaried by the funders of the CONSTANCES cohort. The funders did not have any role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: GA has received speaker and consulting fees from Lundbeck and Pfizer. CL has received speaker and consulting fees from Daiichi-Sankyo, Janssen, Lundbeck, Otsuka Pharmaceuticals and Servier. FL has received speaker and consulting fees from Astra Zeneca, Euthérapie-Servier, Janssen, Lundbeck, Otsuka Pharmaceuticals and Roche. PM, MP, MG, NH, YR and MZ have nothing to declare. No other relationships or activities that could appear to have influenced the submitted work. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

Results

Alcohol, tobacco and cannabis use were associated with job loss, from the second to the highest category: 1.46(95%CI:1.23–1.73) to 1.92(95%CI:1.34–2.75), 1.26(95%CI:1.09–1.46) to 1.78(95%CI:1.26–2.54) and 1.45(95%CI:1.27–1.66) to 2.68(95%CI:2.10–3.42), respectively, and with dose-dependent relationships (all *p* for trend <0.001). When introduced simultaneously, associations remained significant for the three substances without any between-substance interactions. Associations remained significant across almost all stratifications and over a three-year follow-up as well as after adjustment for all the sociodemographic and occupational factors.

Conclusions

Alcohol, tobacco and cannabis use were independently associated with job loss at short-term, with dose-dependent relationships. This knowledge will help refining information and prevention strategies. Importantly, even moderate levels of alcohol, tobacco or cannabis use are associated with job loss at short-term and all sociodemographic and occupational positions are potentially concerned.

Introduction

The costs associated with alcohol amount to more than 1% of the gross national product in high-income and middle-income countries, with the costs of social harm constituting a major proportion in addition to health costs [1]. The social costs of smoking are usually estimated to be even higher than those of alcohol use, and the social costs of illicit drugs could be around billions of Euros in European countries [2]. Substance use is among the leading preventable causes of premature death in the world and this burden is mainly driven by three substances: alcohol, tobacco and cannabis [3]. Regarding social harm, the association between substance use and unemployment, a major issue in most high-income countries, with a mean unemployment rate of 7.1% in Europe [4], is a critical question. Substance use is more prevalent among unemployed subjects compared to employed ones [5,6]. These findings are in accordance with an increased risk of substance use after job loss [7]. The onset of substance use after job loss might contribute to the increased morbidity and mortality observed among unemployed subjects [8]. However, substance use has been associated with detrimental consequences on occupational life and may predate job loss [5–10]. Several longitudinal studies have shown an association between substance use and the risk of subsequent job loss [11,12].

A large prospective study found an association between alcohol problematic use and subsequent unemployment during a 14-year follow-up [10]. An association between high alcohol consumption and job loss at 6 months was found among 658 at-risk drinkers [13]. High alcohol consumption and smoking were associated with long-term unemployment over a 4-year follow-up among 586 male blue-collar workers [14]. Heavy smoking (i.e. 20 cigarettes per day or more) was associated with an increased risk of being unemployed 4 years later, in men but not in women, among 5,355 employees [15]. A chronic use of cannabis in early adolescence was associated with a higher likelihood of unemployment at midlife among 548 participants [16]. Consistently, associations between cannabis use at age 18 and unemployment at age 40 were found among a large cohort of men [17]. Despite these evidence linking substance use and subsequent job loss, quantifying this risk of job loss at short-term in a large national

population-based sample remains underexplored as well as examining if this association persists across various sociodemographic and occupational positions. This knowledge is critical to inform decision-makers in public health and occupational health and refine prevention measures by targeting individuals at higher risk for job loss and by using these findings as a potential motivational tool. However, to the best of our knowledge, no study has examined prospectively the risk of job loss at short-term according to substance use while stratifying by sociodemographic and occupational factors known to be related to substance use or job loss. These factors include sociodemographic variables such as age, gender, education level, marital status and household income, health-related factors such overall health status and depressive symptoms, and occupational factors such as occupational grade, job stress, type of job contract, type of work time and a past history of unemployment [5,6,18–27]. In addition, alcohol, tobacco and cannabis have rarely been studied together, which makes it difficult to know the specific role of each substance.

We took advantage of the CONSTANCES cohort to examine the prospective associations of alcohol, tobacco, cannabis use, and their combination, with job loss among a population-based sample of men and women large enough to allow stratifying for several sociodemographic and occupational factors. First, we searched for an increased risk of job loss according to substance use in the short term (i.e. at one year), and then whether this risk persists over a longer period of time (i.e. a three-year period). Second, we examined whether those associations persist within different subgroups defined based on sociodemographic and occupational factors previously linked to job loss, in order to identify those at greater risk for job loss associated with substance use. Because gender, age, overall health status and depressive symptoms may be potential confounders [5,15,25], we adjusted all the analyses for these variables. We hypothesized that the three substances will be associated with an increased risk of job loss at one-year and that these associations will hold across the different subgroups.

Methods

Participants

The CONSTANCES cohort includes volunteers aged 18–69 years at baseline according to a random sampling scheme stratified on age, gender, socioeconomic status and region of France [28]. Participants fulfill at baseline, and then annually, questionnaires on social and demographic characteristics, health-related behaviors and occupational conditions. All the questionnaires are available at www.constances.fr in both the original language and English. From the participants included between February 2012 and September 2016 ($n = 81,997$), the population study was restricted by the following inclusion criteria consistent with our objectives: being employed at baseline; having been included at least one year ago to allow sufficient follow-up duration; not being retired at one year. Thus, 18,879 participants have been included in the present study (SI Fig). Depending on their inclusion date, the included participants had either a one-year follow-up ($n = 10,145$), a two-year follow-up ($n = 6,454$) or a three-year follow-up ($n = 2,280$). Precisely, among the participants who had only one year of follow-up, 84% ($n = 8,556$) were included too recently to have two years of follow-up. Similarly, among the participants who had two years of follow-up, 89% ($n = 5,741$) were included too recently to have three years of follow-up.

Employment status

Employment status was self-reported at baseline and annually by answering the following question: “What is your current employment situation? 1) employed, including on sick leave, leave without pay or availability, maternity/paternity/adoption/parental leave; 2) job seeker or

looking for a job; 3) retired or withdrawn from business; 4) in training (high school student, student, trainee, apprentice . . .); 5) does not work for health reasons (disability, chronic illness, etc.); 6) without employment; 7) other situation". Participants employed were those who gave the response '1' or '4'. Retired participants were those who gave the response '3'. Participants experiencing job loss were those who reported at follow-up responses '2', '5', '6' or '7'.

Substances

Alcohol use. We used the French version of the Alcohol Use Disorders Identification Test (AUDIT) to assess the risk of alcohol use disorder at baseline [29]. From the total score, we computed a categorical variable based on recommended AUDIT risk levels, as follows: 1) Mild (0–7), 2) Dangerous (8–15), 3) Problematic (16–19), and 4) Dependence (20–40). The two last categories were merged to ensure sufficient subsample size. The AUDIT has been also used after aggregating "Dangerous", "Problematic" and "Dependence" categories and as binary variable as follows: mild risk (0–7) *versus* at-risk (8–40) alcohol use.

Tobacco use. Smoking status (i.e. never smoker, former smoker or current smoker) was self-reported at baseline. Among current smokers, daily tobacco consumption was collected in cigarettes per day. From these two variables, we computed a categorical variable to define: 1) Never smokers, 2) Former smokers, 3) Current light smokers (1–9 cigarettes per day), 4) Current moderate smokers (10–19 cigarettes per day), and 5) Current heavy smokers (>19 cigarettes per day) [30]. Tobacco use has been also used after aggregating "current moderate smokers" and "current heavy smokers" and as binary variable by aggregating the first two categories (i.e. non-smokers) and the last three categories (i.e. current smokers).

Cannabis use. From three questions asked at baseline to characterize the frequency of cannabis consumption, we computed a categorical variable expressing the frequency of lifetime cannabis consumption as follows: 1) Never used; 2) No consumption during the previous 12 months; 3) Less than once a month; and 4) Once a month or more. Cannabis use has been also used as binary variable by aggregating the first two categories (i.e. no consumption during the year) and the last two categories (i.e. at least one consumption in the year).

Adjustment variables

From the baseline questionnaires, we used age, gender, poor self-reported health and depressive state as covariables of adjustment. Age in years was used as a categorical variable in three modalities as follows: 1) <30; 2) ≥ 30 and <50 and 3) ≥ 50 . Self-rated health status was measured from the following question: « How do you judge your general health compared to a person of your entourage of the same age? » to provide a proxy of overall health condition [31]. We computed a binary variable to define poor self-reported health (5 to 8, roughly corresponding to the 90th percentile) *versus* good (1 to 4). Depressive symptoms were assessed with the Center of Epidemiologic Studies Depressive state scale (CESD) taking a global score ≥ 19 to signal clinically significant depressive state [32].

Stratification variables

To search whether the association between substance use and subsequent job loss persist in different subgroups based on sociodemographic and occupational factors previously linked to job loss, we performed stratified analyses according to these factors. Thus, we examined whether these variables could act as effect modifiers. The following sociodemographic and occupational variables were used from the baseline questionnaires: age, gender, household income (i.e. <2800 *versus* ≥ 2800 euros per month), type of job contract (i.e. open-ended contract *versus* other type of contract), type of work time (i.e. full-time *versus* part-time), and

education level based on the International Standard Classification of Education, as follows: levels 0 to 4 (up to upper secondary education or post-secondary non-tertiary education), levels 5 and 6 (short-cycle tertiary education and Bachelor's or equivalent level), and levels 7 and 8 (Master's or equivalent level and Doctoral or equivalent level) [33]. Occupational grade was used in three categories as follows: 1) Blue-collar worker, craftsman or employee; 2) Intermediate worker; 3) Executive. Regarding participants still in initial training ($n = 379$; i.e. 2% of the entire sample), we used the highest occupational grade of their parents, whether it was their father or their mother. As a standardized measure of job stress, the effort-reward imbalance was assessed at baseline [26]. We used a binary variable based on tertiles as follows: 1) first and second tertiles (i.e. low job stress), and 2) third tertile (i.e. high job stress). Regarding history of unemployment, we used data from national administrative databases that are annually linked with the CONSTANCES cohort [28]. From these data, we computed a binary variable for past time unemployed in the three years before inclusion.

Statistical analysis

We first used binomial logistic regressions to compute risks of subsequent job loss according to substance use at baseline. Mild alcohol use risk, non-smoking and never used were chosen as reference for alcohol, tobacco and cannabis use, respectively. All the models were adjusted for age, gender, self-reported health and depressive symptoms. Results are presented as estimated Odds Ratios (OR) with their 95% Confidence Intervals (CI).

Regarding our aim to search for increased risk of job loss according to alcohol, tobacco and cannabis use and their combination over a one-year follow-up, we first introduced alcohol, tobacco and cannabis use separately in the models. Dose-dependent relationships were searched by introducing the substances as continuous variables. Interactions between each substance and each covariable were examined. Then, to further explore the specific and common effects of these three substances, we introduced simultaneously alcohol, tobacco and cannabis use in the same model. We primarily chose to keep at least three categories of use for all substances to inform on a potential gradient in their relations with job loss. However, in order to search for statistical significance while preventing from lack of power due to small sample sizes in some categories, this analysis has also been made after aggregating the “dangerous” and “problematic or dependence” categories regarding alcohol use, and the “moderate” and “heavy” categories regarding tobacco use. Finally, interactions between the three substances were examined by testing in pairs all the interactions in three different models. In this analysis, alcohol, tobacco and cannabis use were introduced in the models as binary variables in order to ease the search and interpretation of potential between-substance interactions.

Regarding our aim to search for at-risk subgroups, separate stratified analyses for each stratification variable were computed. Alcohol, tobacco and cannabis use were introduced in the models as binary variables, in order to ease the search and interpretation of potential effect modifiers.

Sensitivity analyses were conducted. First, since some subjects could have been aware of their risk of job loss one year before, we planned to search for similar associations over a longer duration of follow-up in order to take into account for this potential bias of reverse causality. Thus, we performed generalized estimating equations to search for similar associations while taking into account data at one-year, two-year and three-year follow-up. Alcohol, tobacco and cannabis use were entered successively in the models with occupational status as outcome. Interactions between duration of follow-up and each substance were examined. Second, we adjusted our analyses for all the covariables (i.e. age, gender, depressive symptoms, self-reported health, education level, occupational grade, household income, job stress, type of

contract, type of work time, past time unemployed). Third, we examined whether the associations between substance use and subsequent risk of job loss could persist while excluding unemployed participants at follow-up with another status than job seeking (i.e. those who do not work for health reasons ($n = 103$), those without employment but not in job seeking ($n = 54$) and those in other situations ($n = 383$)).

Included subjects had complete data regarding the variables of interest (S1 Fig). We had missing data for the other variables (from 0.5% to 10.3%), except for age, gender and occupational grade. Assuming a missing at random mechanism, we used stochastic regression imputations rather than complete-case analysis to limit the risk of selection bias [34].

Statistical significance was determined using a two-sided alpha a priori set at 0.05 and analyses were performed with IBM Statistics for Windows, Version 22.0, Released 2013 (Armonk, NY: IBM Corp).

Results

Participants' characteristics

Among the 18,879 participants, 1,129 (6.0%) experienced job loss at one-year. Their characteristics are displayed in Table 1. As expected, job loss was more prevalent among women, participants with low education, low occupational grade, low income, depression, poor self-reported health, job stress, other type of contract than open-ended, part-time job and past-time unemployed.

Associations between alcohol, tobacco and cannabis use and job loss at one-year

When introduced separately, alcohol, tobacco and cannabis use were positively associated with job loss at one-year after adjusting for age, gender, depressive symptoms and self-rated health (Table 2 and S1 Table). We found dose-dependent relationships for alcohol (OR = 1.42(95% CI:1.25,1.62)), tobacco (OR = 1.19(95%CI:1.13,1.26)) and cannabis use (OR = 1.39(95% CI:1.29,1.50)) (Fig 1). We did not find any interaction between alcohol, tobacco and cannabis use and adjustment variables, excepting with depressive state for tobacco use and with age for cannabis use. More precisely, we found an interaction between depressive state and being a heavy smoker; OR = 0.33(95%CI = 0.14,0.76). In stratified analyses according to depressive state, all the associations were significant and in the same direction, except for depressed heavy smoker most probably due to too small sample size (only 8 subjects experienced job loss in this subgroup ($n = 101$), compared to at least 30 in the other subgroups) (S2 Table). We also found an interaction between age more than 50 and having used cannabis more than 12 months ago; OR = 0.63(95%CI:0.42,0.96) and an interaction between age from 30 to 50 and having used cannabis more than 12 months ago; OR = 0.58 (95%CI:0.39,0.86)). In stratified analyses according to age, all the associations were significant and in the same direction, except for consumers less than once month aged over 50, most probably due to too small sample size (only 8 subjects experienced job loss in this subgroup ($n = 74$), compared to at least 30 in the other subgroups) (S3 Table).

When introduced simultaneously, alcohol, tobacco and cannabis use were still positively associated with job loss for alcohol consumers having a dangerous use, for moderate smokers and for all the categories of cannabis consumers after adjustment for age, gender, depressive symptoms and self-reported health (Table 2 and S1 Table). We found dose-dependent relationships for alcohol (OR = 1.20(95%CI:1.05,1.38)), tobacco (OR = 1.08 (95%CI:1.02,1.15)) and cannabis use (OR = 1.30(95%CI:1.20,1.41)). While performing this analysis after

Table 1. Characteristics of the 18,879 included participants from the CONSTANCES cohort and according to their employment status at one-year follow-up.

| N (%) | TOTAL | | EMPLOYED AT ONE-YEAR | | JOB LOSS AT ONE-YEAR | |
|--|---------------|------|----------------------|--------------|----------------------|------|
| | 18,879 (100%) | | 17,750 (94.0%) | | 1,129 (6.0%) | |
| SUBSTANCE USE | N | % | N | % | N | % |
| Alcohol use^a | | | | | | |
| Mild | 16344 | 86.6 | 15438 | 87.0 | 906 | 80.2 |
| Dangerous | 2235 | 11.8 | 2050 | 11.5 | 185 | 16.4 |
| Problematic or dependence | 300 | 1.6 | 262 | 1.5 | 38 | 3.4 |
| Tobacco use^b | | | | | | |
| Non-smokers | 8774 | 46.5 | 8337 | 47.0 | 437 | 38.7 |
| Former smokers | 6374 | 33.8 | 5995 | 33.8 | 379 | 33.6 |
| Light smokers | 2032 | 10.8 | 1871 | 10.5 | 161 | 14.3 |
| Moderate smokers | 1290 | 6.8 | 1177 | 6.6 | 113 | 10.0 |
| Heavy smokers | 409 | 2.2 | 370 | 2.1 | 39 | 3.5 |
| Cannabis use | | | | | | |
| Never used | 9788 | 51.8 | 9327 | 52.5 | 461 | 40.8 |
| No consumption during the previous 12 months | 7513 | 39.8 | 7017 | 39.5 | 496 | 43.9 |
| Less than once a month | 784 | 4.2 | 711 | 4.0 | 73 | 6.5 |
| Once a month or more | 794 | 4.2 | 695 | 4.0 | 99 | 8.8 |
| SOCIODEMOGRAPHIC FACTORS | | | | | | |
| Age (years) | | | | | | |
| Less than 30 | 2245 | 11.9 | 2012 | 11.3 | 233 | 20.6 |
| Between 30 and 50 | 10620 | 56.3 | 10093 | 56.9 | 527 | 46.7 |
| More than 50 | 6014 | 31.9 | 5645 | 31.8 | 369 | 32.7 |
| Gender | | | | | | |
| Men | 8823 | 46.7 | 8340 | 47.0 48.0 | 483 | 42.8 |
| Women | 10056 | 53.3 | 9410 | 53.0 | 646 | 57.2 |
| Education ISCED classification | | | | | | |
| Levels 0 to 4 | 5875 | 31.1 | 5440 | 30.6 | 435 | 38.5 |
| Levels 5 and 6 | 7523 | 39.8 | 7103 | 40.0 | 420 | 37.2 |
| Levels 7 and 8 | 5481 | 29.0 | 5207 | 29.3 | 274 | 24.3 |
| Occupational grade | | | | | | |
| Blue-collar worker, craftsman or employee | 5758 | 30.5 | 5254 | 29.6 | 504 | 44.6 |
| Intermediate worker | 5732 | 30.4 | 5458 | 30.7 | 274 | 24.3 |
| Executive | 7389 | 39.1 | 7038 | 39.7 | 351 | 31.1 |
| Household income (euros per month) | | | | | | |
| Less than 2800 | 5750 | 30.5 | 5201 | 29.3 | 549 | 48.6 |
| More than 2800 | 13129 | 69.5 | 12549 | 70.7 | 580 | 51.4 |
| DEPRESSIVE STATE^c | | | | | | |
| No | 16421 | 87.0 | 15566 | 87.7 | 855 | 75.7 |
| Yes | 2458 | 13.0 | 2184 | 12.3 | 274 | 24.3 |
| SELF-REPORTED HEALTH^d | | | | | | |
| Good | 17116 | 90.7 | 16163 | 91.1 | 953 | 84.4 |
| Poor | 1763 | 9.3 | 1587 | 8.9 | 176 | 15.6 |
| OCCUPATIONAL FACTORS | | | | | | |
| Effort-reward imbalance ratio^e | | | | | | |
| First and second tertiles | 13093 | 69.4 | 12383 | 69.8 | 710 | 62.9 |
| Third tertile | 5786 | 30.6 | 5367 | 30.2 | 419 | 37.1 |

(Continued)

Table 1. (Continued)

| N (%) | TOTAL | | EMPLOYED AT ONE-YEAR | | JOB LOSS AT ONE-YEAR | |
|---|---------------|------|----------------------|------|----------------------|------|
| | 18,879 (100%) | | 17,750 (94.0%) | | 1,129 (6.0%) | |
| SUBSTANCE USE | N | % | N | % | N | % |
| Type of contract | | | | | | |
| Open-ended contract | 15886 | 84.1 | 15123 | 85.2 | 763 | 67.6 |
| Other type of contract | 2993 | 15.9 | 2627 | 14.8 | 366 | 32.4 |
| Type of work time | | | | | | |
| Full-time | 15940 | 84.4 | 15135 | 85.3 | 805 | 71.3 |
| Part-time | 2939 | 15.6 | 261 | 14.7 | 324 | 28.7 |
| Past time unemployed in the last three years | | | | | | |
| No | 17221 | 91.2 | 16426 | 92.5 | 795 | 70.4 |
| Yes | 1658 | 8.8 | 1324 | 7.5 | 334 | 29.6 |

ISCED: 2011 International Standard Classification of Education

^a Categories are defined from Alcohol Use Disorders Identification scores as follows: Mild (0–7), Dangerous (8–15), Problematic (16–19) and Dependence (20–40)

^b Categories of current smokers are defined as follows: Light (1 to 9 cigarettes per day), Moderate (10 to 19) and Heavy (>19) consumers

^c A total score at the Center of Epidemiologic Studies Depression scale ≥ 19 defined a depressive state

^d Self-reported health was used as a binary variable from an 8-points Likert scale

^e Computed from 7 items regarding rewards and from 3 items regarding efforts and with all the items assessed on a 4-points Likert scale.

<https://doi.org/10.1371/journal.pone.0222361.t001>

aggregating the two last categories of alcohol and tobacco use, we found significant associations between these categories and job loss and still with dose-dependent relationships for tobacco use for which there were more than 2 categories left (S4 Table).

Table 2. Associations between alcohol, tobacco and cannabis use and job loss at one-year among 18,879 participants from the CONSTANCES cohort, adjusting for age, gender, self-reported health and depressive symptoms.

| Type of model | Each substance successively entered | | | | All substances simultaneously entered | | | |
|-------------------------------------|-------------------------------------|-------------|-------------|------------------|---------------------------------------|-------------|-------------|------------------|
| | OR | 95%CI | | p value | OR | 95%CI | | p value |
| Alcohol use^a | | | | | | | | |
| Dangerous | 1.46 | 1.23 | 1.73 | <0.001 | 1.22 | 1.02 | 1.46 | 0.030 |
| Problematic or Dependence | 1.92 | 1.34 | 2.75 | <0.001 | 1.42 | 0.98 | 2.05 | 0.067 |
| Tobacco use^b | | | | | | | | |
| Former smoker | 1.26 | 1.09 | 1.46 | 0.002 | 1.08 | 0.93 | 1.26 | 0.320 |
| Light smoker | 1.54 | 1.27 | 1.86 | <0.001 | 1.17 | 0.96 | 1.44 | 0.128 |
| Moderate smoker | 1.69 | 1.36 | 2.10 | <0.001 | 1.29 | 1.02 | 1.63 | 0.034 |
| Heavy smoker | 1.78 | 1.26 | 2.54 | 0.001 | 1.33 | 0.92 | 1.92 | 0.126 |
| Cannabis use^c | | | | | | | | |
| Consumption more than 12 months ago | 1.45 | 1.27 | 1.66 | <0.001 | 1.35 | 1.17 | 1.56 | <0.001 |
| Less than once a month | 1.87 | 1.43 | 2.45 | <0.001 | 1.62 | 1.22 | 2.15 | 0.001 |
| Once a month or more | 2.68 | 2.10 | 3.42 | <0.001 | 2.19 | 1.68 | 2.87 | <0.001 |

OR: Odds ratios; 95%CI: Confidence interval at 95%

^a Categories are defined from Alcohol Use Disorders Identification scores as follows: Mild (0–7), Dangerous (8–15), Problematic (16–19) and Dependence (20–40), with Mild category as reference

^b Categories of current smokers are defined as follows: Light (1 to 9 cigarettes per day), Moderate (10 to 19) and Heavy (>19) consumers, with never smokers as reference category

^c Reference category is never use. Adjustments variables were as follows: gender, age in three categories (<30; ≥ 30 and <50 and ≥ 50), self-reported health was used as a binary variable from an 8-points Likert scale, and depressive state defined as a total score ≥ 19 at the Center for Epidemiologic Studies Depression (CESD). Significant associations are presented in bold (i.e. $p < 0.05$).

<https://doi.org/10.1371/journal.pone.0222361.t002>




Fig 1. Risk of job loss at one-year according to substance use, adjusting for age, gender, self-reported health and depressive symptoms, among 18,879 participants from the CONSTANCES cohort.

<https://doi.org/10.1371/journal.pone.0222361.g001>

We found no interactions between pairs of substances: OR = 0.74 (95%CI:0.51,1.07) with $p = 0.113$ for the interaction between alcohol and cannabis use, OR = 1.09(95%CI:0.79,1.51) with $p = 0.589$ for the interaction between alcohol and tobacco use, and OR = 1.09(95%CI:0.75,1.58) with $p = 0.664$ for the interaction between cannabis and tobacco use. Odds ratios of job loss while using substance use as binary variables are presented in [S5 Table](#).

Stratifications for sociodemographic and occupational factors

In stratified analyses for sociodemographic and occupational factors, all the associations between substance use and job loss were positive ([Table 3](#)). Over a total of 27 stratifications, significant associations persisted in all strata among 20 of them. Regarding alcohol use, the only sub-groups in which the associations were not significant were the youngest (OR = 1.30 (95%CI:0.95,1.79)), those with low income (OR = 1.18(0.94,1.49)), those at part-time (OR = 1.34(95%CI:0.94,1.91)) and those with history of unemployment (OR = 1.00(95%CI:0.73, 1.38)). These associations were significant in all these sub-groups for the two other substances (i.e. tobacco and cannabis). Regarding tobacco use, the only sub-groups in which the associations were not significant were the oldest (OR = 1.19(95%CI:0.90,1.57)), those with the highest occupational grade (OR = 1.20(95%CI:0.92,1.57)) and those with the highest income (OR = 1.23(95%CI:0.99,1.52)). These associations were significant in all these sub-groups for the two other substances (i.e. alcohol and cannabis). Regarding cannabis use, associations were significant in all strata for all the stratifications.

Sensitivity analysis

In generalized estimating equations over a three-year duration of follow-up, all the associations between substance use and job loss remained positive and with similar effects sizes ([S6 Table](#)). We found no time by substance interaction for alcohol (OR = 0.98(95%CI:0.80,1.19); $p = 0.827$), nor for tobacco (OR = 0.99(95%CI:0.83,1.18); $p = 0.886$) and nor for cannabis use (OR = 0.86(95%CI:0.69,1.08); $p = 0.185$).

Associations between alcohol, tobacco and cannabis use and subsequent job loss persist after adjustment for all the covariables ([S7 Table](#)).

Table 3. Associations between alcohol, tobacco and cannabis use and job loss at one-year among 18,879 participants from the CONSTANCES cohort, adjusting for age, gender, self-reported health and depressive symptoms, and while stratifying for sociodemographic and occupational factors.

| SOCIODEMOGRAPHIC FACTORS | Type of substance use | | | | | | | | | | | |
|--|-----------------------|--------------|----------------|----------------------|-------------|--------------|-----------------------|------------------|-------------|--------------|----------------|------------------|
| | Alcohol ^a | | | Tobacco ^b | | | Cannabis ^c | | | | | |
| | OR | 95%CI | p value | OR | 95%CI | p value | OR | 95%CI | p value | | | |
| Age (in years) | | | | | | | | | | | | |
| Less than 30 | 1.30 | 0.95 | 1.79 | 0.105 | 1.86 | 1.40 | 2.47 | <0.001 | 1.51 | 1.11 | 2.04 | 0.008 |
| Between 30 and 50 | 1.48 | 1.17 | 1.89 | 0.001 | 1.47 | 1.21 | 1.79 | <0.001 | 1.99 | 1.55 | 2.56 | <0.001 |
| More than 50 | 1.82 | 1.36 | 2.44 | <0.001 | 1.19 | 0.90 | 1.57 | 0.224 | 2.84 | 1.73 | 4.66 | <0.001 |
| Gender | | | | | | | | | | | | |
| Men | 1.55 | 1.26 | 1.91 | <0.001 | 1.71 | 1.39 | 2.10 | <0.001 | 1.96 | 1.52 | 2.53 | <0.001 |
| Women | 1.45 | 1.12 | 1.87 | 0.005 | 1.30 | 1.08 | 1.57 | 0.006 | 1.78 | 1.37 | 2.31 | <0.001 |
| Education level (2011 ISCED classification) | | | | | | | | | | | | |
| Levels 0 to 4 | 1.45 | 1.12 | 1.87 | 0.004 | 1.51 | 1.22 | 1.88 | <0.001 | 1.82 | 1.33 | 2.50 | <0.001 |
| Levels 5 and 6 | 1.44 | 1.09 | 1.91 | 0.011 | 1.33 | 1.06 | 1.68 | 0.015 | 1.94 | 1.44 | 2.62 | <0.001 |
| Levels 7 and 8 | 1.64 | 1.20 | 2.24 | 0.002 | 1.41 | 1.05 | 1.90 | 0.023 | 1.79 | 1.27 | 2.53 | 0.001 |
| Occupational grade | | | | | | | | | | | | |
| Blue-collar worker, craftsman or employee | 1.37 | 1.07 | 1.76 | 0.013 | 1.52 | 1.25 | 1.86 | <0.001 | 1.78 | 1.35 | 2.36 | <0.001 |
| Intermediate worker | 1.43 | 1.02 | 2.00 | 0.040 | 1.45 | 1.09 | 1.92 | 0.010 | 1.50 | 1.01 | 2.24 | 0.045 |
| Executive | 1.76 | 1.34 | 2.31 | <0.001 | 1.20 | 0.92 | 1.57 | 0.187 | 2.30 | 1.69 | 3.13 | <0.001 |
| Household income(in euros per month) | | | | | | | | | | | | |
| Less than 2800 | 1.18 | 0.94 | 1.49 | 0.164 | 1.54 | 1.27 | 1.85 | <0.001 | 1.63 | 1.28 | 2.07 | <0.001 |
| More than 2800 | 1.82 | 1.46 | 2.27 | <0.001 | 1.23 | 0.99 | 1.52 | 0.054 | 1.92 | 1.45 | 2.55 | <0.001 |
| OCCUPATIONAL FACTORS | OR | 95%CI | p value | | OR | 95%CI | p value | | OR | 95%CI | p value | |
| Effort-reward imbalance ratio^d | | | | | | | | | | | | |
| First and second tertiles | 1.45 | 1.18 | 1.78 | <0.001 | 1.45 | 1.21 | 1.72 | <0.001 | 1.85 | 1.48 | 2.32 | <0.001 |
| Third tertile | 1.61 | 1.24 | 2.09 | <0.001 | 1.50 | 1.20 | 1.88 | <0.001 | 1.87 | 1.37 | 2.55 | <0.001 |
| Type of contract | | | | | | | | | | | | |
| Open-ended contract | 1.50 | 1.12 | 1.82 | <0.001 | 1.43 | 1.20 | 1.69 | <0.001 | 1.77 | 1.40 | 2.25 | <0.001 |
| Other type of contract | 1.47 | 1.10 | 1.96 | 0.010 | 1.51 | 1.19 | 1.93 | 0.001 | 1.57 | 1.18 | 2.10 | 0.002 |
| Type of work time | | | | | | | | | | | | |
| Full-time | 1.56 | 1.30 | 1.87 | <0.001 | 1.47 | 1.25 | 1.72 | <0.001 | 1.82 | 1.47 | 2.24 | <0.001 |
| Part-time | 1.34 | 0.94 | 1.91 | 0.105 | 1.50 | 1.16 | 2.01 | 0.002 | 1.71 | 1.18 | 2.49 | 0.005 |
| History of unemployment during the last three years | | | | | | | | | | | | |
| No | 1.63 | 1.35 | 1.97 | <0.001 | 1.34 | 1.13 | 1.59 | 0.001 | 1.73 | 1.38 | 2.17 | <0.001 |
| Yes | 1.00 | 0.73 | 1.38 | 0.990 | 1.37 | 1.05 | 1.79 | 0.019 | 1.59 | 1.15 | 2.20 | 0.006 |

OR: Odd ratio; 95%CI: Confidence interval at 95%; ISCED: 2011 International Standard Classification of Education

^a Categories are defined from Alcohol Use Disorders Identification scores as follows: Mild (0–7), Dangerous, Problematic and Dependence (8–40), with Mild category as reference

^b Non-smokers is defined as reference category compared to former and current smokers

^c Never used is defined as reference category compared to having already used

^d Computed from 7 items regarding rewards and from 3 items regarding efforts and with all the items assessed on a 4-points likert scale. Adjustments variables were as follows: gender, age in three categories (<30; ≥30 and <50 and ≥50), self-reported health was used as a binary variable from an 8-points Likert scale, and depressive state defined as a total score ≥19 at the Center for Epidemiologic Studies Depression (CESD). Significant associations are presented in bold (i.e. p<0.05).

<https://doi.org/10.1371/journal.pone.0222361.t003>

Finally, we excluded successively, and then simultaneously, the three types of unemployed participants at follow-up having another status than job seeking. All these analyses provided similar results.

Discussion

Summary of the results

Our main objective was to examine the prospective associations between alcohol, tobacco, cannabis use and their combination with job loss at one-year, among a large population-based sample of men and women, while adjusting for age, gender, self-reported health and depressive symptoms. We found that alcohol, tobacco and cannabis use were independently associated with job loss at one-year, showing dose-dependent relationships. We further sought to examine those associations while stratifying for a broad range of sociodemographic and occupational factors previously linked to job loss. Since the associations remained significant across almost all stratifications, all sociodemographic and occupational positions are potentially concerned. Finally, associations remained positive and with similar effect sizes over a three-year follow-up.

Strengths and limitations

To our knowledge, this is the first study to examine the prospective associations of each substance use and their combination and subsequent job loss in the short term in a large national population-based cohort. Over the inclusion period (i.e. 2012 to 2016), the unemployment rate at baseline among the 81,997 participants from the CONSTANCES cohort (11%, including 6.1% who are actively seeking employment) is close to the mean unemployment rate in France (10.2%) while taking the definition of the international labour office [35]. In addition, sociodemographic and occupational factors associated with job loss in CONSTANCES were similar to those described in the general population [5,6,15,16,25–27,36]. Thanks to the large sample, analyses were stratified for several individual and occupational factors to examine whether these associations were specific to subgroups or rather a general phenomenon. In particular, job stress was assessed with a well-validated standardized tool. Finally, all our analyses were adjusted for potential confounders including age, gender, self-reported health and depressive symptoms.

Our study has several limitations. Firstly, even if the CONSTANCES cohort randomly recruited its participants, this population is not representative of the general population due to selection effects associated with voluntary participation. Thus, our findings may not apply to the same extent to other settings. However, the associations were consistent in most stratified analyses, suggesting that potential selection biases related to these sociodemographic and occupational factors may be unlikely to substantially affect our conclusions. Secondly, the listwise deletion of individuals who had missing data for the main variables of interest led to a decrease in statistical power and, potentially, the selection of subjects less likely to display severe substance use or to lose their job. In addition, depression and overall health status might act not only as confounders but also as mediators. Therefore our results might have underestimated the strength of the associations. This issue might explain the lack of significance for some categories of the exposure with small sample. For instance, while introducing simultaneously in the same model the three substances, problematic alcohol use or dependence, as well as heavy smoking, were no longer associated with the risk of job loss. Noteworthy, dose-dependent relationships were found for the three substances in all the models and there were again significant associations after aggregating these categories with the lower category of use. Furthermore, compared with models including only one substance, simultaneously including the three substances did not affect much the association of cannabis with job loss (mean decrease of the odds ratios of 13%). In contrast, the decreases of odds ratios for alcohol and tobacco use were higher but of similar magnitude. Specifically, the decreases were of 21% and of 22% for alcohol

and tobacco use, respectively. This suggests that the association of tobacco with job loss might be partially confounded by the use of the two other substances, but not to a greater extent than alcohol. Thirdly, the prospective design did not rule out the possibility of reverse causality. However, although some participants who experienced job loss at one-year might have anticipated this condition, similar results were found in analyses over a three-year follow-up.

Explanatory hypotheses

Several explanatory hypotheses can be proposed concerning the links between substance use and subsequent job loss. Regarding alcohol use, acute effects rarely go unnoticed, such as disinhibition following intoxication, or tremors, sweats and irritability following withdrawal. In addition, chronic alcohol use may induce long-term neuropsychological impairment such as memory, concentration and attention disorders. All these symptoms may decrease the ability to perform work-related tasks. Regarding tobacco use, smoking is associated with psychological distress including anxiety and irritability which could contribute to a decreased productivity [37]. Moreover, smoking is time-consuming. Regarding the association between former smoking and job loss, they might be explained, at least partially, by the potential increased vulnerability of former smokers to environmental stressors. Anxiety, which is associated with loss productivity [38], has been found to be more prevalent in former smokers than in non-smokers, even after long periods of cessation [39]. Regarding cannabis, its use is associated with symptoms that could substantially interfere with work tasks, such as attentional disorders and persecution delusions [40]. Physical symptoms are also particularly noticeable for the entourage such as the hyperemesis syndrome. Chronic use may also lead to amotivational syndrome and impaired decision-making [41].

Public health implications and future research

As regards information, important messages can be delivered to the population; for instance even moderate levels of alcohol, tobacco or cannabis use are associated with job loss and all sociodemographic and occupational positions are potentially concerned. As regards prevention, measures could be taken within the companies to limit the risks of substance use, for example by no longer offering alcohol in the canteen, and by extending the smoking ban to the whole company (e.g. outdoor work). Preventive and occupational medicine could also play a substantial role by screening at-risk subjects to prevent substance use and its detrimental consequences on occupational life. Moreover, informing on these potential consequences might be a promising motivational tool. Importantly, being excluded from employment is associated with a significant deterioration in physical and psychological health, and substance use makes it more difficult to return to employment. Since unemployment concern a large panel of the general population, from a public health perspective it is of paramount interest to prevent entry into this vicious circle. Future studies should examine which types of jobs might be more concerned than others. Finally, future studies should test whether interventions aiming at preventing or decreasing substance use improve job retention.

Supporting information

S1 Fig. Flow chart.
(DOCX)

S1 Table. Associations between alcohol, tobacco and cannabis use and job loss at one-year among 18,879 participants from the CONSTANCES cohort, adjusting for age, gender,

self-reported health and depressive symptoms.
(DOCX)

S2 Table. Associations between tobacco use and job loss at one-year among 18,879 participants from the CONSTANCES cohort, adjusting for age, gender and self-reported health while stratifying for depressive state.

(DOCX)

S3 Table. Associations between cannabis use and job loss at one-year among 18,879 participants from the CONSTANCES cohort, adjusting for gender, depressive state and self-reported health while stratifying for age.

(DOCX)

S4 Table. Associations between alcohol, tobacco and cannabis use and job loss at one-year among 18,879 participants from the CONSTANCES cohort, adjusting for age, gender, self-reported health and depressive symptoms, with all substances simultaneously entered and while aggregating “dangerous” and “problematic or dependence” categories regarding alcohol use, as well as the “moderate” and “heavy” categories regarding tobacco use.

(DOCX)

S5 Table. Associations between alcohol, tobacco and cannabis use and job loss at one-year among 18,879 participants from the CONSTANCES cohort, adjusting for age, gender, self-reported health and depressive symptoms while using substances as binary variables.

(DOCX)

S6 Table. Associations between alcohol, tobacco and cannabis use and job loss over a three-year follow-up among 18,879 participants from the CONSTANCES cohort, adjusting for age, gender, poor self-reported health and depressive state.

(DOCX)

S7 Table. Associations between alcohol, tobacco and cannabis use and job loss at one-year among 18,225 participants from the CONSTANCES cohort, adjusting for age, gender, education level, occupational grade, household income, effort-reward imbalance ratio, type of contract, type of work time, past time unemployed in the past three years, depressive symptoms and self-reported health.

(DOCX)

Author Contributions

Conceptualization: Guillaume Airagnes, Cédric Lemogne.

Data curation: Marcel Goldberg, Marie Zins.

Formal analysis: Guillaume Airagnes.

Funding acquisition: Marcel Goldberg, Marie Zins.

Methodology: Guillaume Airagnes, Cédric Lemogne, Pierre Meneton, Nicolas Hoertel, Marie Zins.

Project administration: Marcel Goldberg, Marie Zins.

Resources: Marie Zins.

Supervision: Marcel Goldberg, Frédéric Limosin, Marie Zins.

Validation: Marcel Goldberg, Yves Roquelaure, Frédéric Limosin, Marie Zins.

Visualization: Marie Plessz, Marcel Goldberg, Nicolas Hoertel, Yves Roquelaure, Frédéric Limosin, Marie Zins.

Writing – original draft: Guillaume Airagnes.

Writing – review & editing: Cédric Lemogne, Pierre Meneton, Marie Plessz, Marcel Goldberg, Nicolas Hoertel, Yves Roquelaure, Frédéric Limosin, Marie Zins.

References

1. Rehm J, Mathers C, Popova S, Thavorncharoensap M, Teerawattananon Y, Patra J (2009) Global burden of disease and injury and economic cost attributable to alcohol use and alcohol-use disorders. *Lancet* 373: 2223–2233. [https://doi.org/10.1016/S0140-6736\(09\)60746-7](https://doi.org/10.1016/S0140-6736(09)60746-7) PMID: 19560604
2. Shield KD, Rehm MX, Rehm J (2015) Social costs of addiction in Europe. Impact of Addictive Substances and Behaviours on Individual and Societal Well-being: 181–188.
3. WHO (2009) Global Health Risks: Mortality and Burden of Disease Attributable to Selected Major Risks: World Health Organization.
4. Eurostat Février 2018. Le taux de chômage à 8,5% dans la zone euro. Communiqué de presse euroindicateurs. 4 avril 2018. <http://ec.europa.eu/eurostat/documents/2995521/8782909/3-04042018-BP-FR.pdf/df516d72-780a-4db4-9d19-c18edb68f928>.
5. Henkel D (2011) Unemployment and Substance Use: A Review of the Literature (1990–2010). *Current Drug Abuse Reviews* 4: 4–27. PMID: 21466502
6. Compton WM, Gfroerer J, Conway KP, Finger MS (2014) Unemployment and substance outcomes in the United States 2002–2010. *Drug & Alcohol Dependence* 142: 350–353.
7. Eliason M (2014) Alcohol-related morbidity and mortality following involuntary job loss: evidence from Swedish register data. *J Stud Alcohol Drugs* 75: 35–46. <https://doi.org/10.15288/jsad.2014.75.35> PMID: 24411795
8. Eliason M, Storrie D (2009) Job loss is bad for your health—Swedish evidence on cause-specific hospitalization following involuntary job loss. *Social Science & Medicine* 68: 1396–1406.
9. Eliason M, Storrie D (2009) Does Job Loss Shorten Life? *Journal of Human Resources* 44: 277–302.
10. Kaspersen SL, Pape K, Vie GÅ, Ose SO, Krokstad S, Gunnell D, et al. (2016) Health and unemployment: 14 years of follow-up on job loss in the Norwegian HUNT Study. *European Journal of Public Health* 26: 312–317. <https://doi.org/10.1093/eurpub/ckv224> PMID: 26715474
11. Franco S, Olfson M, Wall MM, Wang S, Hoertel N, Blanco C (2019) Shared and specific associations of substance use disorders on adverse outcomes: A national prospective study. *Drug and Alcohol Dependence* 201: 212–219. <https://doi.org/10.1016/j.drugalcdep.2019.03.003> PMID: 31252355
12. Blanco C, Wall MM, Hoertel N, Krueger RF, Liu S-M, Grant BF, et al. (2019) Psychiatric disorders and risk for multiple adverse outcomes: a national prospective study. *Molecular psychiatry*.
13. Booth BM, Feng W (2002) The impact of drinking and drinking consequences on short-term employment outcomes in at-risk drinkers in six southern states. *The Journal of Behavioral Health Services & Research* 29: 157–166.
14. Leino-Arjas P, Liira J, Mutanen P, Malmivaara A, Matikainen E (1999) Predictors and consequences of unemployment among construction workers: prospective cohort study. *BMJ: British Medical Journal* 319: 600–605. <https://doi.org/10.1136/bmj.319.7210.600> PMID: 10473472
15. Jusot F, Khlal M, Rochereau T, Serme C (2008) Job loss from poor health, smoking and obesity: a national prospective survey in France. *Journal of Epidemiology and Community Health* 62: 332–337. <https://doi.org/10.1136/jech.2007.060772> PMID: 18339826
16. Zhang C, Brook JS, Leukefeld CG, Brook DW (2016) Trajectories of marijuana use from adolescence to adulthood as predictors of unemployment status in the early forties. *The American Journal on Addictions* 25: 203–209. <https://doi.org/10.1111/ajad.12361> PMID: 26991779
17. Danielsson A-K, Falkstedt D, Hemmingsson T, Allebeck P, Agardh E (2015) Cannabis use among Swedish men in adolescence and the risk of adverse life course outcomes: results from a 20 year-follow-up study. *Addiction* 110: 1794–1802. <https://doi.org/10.1111/add.13042> PMID: 26172111
18. Zuelke AE, Luck T, Schroeter ML, Witte AV, Hinz A, Engel C, et al. (2018) The association between unemployment and depression—Results from the population-based LIFE-adult-study. *J Affect Disord* 235: 399–406. <https://doi.org/10.1016/j.jad.2018.04.073> PMID: 29677604

19. Wanberg CR, Kanfer R, Hamann DJ, Zhang Z (2016) Age and reemployment success after job loss: An integrative model and meta-analysis. *Psychol Bull* 142: 400–426. <https://doi.org/10.1037/bul0000019> PMID: 26011790
20. Alavinia SM, Burdorf A (2008) Unemployment and retirement and ill-health: a cross-sectional analysis across European countries. *Int Arch Occup Environ Health* 82: 39–45. <https://doi.org/10.1007/s00420-008-0304-6> PMID: 18264715
21. Whooley MA, Kiefe CI, Chesney MA, Markovitz JH, Matthews K, Hulley SB (2002) Depressive symptoms, unemployment, and loss of income: The CARDIA Study. *Archives of Internal Medicine* 162: 2614–2620. <https://doi.org/10.1001/archinte.162.22.2614> PMID: 12456234
22. Åhs A, Westerling R (2005) Self-rated health in relation to employment status during periods of high and of low levels of unemployment. *The European Journal of Public Health* 16: 294–304.
23. Giatti L, Barreto SM, César CC (2010) Unemployment and self-rated health: Neighborhood influence. *Social Science & Medicine* 71: 815–823.
24. Covizzi I (2008) Does Union Dissolution Lead to Unemployment? A Longitudinal Study of Health and Risk of Unemployment for Women and Men Undergoing Separation. *European Sociological Review* 24: 347–361.
25. Dooley D, Catalano R, Wilson G (1994) Depression and unemployment: panel findings from the Epidemiologic Catchment Area study. *American journal of community psychology* 22: 745–765. PMID: 7639201
26. Siegrist J, Li J (2016) Associations of Extrinsic and Intrinsic Components of Work Stress with Health: A Systematic Review of Evidence on the Effort-Reward Imbalance Model. *Int J Environ Res Public Health* 13: 432. <https://doi.org/10.3390/ijerph13040432> PMID: 27104548
27. Fullerton AS, Long MA, Anderson KF (2016) Job Insecurity and Substance Use in the United States: Stress, Strain, and the Gendering of Precarious Employment. *Research in the Sociology of Work: Emerald Group Publishing Limited*. pp. 241–271.
28. Goldberg M, Carton M, Descatha A, Leclerc A, Roquelaure Y, Santin G, et al. (2017) CONSTANCES: a general prospective population-based cohort for occupational and environmental epidemiology: cohort profile. *Occup Environ Med* 74: 66–71. <https://doi.org/10.1136/oemed-2016-103678> PMID: 27884936
29. Gache P, Michaud P, Landry U, Accietto C, Arfaoui S, Wenger O, et al. (2005) The Alcohol Use Disorders Identification Test (AUDIT) as a screening tool for excessive drinking in primary care: reliability and validity of a French version. *Alcohol Clin Exp Res* 29: 2001–2007. <https://doi.org/10.1097/01.alc.0000187034.58955.64> PMID: 16340457
30. Lohse T, Rohrmann S, Bopp M, Faeh D (2016) Heavy Smoking Is More Strongly Associated with General Unhealthy Lifestyle than Obesity and Underweight. *PLoS One* 11: e0148563. <https://doi.org/10.1371/journal.pone.0148563> PMID: 26910775
31. DeSalvo KB, Bloser N, Reynolds K, He J, Muntner P (2006) Mortality prediction with a single general self-rated health question. *J Gen Intern Med* 21: 267. <https://doi.org/10.1111/j.1525-1497.2005.00291.x> PMID: 16336622
32. Morin AJ, Moullec G, Maiano C, Layet L, Just JL, Ninot G (2011) Psychometric properties of the Center for Epidemiologic Studies Depression Scale (CES-D) in French clinical and nonclinical adults. *Rev Epidemiol Sante Publique* 59: 327–340. <https://doi.org/10.1016/j.respe.2011.03.061> PMID: 21925817
33. Schneider SL (2013) The International Standard Classification of Education 2011. In: Birkelund GE, editor. *Class and Stratification Analysis (Comparative Social Research, Volume 30)* Emerald Group Publishing Limited. pp. 365–379.
34. Haukoos JS, Newgard CD (2007) Advanced Statistics: Missing Data in Clinical Research—Part 1: An Introduction and Conceptual Framework. *Academic Emergency Medicine* 14: 662–668. <https://doi.org/10.1197/j.aem.2006.11.037> PMID: 17538078
35. Insee Références, édition 2018—Fiches—Chômage. <https://www.insee.fr/fr/statistiques/fichier/3573767/ecrt18-f2-1-chomage.pdf>.
36. Schunck R, Rogge BG (2012) Unemployment and Smoking: Causation, Selection, or Common Cause?—Evidence from Longitudinal Data. *SOEPpaper* 491.
37. Halpern MT, Shikier R, Rentz AM, Khan ZM (2001) Impact of smoking status on workplace absenteeism and productivity. *Tobacco control* 10: 233–238. <https://doi.org/10.1136/tc.10.3.233> PMID: 11544387
38. Greenberg PE, Sisitsky T, Kessler RC, Finkelstein SN, Berndt ER, Davidson JR, et al. (1999) The economic burden of anxiety disorders in the 1990s. *The Journal of clinical psychiatry* 60: 427–435. <https://doi.org/10.4088/jcp.v60n0702> PMID: 10453795

39. Mykletun A, Overland S, Aarø LE, Liabø H-M, Stewart R Smoking in relation to anxiety and depression: Evidence from a large population survey: The HUNT study. *European Psychiatry* 23: 77–84. <https://doi.org/10.1016/j.eurpsy.2007.10.005> PMID: 18082377
40. Karila L, Roux P, Rolland B, Benyamina A, Reynaud M, Aubin HJ, et al. (2014) Acute and long-term effects of cannabis use: a review. *Curr Pharm Des* 20: 4112–4118. <https://doi.org/10.2174/13816128113199990620> PMID: 24001294
41. Lac A, Luk JW (2018) Testing the Amotivational Syndrome: Marijuana Use Longitudinally Predicts Lower Self-Efficacy Even After Controlling for Demographics, Personality, and Alcohol and Cigarette Use. *Prevention Science* 19: 117–126. <https://doi.org/10.1007/s11121-017-0811-3> PMID: 28620722