

Epidemiology and antimicrobial resistance of methicillin-resistant *Staphylococcus aureus* isolates colonizing pigs with different exposure to antibiotics

Elizeth Lopes, Teresa Conceição, Laurent Poirel, Hermínia de Lencastre, Marta Aires-De-Sousa

► **To cite this version:**

Elizeth Lopes, Teresa Conceição, Laurent Poirel, Hermínia de Lencastre, Marta Aires-De-Sousa. Epidemiology and antimicrobial resistance of methicillin-resistant *Staphylococcus aureus* isolates colonizing pigs with different exposure to antibiotics. PLoS ONE, Public Library of Science, 2019, 14 (11), pp.e0225497. 10.1371/journal.pone.0225497. inserm-02409006

HAL Id: inserm-02409006

<https://www.hal.inserm.fr/inserm-02409006>

Submitted on 13 Dec 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH ARTICLE

Epidemiology and antimicrobial resistance of methicillin-resistant *Staphylococcus aureus* isolates colonizing pigs with different exposure to antibiotics

Elizeth Lopes¹, Teresa Conceição¹, Laurent Poirel^{2,3,4}, Hermínia de Lencastre^{1,5}, Marta Aires-de-Sousa^{1,6*}

1 Laboratory of Molecular Genetics, Instituto de Tecnologia Química e Biológica António Xavier, Universidade Nova de Lisboa, Oeiras, Portugal, **2** Emerging Antibiotic Resistance Unit, Medical and Molecular Microbiology, Department of Medicine, University of Fribourg, Fribourg, Switzerland, **3** French INSERM European Unit, University of Fribourg (LEA-IAME), Fribourg, Switzerland, **4** National Reference Center for Emerging Antibiotic Resistance, Fribourg, Switzerland, **5** Laboratory of Microbiology and Infectious Diseases, The Rockefeller University, New York, New York, United States of America, **6** Escola Superior de Saúde da Cruz Vermelha Portuguesa, Lisbon, Portugal

* msousa@esscvp.eu



OPEN ACCESS

Citation: Lopes E, Conceição T, Poirel L, de Lencastre H, Aires-de-Sousa M (2019) Epidemiology and antimicrobial resistance of methicillin-resistant *Staphylococcus aureus* isolates colonizing pigs with different exposure to antibiotics. PLoS ONE 14(11): e0225497. <https://doi.org/10.1371/journal.pone.0225497>

Editor: Tara C. Smith, Kent State University, UNITED STATES

Received: September 6, 2019

Accepted: November 6, 2019

Published: November 20, 2019

Copyright: © 2019 Lopes et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

Funding: This work was partly supported by project PTDC/DTP-EPI/0842/2014 from Fundação para a Ciência e a Tecnologia (FCT), Portugal, and Project LISBOA-01-0145-FEDER-007660 (Microbiologia Molecular, Estrutural e Celular) funded by FEDER funds through COMPETE2020 - Programa Operacional Competitividade e

Abstract

Background

In 2016, very high rates of methicillin-resistant *Staphylococcus aureus* (MRSA)-ST398 (99%) were found in Portuguese pig farms that used colistin, amoxicillin, and zinc oxide as feed additives. Since then, farms A and B banned the use of colistin, and farm C banned the use of both antibiotics.

Objective

The aim of the present study was to evaluate the impact of the ban of colistin and amoxicillin on pig MRSA carriage rates, clonal types and antimicrobial resistance, compared to the results obtained in 2016.

Methods

In 2018, 103 pigs (52 from farm B using amoxicillin only as a feed additive and 51 from farm C where no antibiotics were included in the feed regimen) were nasally swabbed for MRSA colonization. Isolates were tested for antimicrobial susceptibility, and characterised by *spa* typing, SCC*mec* typing and MLST. Whole genome sequencing (WGS) was performed for representative isolates.

Results

Overall, 96% of the pigs swabbed in 2018 carried MRSA, mostly ST398-SCC*mec* V-*spa* types t011/t108. MRSA from pigs not receiving antibiotics in the feed regimen showed susceptibility to a higher number of antibiotics, namely erythromycin, ciprofloxacin, gentamicin,

Internacionalização (POCI) and by national funds through FCT. This work was also partially supported by ONEIDA project (LISBOA-01-0145-FEDER-016417) co-funded by FEEL - "Fundos Europeus Estruturais e de Investimento" from "Programa Operacional Regional Lisboa 2020" and by national funds from FCT. Elizeth Lopes was supported by grant 03/BI/2017 from FCT, Portugal.

Competing interests: The authors have declared that no competing interests exist.

and chloramphenicol. Notably, most of these isolates ($n = 52$) presented an unusual erythromycin-susceptibility/clindamycin-resistance phenotype. WGS showed that these isolates lacked the *erm* and the *lnu* genes encoding resistance to macrolides and lincosamides, respectively, but carried the *vgaA_{LC}* gene encoding resistance to lincosamides, which is here firstly identified in *S. aureus* ST398.

Conclusion

After two years the ban of colistin and amoxicillin as feed additives had no significant impact on the MRSA nasal carriage rates. Nevertheless, the MRSA strains circulating in those farms showed resistance to a lower number of antibiotic classes.

Introduction

There is increasing concern about the use of antibiotics in food-producing animals that may lead to elevated resistance rates, and therefore ultimately impact the treatment of human infections. By consequence, several countries in the European Union have made efforts to reduce the use of antibiotics in livestock, in particular limiting their use as growth promoters and prophylaxis in healthy animals.

Livestock-associated methicillin-resistant *Staphylococcus aureus* (LA-MRSA) have been widely reported as nasal colonizers of pigs in many geographical areas [1]. Porcine MRSA in Europe and the United States mainly belong to clonal complex (CC)398 [1]. In 2016, we evaluated the occurrence of MRSA isolates in two pig farms in Portugal (farms A and B) that supplemented the feed regimen of the animals with colistin and amoxicillin. Very high rates of MRSA (99%) were found in both farms, and all strains belonged to ST398 [2]. Since then, as a consequence of the Portuguese national action plan for the reduction of the use of antibiotics in animals [3], several farms stopped feeding the pigs with colistin-supplemented regimens, including farms A and B, and some farms completely abolished the routine use of any antibiotic prophylaxis.

Although there is clear evidence about the relationship between a high antimicrobial usage in pig farms and the increased rates of Gram-negative resistant bacteria in their digestive tract [4–7], there is no study evaluating the impact of the use of antimicrobials on the nasal carriage of multidrug-resistant MRSA in pigs.

The aim of the present study was to evaluate the impact of the ban of colistin and amoxicillin from the feed regimens of healthy pigs on MRSA carriage rates, MRSA clonal types and antimicrobial resistance, compared to the isolates obtained in 2016 from animals receiving both antibiotics.

Materials and methods

Farms and study design

Two independent Portuguese pig farms (farms B and C), all located in the Alentejo region, were included in the study. All pigs are born in these farms and further delivered to slaughterhouses. The two farms used amoxicillin (0.5%), colistin (0.5%), and zinc oxide (0.15%) in the feed regimen of all animals until 2016. Since then, farm B banned colistin from the feed regimen, maintaining amoxicillin (0.5%) and zinc oxide (0.15%), while farm C did not use either antibiotic, keeping zinc oxide (0.15%) as a feed supplement for the prevention of gastrointestinal diseases. No other feed additives were included in the regular feed regimen at any time.




Fig 1. Diagram representing the three groups of pigs, considering the sampling period, the farm, and the feed regimen (amoxicillin, colistin, and zinc oxide).

<https://doi.org/10.1371/journal.pone.0225497.g001>

However, the two farms administered tetracycline in the feed regimens of all animals whenever more than 10% of the pigs developed a gastrointestinal infection.

A total of 154 piglets, aged 10–12 weeks, were randomly selected from different stockyards in each farm and nasally swabbed for MRSA colonization. Fifty-one pigs from farm B were swabbed in 2016 (group 1) and 103 pigs were swabbed in 2018 (52 pigs from farm B [group 2] and 51 pigs from farm C [group 3])—Fig 1. Unfortunately, we could not obtain samples from farm C in 2016. MRSA isolates obtained from group 1 were isolated and characterized in our previous study [2].

Ethics statement

The protocol was approved by the Research Board of Escola Superior de Saúde da Cruz Vermelha Portuguesa.

Sampling and MRSA identification

Samples were taken by swabbing a single nasal cavity of each animal with a sterile cotton swab, which was stored in Stuart transport medium. After overnight enrichment growth at 37°C in Mueller-Hinton broth (Becton, Dickinson & Co, New Jersey, USA), the overnight samples were inoculated on Tryptic Soy Agar (TSA) (Becton, Dickinson & Co, New Jersey, USA) and Chromagar MRSA (ChromAgar, Paris, France). MRSA was confirmed by PCR amplification of the *spa* gene for species identification, and the detection of the *mecA* gene [8, 9].

Molecular typing

The isolates were characterized by a combination of three typing methods. *Spa* typing was performed as previously described [9] and *spa* types were assigned through the Ridom web server

(<http://spaserver.ridom.de>). Multilocus sequence typing (MLST) was performed as previously described [10] and the allelic profiles and sequence types (ST) were defined using the MLST online database (<https://pubmlst.org/saureus/>). Staphylococcal cassette chromosome *mec* (SCC*mec*) was characterized by multiplex PCR [11].

Susceptibility testing

Antimicrobial susceptibility testing was performed by disk diffusion, according to the European Committee on Antimicrobial Susceptibility Testing (<http://www.eucast.org>), for the following antibiotics: ceftiofur (FOX), ciprofloxacin (CIP), chloramphenicol (CHL), clindamycin (CLIN), erythromycin (ERY), fusidic acid (FUS), gentamicin (GEN), linezolid (LZD), mupirocin (MUP), penicillin (PEN), quinupristin-dalfopristin (QD), rifampin (RIF), tetracycline (TET), and trimethoprim-sulfamethoxazole (SXT). Vancomycin (VAN) resistance was tested by E-test.

Whole genome sequencing

Whole genomic DNA of six MRSA isolates belonging to the two major *spa* types, t011 (n = 3) and t108 (n = 3), was extracted with the Sigma-Aldrich GenElute™ Bacterial Genomic DNA Kit. Genomic libraries were assessed using the NexteraXT library preparation kit (Illumina Inc., San Diego, CA) and sequencing was performed using the Illumina MiniSeq system with 150-bp paired-end reads and a coverage of 50X. Generated FastQ data were compiled and analyzed using the CLC genomic workbench 7.5.1 (CLC bio, Aarhus, Denmark). Reads were de novo assembled with automatic bubble and word size and contigs were generated using the mapping mode “map reads back to contigs” with a minimum contig length of 800 nucleotides. Antimicrobial resistance genes were identified using the ResFinder database [12] and the Comprehensive Antibiotic Resistance Database (CARD) platform [13]. Since all pigs received zinc oxide in their feed regimen, the presence of the *czrC* gene encoding resistance to zinc and cadmium [14] was evaluated by a BLAST (Basic Local Alignment Search Tool) analysis over the GenBank database.

Detection of resistance genes by PCR

Resistance genes detected by WGS on representative isolates, namely genes *fexA*, *dfrG*, *aac(6′)-Ie-aph(2′′)-Ia*, *ermA*, *ermC*, and *ermT* have been additionally confirmed by PCR [15–17].

Statistical analysis

Chi-squared or Fisher’s exact test were used to determine the differences between the MRSA prevalence and resistance to the different antibiotics among isolates recovered from the three groups of pigs with different antibiotic feed regimens. *P*-values <0.01 were considered statistically significant.

Results

MRSA prevalence

Overall, 96% (99 out of 103) of the piglets swabbed in 2018 (farm B n = 48/52 and farm C n = 51/51) were nasally colonized with MRSA. This rate was similar to the one found in 2016 (99%) [2]. No differences in MRSA prevalence were observed regarding the three groups of pigs under different antibiotic feed regimens (*p*<0.01).

Molecular characterization of MRSA

All isolates ($n = 157$) belonged to ST398, independently of the farm, sampling period, and antibiotic administration in the feed regimen. Most of the isolates harbored SCC*mec* type V ($n = 146/157$; 97%), while 3% were SCC*mec* non-typeable by the multiplex strategy (amplification of *mecA* was obtained only). Three *spa* types were detected within the whole collection (2016 and 2018), namely t011 ($n = 108$; 70%), t108 ($n = 45$; 29%), and a novel type t18272 ($n = 4$; 3%)—Table 1. The two major types, t011 and t108, were present in the four groups of isolates, while t18272 was exclusively found in group 2. Interestingly, t011 was the predominant clone in all groups with the exception of group 1 in which 62% of the isolates corresponded to t108.

Antimicrobial susceptibility

None of the isolates showed decreased susceptibility to fusidic acid, linezolid, rifampicin, mupirocin, and vancomycin, while all isolates were resistant to cefoxitin and tetracycline, and 99% were resistant to clindamycin and QD.

Analysis of the antimicrobial susceptibility profiles of the MRSA isolates collected in the three groups, showed striking differences (Table 2). Overall, in group 1 (receiving colistin and amoxicillin in the feed regimen in 2016), ciprofloxacin was the single antibiotic to which more than 50% of the isolates were susceptible, while in group 2 (receiving amoxicillin only) the large majority of the isolates was susceptible to gentamicin and chloramphenicol, and in group 3 (pigs not receiving antibiotics) the majority of the isolates remained susceptible to four antibiotics (erythromycin, ciprofloxacin, gentamicin, and chloramphenicol).

Antimicrobial susceptibility by *spa* type

By comparing the antimicrobial susceptibility profiles of all isolates belonging to each of the two major *spa* types (Table 2), resistance to ciprofloxacin and SXT was significantly higher among t011 isolates compared to t108 isolates (58% vs 0%; $p < 0.01$ and 92% vs 24%; $p < 0.001$, respectively), while resistance to erythromycin, gentamicin and chloramphenicol was lower in the former group (56% vs 91%, $p < 0.001$; 20% vs 71%, $p < 0.01$; and 14% vs 82%, $p < 0.001$, respectively).

By comparing t011 isolates recovered from farm B pigs that received colistin and amoxicillin in 2016 (group 1) and isolates from the same farm, from pigs receiving only amoxicillin two years later (group 2), a difference in chloramphenicol susceptibility was demonstrated, with isolates from group 2 being susceptible while those of group 1 being resistant. Similarly, most of the t108 isolates from group 2 showed higher rates of susceptibility to gentamicin and chloramphenicol compared to those of group 1 (Table 2).

Table 1. Distribution of *spa* types of the 157 ST398-MRSA isolates from the three groups of pigs.

<i>spa</i> type	Group 1 [2] [Farm B; 2016; AMX+COL]*	Group 2 [Farm B; 2018; AMX]*	Group 3 [Farm C; 2018; none]*	Total
t011	22 (38%)	38 (79%)	48 (94%)	108 (70%)
t108	36 (62%)	6 (13%)	3 (6%)	45 (29%)
t18272		4 (8%)		4 (3%)
Total	58	48	51	157

* Antibiotics included in the feed regimen. AMX—Amoxicillin; COL—Colistin.

The prevalent *spa* type in each group is displayed in bold.

Percentages referred to the total number of isolates in each group.

<https://doi.org/10.1371/journal.pone.0225497.t001>

Table 2. Antibiotic resistance of the 157 ST398-MRSA isolates from the three groups of pigs.

	Total	Group 1 [2] [Farm B; 2016; AMX+COL] ^a	Group 2 [Farm B; 2018; AMX] ^a	Group 3 [Farm C; 2018; none] ^a
Total collection	157	58	48	51
FOX	157 (100%)	58 (100%)	48 (100%)	51 (100%)
ERY	106 (68%)	58 (100%)*	46 (96%)*	2 (4%)*
CLIN	155 (99%)	58 (100%)	48 (100%)	49 (96%)
CIP	67 (43%)	22 (38%)	42 (88%)*	3 (6%)*
TET	157 (100%)	58 (100%)	48 (100%)	51 (100%)
SXT	114 (73%)	31 (53%)*	44 (92%)*	39 (76%)
GEN	54 (34%)	29 (50%)	2 (4%)*	23 (45%)
QD	156 (99%)	58 (100%)	48 (100%)	50 (98%)
CHL	52 (33%)	51 (88%)*	1 (2%)*	0*
<i>spa</i> t011	108	22	38	48
FOX	108 (100%)	22 (100%)	38 (100%)	48 (100%)
ERY	60 (56%)	22 (100%)*	36 (95%)*	2 (4%)*
CLIN	106 (98%)	22 (100%)	38 (100%)	46 (96%)
CIP	63 (58%)	22 (100%)*	38 (100%)*	3 (6%)*
TET	108 (100%)	22 (100%)	38 (100%)	48 (100%)
SXT	99 (92%)	22 (100%)	38 (100%)	39 (81%)*
GEN	22 (20%)	0*	0*	22 (46%)
QD	107 (99%)	22 (100%)	38 (100%)	47 (98%)
CHL	15 (14%)	15 (68%)*	0*	0*
<i>spa</i> t108	45	36	6	3
FOX	45 (100%)	36 (100%)	6 (100%)	3 (100%)
ERY	41 (91%)	36 (100%)*	5 (83%)	0*
CLIN	45 (100%)	36 (100%)	6 (100%)	3 (100%)
CIP	0 (0%)	0*	0*	0*
TET	45 (100%)	36 (100%)	6 (100%)	3 (100%)
SXT	11 (24%)	9 (25%)	2 (33%)	0*
GEN	32 (71%)	29 (81%)*	2 (33%)	1 (33%)
QD	45 (100%)	36 (100%)*	6 (100%)	3 (100%)
CHL	37 (82%)	36 (100%)*	1 (17%)*	0*

FOX–Cefoxitin; ERY–Erythromycin; CLIN–Clindamycin; CIP–Ciprofloxacin; TET–Tetracycline; SXT–Trimethoprim-sulfamethoxazole; GEN–Gentamicin; QD–Quinupristin-dalfopristin; CHL–Chloramphenicol.

^aAntibiotics included in the feed regimen. AMX–Amoxicillin; COL–Colistin.

*Significant difference (p<0.01).

Numbers in bold indicate that ≥50% of the isolates are resistant to the antibiotic.

<https://doi.org/10.1371/journal.pone.0225497.t002>

Likewise, the large majority of the isolates recovered from pigs that did not receive any antibiotic in the feed regimen (group 3), for both *spa* types, remained susceptible to erythromycin, ciprofloxacin, gentamicin, and chloramphenicol, and also to SXT for *spa* t108 (Table 2).

Whole genome sequencing

To gain insights into the antimicrobial resistance genotypes that may explain the different phenotypes observed among the different groups of isolates, WGS was performed for six isolates representatives of the three groups (Fig 2).

Sequence analysis followed by PCR showed that, independently of the *spa* type, resistance to chloramphenicol was due to the presence of the phenicol exporter encoding gene *fexA*,

Also, all isolates harbored the *czrC* gene, which is frequently localized together with *mecA* on SCC*mec* elements from LA-MRSA, in particular SCC*mec* type V [24]. The fact that zinc oxide was given as a feed supplement constituted a selective pressure for acquisition of β -lactam resistance, and therefore selection of MRSA, despite the lack of β -lactam selective pressure. Future studies in farms that will ban the use of zinc oxide in the feed regimen will be of interest to confirm this phenomenon. Moreover, it has been shown that the presence of both *tet(K)* and *tet(M)* confers a fitness advantage to LA-MRSA CC398, which associated to *czrC* might drive the expansion of this clone [25].

In our collection, MRSA isolates recovered from pigs receiving no antibiotic in the feed regimen showed susceptibility to a higher number of antimicrobial agents compared to isolates from pigs receiving colistin and/or amoxicillin, suggesting that a lower antimicrobial exposure correlates with a lower rate of antibiotic resistance among MRSA colonizing the anterior nares of healthy pigs. This is in agreement with previous studies that showed that any form of antimicrobial exposure in swine, including different modes of administration, actually increases the prevalence of antibiotic-resistant bacteria in their gut [5, 6]. Of note, many of the antibiotic resistance genes found among isolates recovered from pigs receiving antibiotics in the feed regimen and absent in isolates recovered from pigs not receiving antibiotics are plasmid-encoded, namely *ermT*, *tetL*, *fexA*, *spd*, *apmA*, and *dfrG* [15, 26]. Therefore, the absence of antibiotic selective pressure might have driven the loss of these genetic elements over time, that in most cases are of small size (<15 kb). However, given that in some cases the observed differences in antibiotic susceptibility are outside of the classes of drugs no longer administered, these changes might be potentially just happenstance.

Another important finding from this surveillance study was the detection of a high proportion of MRSA isolates presenting erythromycin susceptibility and clindamycin resistance (25%), mostly in isolates recovered from pigs receiving no antibiotic in the feed regimen (90%). Previous studies have found this uncommon phenotype among MRSA-ST398 swine isolates associated with the *lnu(A)* or *lnu(B)* genes and this phenotype seems to be related to *S. aureus* animal-associated clonal lineages [27, 28]. Our isolates did not carry any of the *lnu* genes but instead harbored the *vga_{ALC}* gene that encodes resistance to lincosamides. This variant of the *vgaA* gene has substrate specificity towards lincosamides and has been previously found in clinical isolates of *Staphylococcus haemolyticus* resistant to lincomycin/clindamycin but susceptible to erythromycin, for which no relevant lincosamide resistance gene was found [19]. The *vga_{ALC}* gene has been previously reported in two *S. aureus* human clinical isolates responsible for skin and soft tissue infection that showed the same erythromycin susceptibility and clindamycin resistance phenotype [29]. Moreover, *vga_{ALC}* was detected in the genome of a single swine LA-MRSA ST5 isolate recovered in the United States [30, 31], but to our knowledge, this is the first identification of the *vga_{ALC}* gene in the widely widespread LA-MRSA lineage ST398.

In summary, MRSA currently colonizing the nares of healthy pigs in Portugal belong to ST398-V, mainly associated with *spa* types t011 and t108. A considerable proportion of MRSA-ST398-t011 isolates presented the unusual phenotype macrolide-susceptibility/lincosamide-resistance associated to the presence of the *vga_{ALC}* gene. Pigs receiving less antibiotics as feed additives but still receiving zinc oxide maintained high MRSA nasal carriage rates, which was likely related to this heavy metal selective pressure. However, those MRSA isolates colonizing the pigs were resistant to less classes of antibiotics.

Acknowledgments

We thank Nicolas Kieffer and José Manuel Ortiz-de-la-Rosa for technical assistance.

Author Contributions

Conceptualization: Marta Aires-de-Sousa.

Funding acquisition: Marta Aires-de-Sousa.

Investigation: Elizeth Lopes, Teresa Conceição.

Supervision: Teresa Conceição, Laurent Poirel, Hermínia de Lencastre, Marta Aires-de-Sousa.

Writing – original draft: Marta Aires-de-Sousa.

Writing – review & editing: Laurent Poirel, Hermínia de Lencastre.

References

1. Aires de Sousa M. Methicillin-resistant *Staphylococcus aureus* among animals: current overview. *Clin Microbiol Infect*. 2016; 23(6):373–80. <https://doi.org/10.1016/j.cmi.2016.11.002> PMID: 27851997
2. Conceição T, de Lencastre H, Aires de Sousa M. Frequent isolation of methicillin resistant *Staphylococcus aureus* (MRSA) ST398 among healthy pigs in Portugal. *PLoS One*. 2017; 12(4):e0175340. <https://doi.org/10.1371/journal.pone.0175340> PMID: 28399155
3. Direção Geral de Alimentação e Veterinária Ministério da Agricultura e do Mar. Plano de ação nacional para a redução do uso de antibióticos nos animais. 2014.
4. Blake DP, Humphry RW, Scott KP, Hillman K, Fenlon DR, Low JC. Influence of tetracycline exposure on tetracycline resistance and the carriage of tetracycline resistance genes within commensal *Escherichia coli* populations. *Journal of applied microbiology*. 2003; 94(6):1087–97. Epub 2003/05/20. <https://doi.org/10.1046/j.1365-2672.2003.01937.x> PMID: 12752819.
5. Gellin G, Langlois BE, Dawson KA, Aaron DK. Antibiotic resistance of gram-negative enteric bacteria from pigs in three herds with different histories of antibiotic exposure. *Appl Environ Microbiol*. 1989; 55(9):2287–92. Epub 1989/09/01. PMID: 2802608; PubMed Central PMCID: PMC203070.
6. Bibbal D, Dupouy V, Ferre JP, Toutain PL, Fayet O, Prere MF, et al. Impact of three ampicillin dosage regimens on selection of ampicillin resistance in *Enterobacteriaceae* and excretion of *bla*TEM genes in swine feces. *Appl Environ Microbiol*. 2007; 73(15):4785–90. Epub 2007/06/15. <https://doi.org/10.1128/AEM.00252-07> PMID: 17557857; PubMed Central PMCID: PMC1951005.
7. Langlois BE, Cromwell GL, Stahly TS, Dawson KA, Hays VW. Antibiotic resistance of fecal coliforms after long-term withdrawal of therapeutic and subtherapeutic antibiotic use in a swine herd. *Appl Environ Microbiol*. 1983; 46:1433–4. PMID: 6660878
8. Okuma K, Iwakawa K, Turnidge JD, Grubb WB, Bell JM, O'Brien FG, et al. Dissemination of new methicillin-resistant *Staphylococcus aureus* clones in the community. *J Clin Microbiol*. 2002; 40(11):4289–94. <https://doi.org/10.1128/JCM.40.11.4289-4294.2002> PMID: 12409412
9. Aires-de-Sousa M, Boye K, de Lencastre H, Deplano A, Enright MC, Etienne J, et al. High interlaboratory reproducibility of DNA sequence-based typing of bacteria in a multicenter study. *J Clin Microbiol*. 2006; 44(2):619–21. <https://doi.org/10.1128/JCM.44.2.619-621.2006> PMID: 16455927.
10. Enright MC, Day NP, Davies CE, Peacock SJ, Spratt BG. Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. *J Clin Microbiol*. 2000; 38(3):1008–15. PMID: 10698988
11. Milheiro C, Oliveira DC, de Lencastre H. Update to the multiplex PCR strategy for assignment of *mec* element types in *Staphylococcus aureus*. *Antimicrob Agents Chemother*. 2007; 51(9):3374–7. <https://doi.org/10.1128/AAC.00275-07> PMID: 17576837.
12. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, et al. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother*. 2012; 67(11):2640–4. Epub 2012/07/12. <https://doi.org/10.1093/jac/dks261> PMID: 22782487; PubMed Central PMCID: PMC3468078.
13. Jia B, Raphenya AR, Alcock B, Waglechner N, Guo P, Tsang KK, et al. CARD 2017: expansion and model-centric curation of the comprehensive antibiotic resistance database. *Nucleic Acids Res*. 2017; 45(D1):D566–D73. Epub 2016/10/30. <https://doi.org/10.1093/nar/gkw1004> PMID: 27789705; PubMed Central PMCID: PMC5210516.
14. Cavaco LM, Hasman H, Stegger M, Andersen PS, Skov R, Fluit AC, et al. Cloning and occurrence of *czrC*, a gene conferring cadmium and zinc resistance in methicillin-resistant *Staphylococcus aureus* CC398 isolates. *Antimicrob Agents Chemother*. 2010; 54(9):3605–8. Epub 2010/06/30. <https://doi.org/10.1128/AAC.00058-10> PMID: 20585119; PubMed Central PMCID: PMC2934997.

15. Kehrenberg C, Schwarz S. Distribution of florfenicol resistance genes *fxaA* and *cfr* among chloramphenicol-resistant *Staphylococcus* isolates. *Antimicrob Agents Chemother*. 2006; 50(4):1156–63. Epub 2006/03/30. <https://doi.org/10.1128/AAC.50.4.1156-1163.2006> PMID: 16569824; PubMed Central PMCID: PMC1426988.
16. Argudin MA, Tenhagen BA, Fetsch A, Sachsenroder J, Kasbohrer A, Schroeter A, et al. Virulence and resistance determinants of German *Staphylococcus aureus* ST398 isolates from nonhuman sources. *Appl Environ Microbiol*. 2011; 77(9):3052–60. Epub 2011/03/08. <https://doi.org/10.1128/AEM.02260-10> PMID: 21378035; PubMed Central PMCID: PMC3126402.
17. Fessler A, Scott C, Kadlec K, Ehrlich R, Monecke S, Schwarz S. Characterization of methicillin-resistant *Staphylococcus aureus* ST398 from cases of bovine mastitis. *J Antimicrob Chemother*. 2010; 65(4):619–25. Epub 2010/02/19. <https://doi.org/10.1093/jac/dkq021> PMID: 20164198.
18. Trong HN, Prunier AL, Leclercq R. Hypermutable and fluoroquinolone-resistant clinical isolates of *Staphylococcus aureus*. *Antimicrob Agents Chemother*. 2005; 49(5):2098–101. Epub 2005/04/28. <https://doi.org/10.1128/AAC.49.5.2098-2101.2005> PMID: 15855537; PubMed Central PMCID: PMC1087674.
19. Novotna G, Janata J. A new evolutionary variant of the streptogramin A resistance protein, Vga(A)LC, from *Staphylococcus haemolyticus* with shifted substrate specificity towards lincosamides. *Antimicrob Agents Chemother*. 2006; 50(12):4070–6. Epub 2006/10/04. <https://doi.org/10.1128/AAC.00799-06> PMID: 17015629; PubMed Central PMCID: PMC1693986.
20. Tang Y, Larsen J, Kjeldgaard J, Andersen PS, Skov R, Ingmer H. Methicillin-resistant and -susceptible *Staphylococcus aureus* from retail meat in Denmark. *Int J Food Microbiol*. 2017; 249:72–6. Epub 2017/03/23. <https://doi.org/10.1016/j.ijfoodmicro.2017.03.001> PMID: 28324679.
21. van Alen S, Ballhausen B, Peters G, Friedrich AW, Mellmann A, Köck R, et al. In the centre of an epidemic: Fifteen years of LA-MRSA CC398 at the University Hospital Münster. *Vet Microbiol*. 2016; 200:19–24. <https://doi.org/10.1016/j.vetmic.2016.01.021> PMID: 26878970
22. Wulf MW, Sorum M, van Nes A, Skov R, Melchers WJ, Klaassen CH, et al. Prevalence of methicillin-resistant *Staphylococcus aureus* among veterinarians: an international study. *Clin Microbiol Infect*. 2008; 14(1):29–34. Epub 2007/11/08. <https://doi.org/10.1111/j.1469-0691.2007.01873.x> PMID: 17986212.
23. Agency EM. Sales of veterinary antimicrobial agents in 30 European countries in 2016—Trends from 2010 to 2016 Eighth ESVAC report. 2018.
24. Monecke S, Slickers P, Gawlik D, Muller E, Reissig A, Ruppelt-Lorz A, et al. Variability of SCCmec elements in livestock-associated CC398 MRSA. *Vet Microbiol*. 2018; 217:36–46. Epub 2018/04/05. <https://doi.org/10.1016/j.vetmic.2018.02.024> PMID: 29615254.
25. Larsen J, Clasen J, Hansen JE, Paulander W, Petersen A, Larsen AR, et al. Copresence of *tet(K)* and *tet(M)* in livestock-associated methicillin-resistant *Staphylococcus aureus* clonal complex 398 is associated with increased fitness during exposure to sublethal concentrations of tetracycline. *Antimicrob Agents Chemother*. 2016; 60(7):4401–3. Epub 2016/05/11. <https://doi.org/10.1128/AAC.00426-16> PMID: 27161637; PubMed Central PMCID: PMC4914685.
26. Fessler A, Kadlec K, Wang Y, Zhang WJ, Wu C, Shen J, et al. Small antimicrobial resistance plasmids in livestock-associated methicillin-resistant *Staphylococcus aureus* CC398. *Frontiers in microbiology*. 2018; 9:2063. Epub 2018/10/05. <https://doi.org/10.3389/fmicb.2018.02063> PMID: 30283407; PubMed Central PMCID: PMC6157413.
27. Lozano C, Aspiroz C, Ara M, Gomez-Sanz E, Zarazaga M, Torres C. Methicillin-resistant *Staphylococcus aureus* (MRSA) ST398 in a farmer with skin lesions and in pigs of his farm: clonal relationship and detection of *Inu(A)* gene. *Clin Microbiol Infect*. 2011; 17(6):923–7. Epub 2011/06/21. <https://doi.org/10.1111/j.1469-0691.2010.03437.x> PMID: 21682806.
28. Lozano C, Aspiroz C, Saenz Y, Ruiz-Garcia M, Royo-Garcia G, Gomez-Sanz E, et al. Genetic environment and location of the *Inu(A)* and *Inu(B)* genes in methicillin-resistant *Staphylococcus aureus* and other staphylococci of animal and human origin. *J Antimicrob Chemother*. 2012; 67(12):2804–8. Epub 2012/08/18. <https://doi.org/10.1093/jac/dks320> PMID: 22899804.
29. Qin X, Poon B, Kwong J, Niles D, Schmidt BZ, Rajagopal L, et al. Two paediatric cases of skin and soft-tissue infections due to clindamycin-resistant *Staphylococcus aureus* carrying a plasmid-encoded vga (A) allelic variant for a putative efflux pump. *Int J Antimicrob Agents*. 2011; 38(1):81–3. Epub 2011/05/10. <https://doi.org/10.1016/j.ijantimicag.2011.03.007> PMID: 21549571.
30. Otarigho B, Falade MO. Analysis of antibiotics resistant genes in different strains of *Staphylococcus aureus*. *Bioinformatics*. 2018; 14(3):113–22. Epub 2018/05/23. <https://doi.org/10.6026/97320630014113> PMID: 29785070; PubMed Central PMCID: PMC5953858.
31. Hau SJ, Bayles DO, Alt DP, Frana TS, Nicholson TL. Complete genome sequence of a livestock-associated methicillin-resistant *Staphylococcus aureus* Sequence Type 5 Isolate from the United States. *Genome Announc*. 2017; 5(32). Epub 2017/08/12. <https://doi.org/10.1128/genomeA.00791-17> PMID: 28798188; PubMed Central PMCID: PMC5552997.