

**Human basophils may not undergo modulation by
DC-SIGN and mannose receptor–targeting
immunotherapies due to absence of receptors**

Mrinmoy Das, Caroline Galeotti, Emmanuel Stephen-Victor, Anupama
Karnam, Srini Kaveri, Jagadeesh Bayry

► **To cite this version:**

Mrinmoy Das, Caroline Galeotti, Emmanuel Stephen-Victor, Anupama Karnam, Srini Kaveri, et al.. Human basophils may not undergo modulation by DC-SIGN and mannose receptor–targeting immunotherapies due to absence of receptors. *Journal of Allergy and Clinical Immunology*, Elsevier, 2017, 139 (4), pp.1403-1404.e1. 10.1016/j.jaci.2016.09.062 . inserm-02455463

HAL Id: inserm-02455463

<https://www.hal.inserm.fr/inserm-02455463>

Submitted on 26 Jan 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Human basophils may not undergo modulation by DC-SIGN and mannose receptor–targeting immunotherapies due to absence of receptors



To the Editor:

Sirvent et al¹ recently showed that novel vaccines targeting dendritic cells (DCs) by coupling glutaraldehyde-polymerized grass pollen allergoids to nonoxidized mannan enhance allergen uptake and induce functional regulatory T cells through programmed death ligand 1. Mechanistically, they found that nonoxidized mannan-coupled glutaraldehyde-polymerized grass pollen allergoids are captured and internalized by 2 lectin receptors on DCs: mannose receptor (CD206) and DC-specific intercellular adhesion molecule-3-grabbing nonintegrin (DC-SIGN or CD209). These data thus indicated that DCs could be targeted by C-type lectin receptors for efficient allergen immunotherapy.

Basophils are one of the key players of allergic responses. They mediate allergic inflammation by secretion of T_H2-polarizing cytokines IL-4 and IL-13 and by the release of effector molecules such as histamines and leukotrienes upon FcεRI signaling by IgE-allergen complexes.² Basophils receive activation signals not only via allergen-IgE complexes³ but also via Toll-like receptors⁴ and possibly C-type lectin receptors. In fact, basophils express several lectin receptors such as C-type lectin domain family 12 member A and DC immunoreceptor.^{5,6} Thus, it is likely that in addition to DCs, nonoxidized mannan-coupled allergoids might modulate basophil functions to exert immunotherapeutic benefits. Therefore, we analyzed the expression of mannose receptor and DC-SIGN on steady-state circulating human basophils and on stimulated basophils.

We analyzed basophils in whole blood of healthy donors without their purification to avoid any loss of cells and consequently misinterpretation of data (see this article's Online Repository at www.jacionline.org). Furthermore, erythrocyte-lysed whole blood cells were stimulated with IL-3 (100 ng/10⁶ cells) for 24 hours. IL-3-stimulated basophils were also stimulated for degranulation with anti-IgE antibodies (100 ng/10⁶ cells) for 30 minutes. As controls for the expression of DC-SIGN and CD206, we used CD14⁺ PBMCs (negative control), and rhIL-4 (500 IU/10⁶ cells) and rhGM-CSF (1000 IU/10⁶ cells)-differentiated monocyte-derived DCs (positive control).⁷

Circulating basophils were identified as positive for FcεRI and CD123 and negative for BDCA-4. We found that human basophils at steady state are negative for DC-SIGN and CD206 (Fig 1, A and B). Because basophils display enhanced expression of various receptors on receiving activation stimuli, we explored whether they express these lectin receptors on activation. However, irrespective of stimulation (IL-3 or degranulation stimuli), basophils remained negative for DC-SIGN and CD206 (Fig 1, B and C). Absence of DC-SIGN was also confirmed on isolated basophils. Furthermore, the absence of DC-SIGN and CD206 on basophils in our report is not due to nonreactivity of antibodies used in flow cytometry

because monocyte-derived DCs, used as positive control, uniformly expressed CD206 and DC-SIGN (Fig 1, A and B). As expected, CD14⁺ circulating monocytes, used as negative control, did not stain for both the markers, thus confirming lack of nonspecific binding of antibodies (Fig 1, A and B).

Our results thus indicate that human basophils lack DC-SIGN and mannose receptors and hence unlike DCs, they may not directly respond and be modulated by DC-SIGN- and mannose receptor-binding nonoxidized mannan-coupled allergoids. In addition, our data suggest that basophils do not get activated by DC-SIGN- and mannose receptor-binding allergens unless they are IgE-bound. Thus, the expression pattern of DC-SIGN and mannose receptor among innate cells diversifies allergic as well as tolerogenic responses.

Mrimoy Das, MSc^{a,b,c,*}
 Caroline Galeotti, MD^{a,b,c,d,*}
 Emmanuel Stephen-Victor, MSc^{a,b,c,*}
 Anupama Karnam, MSc, PhD^{a,b,c}
 Sridi V. Kaveri, DVM, PhD^{a,b,c,e}
 Jagadeesh Bayry, DVM, PhD^{a,b,c,e}

From ^aInstitut National de la Santé et de la Recherche Médicale Unité 1138, ^bSorbonne Universités, UPMC Univ Paris 06, UMR S 1138, ^cCentre de Recherche des Cordeliers, Equipe-Immunopathologie et immuno-intervention thérapeutique, ^dthe Department of Pediatric Rheumatology, National Referral Centre of Auto-inflammatory Diseases, CHU de Bicêtre, le Kremlin Bicêtre and ^eUniversité Paris Descartes, Sorbonne Paris Cité, UMR S 1138, Paris, France. E-mail: jagadeesh.bayry@crc.jussieu.fr.

*These authors contributed equally to this work.

This study was supported by Institut National de la Santé et de la Recherche Médicale (INSERM), Université Pierre et Marie Curie, Université Paris Descartes, and CSL Behring, Switzerland. C.G. is a recipient of fellowship from La Fondation pour la Recherche Médicale (FDM20150633674), France, and E.S.-V. and A.K. are recipients of fellowships from the Indo-French Center for Promotion of Advanced Research.

Disclosure of potential conflict of interest: C. Galleotti has received a PhD fellowship from La Fondation pour la Recherche Médicale. E. Stephen-Victor and A. Karnam have received PhD fellowships from the Indo-French Center for Promotion of Advanced Research. J. Bayry has received a grant from CSL Behring. The rest of the authors declare that they have no relevant conflicts of interest.

REFERENCES

1. Sirvent S, Soria I, Cirauqui C, Cases B, Manzano AI, Diez-Rivero CM, et al. Novel vaccines targeting dendritic cells by coupling allergoids to nonoxidized mannan enhance allergen uptake and induce functional regulatory T cells through programmed death ligand 1. *J Allergy Clin Immunol* 2016;138:558-67.e11.
2. Voehringer D. Protective and pathological roles of mast cells and basophils. *Nat Rev Immunol* 2013;13:362-75.
3. Oettgen HC. Fifty years later: emerging functions of IgE antibodies in host defense, immune regulation, and allergic diseases. *J Allergy Clin Immunol* 2016;137:1631-45.
4. Suurmond J, Stoop JN, Rivellese F, Bakker AM, Huizinga TW, Toes RE. Activation of human basophils by combined Toll-like receptor- and FcεRI-triggering can promote Th2 skewing of naive T helper cells. *Eur J Immunol* 2014;44:386-96.
5. Bates EE, Fournier N, Garcia E, Valladeau J, Durand I, Pin JJ, et al. APCs express DCIR, a novel C-type lectin surface receptor containing an immunoreceptor tyrosine-based inhibitory motif. *J Immunol* 1999;163:1973-83.
6. Divekar A, Conklin J, Yang X, Ransom J. Role of C-type lectin receptor CLEC12A in human basophil activation (48.6). *J Immunol* 2012;188:48.6.
7. Maddur MS, Sharma M, Hegde P, Stephen-Victor E, Pulendran B, Kaveri SV, et al. Human B cells induce dendritic cell maturation and favour Th2 polarization by inducing OX-40 ligand. *Nat Commun* 2014;5:4092.

Available online January 11, 2017.
<http://dx.doi.org/10.1016/j.jaci.2016.09.062>




FIG 1. Human basophils are deficient for DC-SIGN and CD206. **A-C,** Flow cytometric analysis of DC-SIGN and CD206 on steady-state basophils, stimulated basophils (IL-3 or IL-3 and anti-IgE), and monocytes and monocyte-derived DCs. Representative dot-plots and percentage of cells (mean ± SEM, n = 3) positive for DC-SIGN and CD206. ****P* < .001.

METHODS

Cells and stimulation

Buffy bags of healthy donors were obtained from Centre Necker-Cabanel (EFS, Paris, France), and INSERM-EFS ethical committee approval (N°15/EFS/012) for the use of such material was obtained. Experiments were performed in accordance with the approved guidelines of INSERM.

Red blood cells were lysed using Ammonium-Chloride-Potassium Lysing Buffer (Lonza, Levallois, France). Briefly, blood was spun down and resuspended and incubated in Ammonium-Chloride-Potassium lysing buffer for 30 to 60 seconds. Cells were washed with medium and resuspended in serum-free X-VIVO medium. Cells were stimulated with IL-3 (100 ng/million cells; ImmunoTools, Friesoythe, Germany) for 24 hours. In addition, cells were cultured with IL-3 (100 ng/million cells) for up to 24 hours and during the last 30 minutes, cells were treated with anti-IgE antibodies (100 ng/million cells; Sigma-Aldrich, St Quentin Fallavier, France). Phenotype of basophils was analyzed in steady state and stimulated conditions by flow cytometry (LSR II, BD Biosciences, Le Pont De Claix, France) and the data were analyzed using FACSDiva software (BD Biosciences).

PBMCs were obtained from buffy bags of healthy donors by Ficoll density gradient centrifugation. Monocytes were isolated from PBMCs by using CD14 microbeads (Miltenyi Biotec, Paris, France) and were cultured for 5 days in rhIL-4 (500 IU/10⁶ cells) and rhGM-CSF (1000 IU/10⁶ cells) (both from Miltenyi Biotec) to obtain monocyte-derived DCs.

Antibodies for flow cytometry

The following antibodies were used for flow cytometry. FcεRIα-BV510 (Clone AER37 [CRA-1]) was from BioLegend (London, United Kingdom) and BDCA-4 (CD304)-APC (Clone AD5-17F6) was obtained from Miltenyi Biotec. CD123-BV421 (Clone 9F5), CD209-FITC (Clone DCN46), CD206-PE (Clone 19.2), HLA-DR-APC or PE (G46-6, BD Biosciences), and CD14-APC (Clone M5E2) antibodies were from BD Biosciences.

Statistical analysis

Levels of significance for comparison between samples were determined by 1-way ANOVA (repeated measures with Tukey's multiple comparison test). A *P* value of less than .05 was considered significant. Statistical analysis was performed by using Prism 5 GraphPad Software (La Jolla, Calif). Data are presented as mean ± SEM.