



HAL
open science

Interleukin-1 Family Cytokines: Keystones in Liver Inflammatory Diseases

Louise Barbier, Maroua Ferhat, Ephrem Salamé, Aurélie Robin, André Herbelin, Jean-Marc Gombert, Christine Silvain, Alice Barbarin

► **To cite this version:**

Louise Barbier, Maroua Ferhat, Ephrem Salamé, Aurélie Robin, André Herbelin, et al.. Interleukin-1 Family Cytokines: Keystones in Liver Inflammatory Diseases. *Frontiers in Immunology*, 2019, 10, pp.2014. 10.3389/fimmu.2019.02014 . inserm-02509347

HAL Id: inserm-02509347

<https://inserm.hal.science/inserm-02509347>

Submitted on 16 Mar 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Interleukin-1 Family Cytokines: Keystones in Liver Inflammatory Diseases

Louise Barbier^{1,2}, Maroua Ferhat³, Ephrem Salamé^{1,2}, Aurélie Robin⁴, André Herbelin^{3*}, Jean-Marc Gombert^{1,5}, Christine Silvain⁶ and Alice Barbarin¹

¹INSERM U1082, Poitiers, France, ²Department of Digestive Surgery and Liver Transplantation, Trousseau University Hospital, Tours University, Tours, France, ³INSERM U1082, Poitiers University, Poitiers, France, ⁴INSERM U1082, Poitiers University Hospital, Poitiers, France, ⁵Department of Immunology and Inflammation, Poitiers University Hospital, University of Poitiers, Poitiers, France, ⁶Department of Hepatology and Gastroenterology, Poitiers University Hospital, University of Poitiers, Poitiers, France

OPEN ACCESS

Edited by:

Elizabeth Brint,
University College Cork, Ireland

Reviewed by:

Aldo Tagliabue,
Institute for Genetic and Biomedical
Research (IRGB), Italy
Parameswaran Ramakrishnan,
Case Western Reserve University,
United States

*Correspondence:

André Herbelin
andre.herbelin@inserm.fr

Specialty section:

This article was submitted to
Cytokines and Soluble Mediators in
Immunity,
a section of the journal
Frontiers in Immunology

Received: 07 March 2019

Accepted: 08 August 2019

Published: 27 August 2019

Citation:

Barbier L, Ferhat M, Salamé E,
Robin A, Herbelin A, Gombert J-M,
Silvain C and Barbarin A (2019)
Interleukin-1 Family Cytokines:
Keystones in Liver Inflammatory
Diseases. *Front. Immunol.* 10:2014.
doi: 10.3389/fimmu.2019.02014

The pyrogenic property being the first activity described, members of the interleukin-1 superfamily (IL-1 α , IL-1 β , IL-18, and the newest members: IL-33, IL-36, IL-37, and IL-38) are now known to be involved in several inflammatory diseases such as obesity, atherosclerosis, cancer, viral and parasite infections, and auto-inflammatory syndromes as well as liver diseases. Inflammation processes are keystones of chronic liver diseases, of which the etiology may be viral or toxic, as in alcoholic or non-alcoholic liver diseases. Inflammation is also at stake in acute liver failure involving massive necrosis, and in ischemia-reperfusion injury in the setting of liver transplantation. The role of the IL-1 superfamily of cytokines and receptors in liver diseases can be either protective or pro-inflammatory, depending on timing and the environment. Our review provides an overview of current understanding of the IL-1 family members in liver inflammation, highlighting recent key investigations, and therapeutic perspectives. We have tried to apply the concept of trained immunity to liver diseases, based on the role of the members of the IL-1 superfamily, first of all IL-1 β but also IL-18 and IL-33, in modulating innate lymphoid immunity carried by natural killer cells, innate lymphoid cells or innate T- $\alpha\beta$ lymphocytes.

Keywords: interleukin-1 family cytokines, liver diseases, inflammation, innate immunity, trained immunity, invariant natural killer T-cells, natural killer cells, innate lymphoid cells

THE EXPANDING FAMILY OF INTERLEUKIN (IL)-1 CYTOKINES AND THEIR RECEPTORS

From “Endogenous Pyrogens” to the IL-1 Superfamily of Cytokines

The story of the IL-1 superfamily of cytokines began in the 1940s when Menkin (1), Beeson (2), and then Atkins (3) described a group of proteins, named “endogenous pyrogens,” responsible for fever, produced by leucocytes, and able to mediate multiple biological activities (4–6). By the late 1970s, the term “interleukin” was being used to describe the pleiotropic factors implicated in inflammatory immune response and the designation “IL-1” was then used to define the factors secreted by macrophages (4, 6). The discovery of IL-1, and its two forms IL-1 α and IL-1 β , is considered the birth of “cytokine biology,” and it led to the discovery of more members. This

is how, over the years, IL-18, IL-33, IL-36, IL-37, and IL-38, which share several functional and structural properties, have been added to and included in the IL-1 superfamily of cytokines.

IL-1 Sub-Families

IL-1 superfamily (7) is divided into three sub-families according to the length of the N-term pro-peptides: IL-1 sub-family (IL-1 α , IL-1 β and IL-33, IL-1Ra), IL-18 sub-family (IL-18 and IL-37), and IL-36 sub-family (IL-36 α , β , γ , and IL-38) (**Table 1**). Among these cytokines, in each sub-family one may distinguish “pro-inflammatory” members such as IL-1 α , IL-1 β , IL-33 or IL-18 from “anti-inflammatory” members such as IL-1Ra, IL-36Ra, IL-37, or IL-38 [**Table 1**, and for reviews: (7–9)].

All of their receptors are heteropolymers of which at least one sub-unit is a member of the family of IL-1 receptors (IL-1R), which are characterized by extracellular immunoglobulin-like domains and an intracellular Toll/Interleukin-1R (TIR) domain within their cytoplasmic tail. Upon cell stimulation, IL-1R sub-units are oligomerized through the TIR domains. Secondly, MyD88 binds to the TIR domain, triggering nuclear factor- κ B (NF- κ B) translocation to the nucleus and activation of mitogen-activated protein kinases (MAPK) such as p38 and JNK, thereby leading to pro-inflammatory responses.

The combination of receptor sub-units conditions the type of signal transduced. For example, while association of IL-1RAcP (chain common to IL-1 and IL-1R3 receptors in the new nomenclature; **Table 1**) with the IL-1R1 sub-unit transduces an activation signal following binding of IL-1 α or IL-1 β , association of IL-1RAcP with IL-1R2 constitutes an IL-1 β receptor that transduces an inhibitory signal.

IL-1RAcP is also a sub-unit of the IL-33 and IL-36 receptors, which associate IL-1RAcP with the IL-1RL1 chain (also called ST2: “suppression of tumorigenicity 2”, T1/ST2 or IL-1-R4) and with the IL-1Rrp chain (also called IL-36 R or IL-1R6), respectively.

IL-1R8 (SIGIRR for Single Ig IL-1 related receptor), another receptor of the IL-1 family, has no known ligands but transduces

an inhibitory signal and has been described as a checkpoint for terminal maturation and acquisition of the effector functions of NK cells (10). Mantovani et al. have suggested that IL-1R8 interferes with the TIR-domain oligomerization of the IL-1 family receptors engaged by a ligand agonist, thereby leading to blocked recruitment of the MyD88 adapter (9, 11). Lastly, IL-1R8 is a sub-unit of the IL-37 receptor. IL-37 can reduce production of the pro-inflammatory cytokines IL-1 α and IL-1 β , IL-1Ra, IL-8, IL-17, IL-23, tumor necrosis factor (TNF α), and interferon γ (IFN γ) through IL-18R α and IL-1R8 (SIGIRR) (12).

It is important to note that there exist soluble forms of the IL-1, sIL-1R1, and sIL-1R2 receptors that possess the inhibitory functions of IL-1. On the same token, IL-18BP (Binding Protein), which consists in a single IgG domain such as the transmembrane receptor IL-1R8, is a soluble factor interfering with the liaison of IL-18 and its receptor. IL-18BP possesses a particularly strong affinity with IL-18. Seric concentrations of IL-18BP in a healthy individual generally range from 2,000 to 4,000 pg/ml, which means that in steady state circulation, there is always a major excess of IL-18BP compared to IL-18. In a pathological situation, however, the anti-inflammatory effects of IL-18BP are liable to be attenuated due to the fact that IL-18BP likewise binds to IL-37. While blocking the anti-inflammatory functions of IL-37, this association is liable to render excessive the active free form of IL-18.

As regards the IL-33 receptor, there exists a soluble free form of the chain specific to the ST2 receptor, sST2. It is derived from alternative splicing of the ST2 mRNA (13). The biological relevance of the sST2 chain has remained elusive: Is it, as suggested in some studies, a decoy receptor enabling IL-33 neutralization, or is it simply the signature of the bringing into play of the IL-33/ST2 axis, with the seric sST2 concentrations being correlated with the severity of the pathology under consideration (13–16).

All in all, these different elements demonstrate the existence of cross-regulation between the different IL-1 family cytokines, of which each member possesses a counter-regulatory receptor or ligand.

IL-1 Superfamily Members Differentially Depend on Processing Mechanisms to Function

IL-1 superfamily members (cytokines and receptors) contribute to a wide spectrum of immunological and inflammatory responses. Most of their members lack a signal peptide and are not immediately secreted. They are found diffusely in the cytoplasm as precursors containing cleavage sites and the difference that exists between members in terms of expression and proteolytic processing impact their respective roles *in vivo*. Indeed, to be biologically active, full-length precursors of IL-1 β , IL-18, and IL-37 require intracellular processing, which depends on caspase-1, and the activation of the NOD-like receptor family, pyrin domain containing 3 (NLRP3)-inflammasome, required for conversion of the procaspase-1 into the active caspase-1. While pro-IL-36 and pro-IL-38 also need to be cleaved to

Abbreviations: AID, induced cytidine deaminase; ACLF, HBV acute-on-chronic liver failure; ALD, alcoholic liver disease; ALF, acute liver failure; Areg, Amphiregulin; BP, binding protein; CHB, chronic HBV; ConA, concanavalin A; DAMP, damage-associated molecular pattern; DT2, type 2 diabetes; EMT, epithelial-mesenchymal transition; ERK, extracellular signal-regulated kinases; HAV, hepatitis virus A; HBV, hepatitis virus B; HCV, hepatitis virus C; HCC, hepatocellular carcinoma; HFD, high fat diet; HIF, hypoxia-inducible factor; ICAM-1, intercellular adhesion molecule 1; I κ B, inhibitor of kappa B; IL, interleukin; IL-1Ra, IL-1 receptor antagonist; IL-1R, IL-1 receptor; IL-1RAcP, IL-1 receptor accessory protein; IFN γ , interferon γ ; ILC, innate lymphoid cell; iNKT, invariant Natural Killer T; IR, ischemia-reperfusion; IRS-1, insulin receptor substrate-1; KC, Kupffer cells; LSEC, liver sinusoidal endothelial cells; MAPK, mitogen-activated protein kinases; miR, microRNA; NAFLD, non-alcoholic fatty liver disease; NF- κ B, nuclear factor- κ B; NLRP3, NOD-like receptor family, pyrin domain containing 3; NLRP6, NOD-like receptor family, pyrin domain containing 6; NAPQI, N-acetyl-*p*-benzoquinone-imine; NK, natural killer; PAMP, pathogen-associated molecular pattern; PMN, polymorphonuclear neutrophils; PR3, proteinase 3; ROS, Reactive oxygen species; SIGIRR, single Ig IL-1 related receptor; ST2L, suppression of tumorigenicity 2 ligand; sST2, soluble ST2; TAM, Tumor-associated macrophages; TCR, T-cell receptor; Th, T helper; TLR, toll-like receptor; TLR/IL-1R, TIR domain TLR/IL-1R; TNE, tumor necrosis factor; Treg, T regulatory; TXNIP, thioredoxin-interacting protein; WT, wild-type.

TABLE 1 | IL-1 cytokine superfamily characteristics.

Cytokine name	Specific receptor chain name	Alternative receptor name	Co-receptor	Alternative co-receptor name	Function	Inhibitory ligand/Interference ligand
IL-1 α , IL-1 β	IL-1R1	CD121a	IL-1R3	IL-1RAcP	Pro-inflammatory	IL-1Ra, sIL-1R1
IL-1 β	IL-1R2	CD121b	IL-1R3	IL-1RAcP	Anti-inflammatory	sIL-1R2
IL-18	IL-1R5	IL-18R α , IL-2Rrp, CD218a	IL-1R7, and/or IL-R8 (inhibitory)	IL-18R β , IL-18RAcP, CD218b	Pro-inflammatory	IL-18BP
IL-33	IL-1R4	T1/ST2, ST2L, IL-1RL1, IL-33R	IL-1R3 and/or IL-R8 (inhibitory)	IL-1RAcP	Pro-inflammatory	sST2 (sIL1R4)
IL-36 α , IL-36 β , IL-36 γ	IL-1R6	IL-1Rrp2, IL-1RL2, IL-36R	IL-1R3	IL-1RAcP	Pro-inflammatory	IL-36Ra
IL-36Ra	IL-1R6	IL-1Rrp2, IL-1RL2, IL-36R	IL-1R3	IL-1RAcP	Anti-inflammatory	
IL-37	IL-1R5	IL-18R α , IL-2Rrp, CD218a	IL-1R8	TIR8, SIGIRR	Anti-inflammatory	
IL-38	IL-1R6	IL-1Rrp2, IL-1RL2, IL-36R	IL-1R9	IL-1RAPL, TIGIRR-2	Anti-inflammatory	
Not known	NA		IL-1R8	TIR8, SIGIRR	Anti-inflammatory	
Not known	NA		IL-1R9 and IL-1R10	IL-1RAPL, TIGIRR-2, TIGIRR, TIGIRR-1	Not known	

IL-1RAcP, Interleukin-1 Receptor Associated Protein; IL-1RAPL, Interleukin-1 Associated Protein Like 1/2; IL-1Ra, Interleukin-1 antagonist; IL-18BP, IL-18 Binding Protein; SIGIRR, Single Ig Interleukin-1 Related Receptor; TIGIRR, Three Ig Interleukin-1 Related Receptor; NA, not applicable.

generate a mature active form, the underlying mechanism does not depend on caspase-1 and remains unclear.

Remarkably, IL-1 α and IL-33 are considered dual-function cytokines, meaning that in addition to their function as a classical cytokine, full-length IL-1 α and full-length IL-33 (as part of damage-associated molecular pattern or DAMP) act as an “alarmin.” IL-1 α and IL-33 precursors are constitutively expressed in the nuclei of epithelial and endothelial cells from different organs such as kidneys, liver, and lung. Moreover, the nuclear location of IL-33 seems to correspond to a storage function, a supposition confirmed by observation showing that mice expressing IL-33 from which the nuclear anchoring area has been eliminated have a fatal systemic inflammatory response syndrome, which is due to the absence of cellular retention of IL-33 (17). As regards IL-1 α , a steady-state shuttle of cytokines between nucleus and cytoplasm has been described (18). In the event of a pro-apoptotic signal, IL-1 α leaves the cytoplasm, is closely bound with the chromatin complex, and has no pro-inflammatory biological activity. By contrast, in the event of a necrotic signal, IL-1 α leaves the nucleus and moves toward the cytoplasm, as the necrotic cell releases pro-inflammatory IL-1 α (18). To this day no similar phenomenon has been documented regarding IL-33.

Consequent to tissue damage, biologically active full-length IL-1 α and IL-33 are rapidly released by necrotic cells to alert the immune system to the danger, resulting in rapid production of pro-inflammatory mediators and infiltration of polymorphonuclear neutrophils (PMN) initially followed by monocytes/macrophages to the insult site.

With the exception of IL-1 α and IL-33, cytokine precursors of the IL-1 family generally require intracellular processing to

generate active forms. However, extracellular processing also occurs, involving different proteases. Protease driven from PMN, including elastase and cathepsin G, as well as chymase from mast cells and macrophages, proteinase 3 from PMN and macrophages, granzyme B from natural killer (NK) cells, and meprims from epithelial cells, are able to convert pro-IL-1 β , pro-IL-18, pro-IL-36 α , β , and γ into their corresponding fully active forms. Active pro-IL-33 can also be cleaved extracellularly by these proteases to generate a super active form (Figure 1).


Immune Cell Targets of IL-1 Superfamily Members

IL-1 Superfamily Members and Myelopoiesis Control

While IL-1 superfamily members have a role in the activation and mobilization of monocytes and in the recruitment of PMN, they also fulfill a critical function in the induction of emergency hematopoiesis leading to sustained myeloid M1 skewing. In this setting, IL-1 α and IL-1 β are both liable to be produced by the medullar environment. However, emergency hematopoiesis and its pro-inflammatory component are strongly associated with IL-1 β , which mainly acts systemically (see section Concept of Trained Immunity and Innate Immunological Memory: The Place of the IL-1 Cytokine Family: trained immunity), whereas IL-1 α in its role as cytokine/alarmin seems to exercise juxtacrine action (9, 19–21).

Cytokines of the IL-1 Superfamily and Control of T Helper 1 Effectors and NK Cells

IL-18 and IL-1 are implicated in the genesis of type 1 responses through the recruitment and differentiation of T helper (Th)1 lymphocytes and the activation of NK cells and of innate



lymphoid cells (ILC)1, particularly by inducing IFN γ production. In a mouse model, absence of IL-18 production or of its mature form leads to loss of anti-tumoral activity, loss that is partially objectified by the absence of the FasL-dependent cytotoxic function by hepatic NK cells, a circumstance favoring pulmonary and colorectal cancers (22).

The existence of antigen-specific and/or NK memory cells was recently described [for review (23)]. Remarkably, these are liver-resident NK cells serving as the support for memory responses in mice and, quite probably, in humans as well (23–25). That much said, in men as well as mice there exist several models suggesting that IL-18 is a factor implicated in the generation of antigen-specific and/or NK memory cells (26). However, study of mice genetically deficient in IL-18 or IL-18R α shows that IL-18 is not essential in this function (24).

IL-1R8 seems to be an antagonist of IL-18 insofar as the effector/cytotoxic function of NK cells increase pronouncedly in mice genetically deficient in IL-1R8. Remarkably, these

mice are resistant to the induction of hepatocellular carcinoma (HCC) or hepatic metastases of colorectal or pulmonary cancer. IL-1R8 at least partially exercises its antagonistic effects on the effector functions of NK cells by interfering with the IL-18/IL-18R α /Myd88 signaling pathway. In humans, IL-37, of which the receptor consists in the IL-1R8 and IL-18R α sub-units, is suppressive with regard to NK CD56(dim)CD57(+)Fc ϵ R γ (+)NKG2C(–) cells.

So it is that the cytokine/receptor couples IL-18/IL-1R5 and IL-37/IL-1R8 have opposed effects in the effector/cytotoxic functions of NK/cytotoxic cells.

According to organism site or the tissue, the effects of IL-1R8 favor or disfavor the emergence of infections and/or cancers. For example, IL-1R8 represents not only a protective factor as concerns the development of some colon cancers (11), infectious complications of keratitis (27), and acute *Pseudomonas aruginosa* pneumopathies (28), but also a promoting factor as concerns mammary cancer and HCC (10, 29). One interpretation of these

opposed effects is that in the absence of IL-1R8, an exacerbated inflammatory response (bringing into play the different cell types sensitive to IL-18/IL-33 and IL-37) favors solid tumorigenesis and tissue lesions during infections, while censoring of cancers in liver, particularly hepatic metastases, is partially counteracted by IL-1R8.

IL-18BP (Binding Protein) is also involved and blocks IL-18 action and seems critical as a means of limiting deleterious hepatic NK responses in hepatic aggression models (30, 31) (see Acetaminophen-Induced ALF and Post-Viral Hepatitis A ALF).

IL-1 α / β and Th17 Lymphocyte Differentiation

IL-1 superfamily members are cytokines of critical importance in control of IL-17 production and in differentiation of Th17 lymphocytes and ILC3 (9, 32, 33). This function of IL-1 superfamily members is synergic with that of IL-23. As a result, IL-1R1-deficient mice are lacking in differentiation of Th17 lymphocytes/ILC3 and protected from experimental autoimmune encephalitis induction (34, 35).

The Complex Effects of IL-33 on the Dynamics of Immune Response

- Pro-Th2 effects: IL-33 is a key factor in ILC2 and M2 macrophage differentiation/maturation and also intervenes in Th2 lymphocyte differentiation. From that standpoint, IL-33 has a critical role in the immune responses of mucus epitheliums, as has been demonstrated in mouse models of bronchial hyper-reactivity, pulmonary allergies, and pulmonary response to *A fumigatus* (13). Remarkably, IL-33 on ILC2 induces expression of amphiregulin (Areg), a factor implicated in tissue repair (36, 37).
- The facilitating effects of Th1/cytotoxic/NK cell response: In addition to its effects on type 2 immunity, IL-33 plays a role in control of T cytotoxic and NK cell responses. Associated with IL-12, IL-33 induces IFN γ production by NK cells and seems to have a spectrum of action similar to that of IL-18 (25, 38, 39). According to the timing of stimulation and its chronicity at the level of the bronchial mucus, IL-33 will either favor a type 2 response by means of ILC2 or, on the contrary, induce a pro-Th1 NK cell response, as in the case of chronic lung inflammation due to tobacco in chronic obstructive pneumopathy disease (40). In a context of antiviral response, IL-33 serves as an amplification factor of effector cytotoxic T CD8 cell response by favoring the expansion of short live effector cells (41). Lastly, IL-33, in combination with IL-12, induces IFN γ production by invariant natural killer T (iNKT) cells (38, 42). From this standpoint, we may note the role of the IL-33/iNKT/IFN γ axis in the genesis of renal ischemia-reperfusion (IR) injury in animal models and in humans following transplantation (43–45).
- A role in the recruitment and acquisition of T regulatory lymphocytes' innate competences IL-33 is implicated in the recruitment of T regulatory (Treg) lymphocytes during renal IR injury (46), and is associated with improved renal IR injury tolerance. Furthermore, the eutrophic functions of Treg lymphocytes present in visceral adipose tissue are dependent on the production of IL-33 by the stromal cells in adipose

tissue (47, 48). Lastly, IL-33, and to a certain extent IL-18, are liable to induce Areg production independently of the T cell receptor (TCR) (innate Treg function), thereby contributing to restorative functions, particularly those of the epitheliums, and also favoring tumor promotion and growth (36, 49).

Concept of Trained Immunity and Innate Immunological Memory: the Place of the IL-1 Cytokine Family

A series of recent studies has shown that non-specific stimulation of the immune system (pathogenic agent through *Mycobacterium tuberculosis* or BCG, oxidized low-density lipoproteins, Western diets...) induces modification of the pro-inflammatory immune response program driven by or dependent on myeloid cells (monocytes/macrophages) (19–21, 50, 51). Acquisition of this imprint depends on IL-1 β . This inflammatory emergency hematopoiesis is implicated in the genesis of atherosclerosis lesions, metabolic syndrome, NAFLD or induction of tumor transformation, in particular by a promoting effect of epithelial-mesenchymal transition (EMT). It is the demonstration that this phenomenon leads to the constitution of an innate memory of cells of myeloid origin that led to the proposal of the new concept of “trained immunity.” We have attempted here to apply the concept of trained immunity to liver diseases, based on the role of the IL-1 superfamily, first of all IL-1 β but also IL-18 and IL-33, in modulating innate lymphoid immunity carried by NK cells, ILC or innate T- $\alpha\beta$ lymphocytes.

CONTRIBUTION OF IL-1 SUPERFAMILY OF CYTOKINES TO HEPATIC DISEASES

Constitutive and Inducible Expression of Members of the IL-1 Superfamily in the Liver

In steady-state, the liver expresses the IL-1 α and IL-33 cytokines/alarmins, IL-18 cytokine, the receptors functioning as decoys or inhibitors of these three cytokines: IL-1Ra, IL-18BP, and IL-1R2. While pro-IL-1 α is expressed in hepatocytes, pro-IL-18 is expressed in Kupffer cells (KC). IL-33 is constitutively expressed in liver vascular endothelial and liver sinusoidal endothelial cells (LSEC) (52). Such constitutive expression may have major implications during liver injuries in which cell death by necrosis is predominant and a rapid immune response is required.

Expression in steady-state in liver also occurs for anti-inflammatory factors such as IL-1Ra, which antagonizes IL-1-related functions and modulates a variety of immune and inflammatory responses (53) or IL-18BP, of which the spontaneous production neutralizes locally produced basal IL-18. The sources of IL-18BP in the liver seem to be the hepatocytes themselves, KC and hepatic stellate cells (HSC) (31). The soluble form of IL-1 receptor accessory protein (IL-1RAcP) is constitutively produced by the liver and forms a complex with the

soluble IL-1RII, which binds and neutralizes IL-1 β (8, 54). IL-33 and ST2 are also constitutively expressed in the normal liver (55).

Under pathological conditions that drive liver damage, IL-1 superfamily members are up-regulated. IL-1 and IL-1Ra/IL-1R1 are expressed in almost all cells, especially in activated non-parenchymal cells such as KC. As an alarmin, IL-33 is rapidly released from LSEC and vascular endothelial cells following carbon tetrachloride or Concanavalin A (ConA)-induced hepatitis (52) to mediate a pro-inflammatory response. In response to these danger signals, IL-33 may also be newly secreted in hepatocytes and/or HSC. Its specific receptor ST2 (IL-1 receptor-like 1) and co-receptor IL-1RAcP constitutively expressed on innate immune cells contribute to a rapid immune response.

Acute Liver Failure

Acute liver failure (ALF) is defined as the rapid development (within days or weeks) of severe liver injury with impaired liver function and hepatic encephalopathy (56). ALF is a life-threatening condition and refers to a wide variety of causes, among which toxin/drug-induced (usually acetaminophen) liver damage or viral hepatitis (hepatitis A, B, and E) are most common in the United States and Europe (56, 57).

The Contributions of Experimental Models

In vivo studies in animals have shown the existence of mechanisms common to the different ALF models. So it is that cell death by necrosis results in rapid IL-1 α precursor

release, upregulation of IL-1 β and IL-18 leading to tissue injury through the IL-1R/IL-18R-MyD88 pathway (58–60) with massive upregulation of anti-inflammatory molecule IL-1Ra (61). Following ALF, IL-1 α , IL-1 β , and IL-18 all up-regulate the pro-inflammatory process through a dramatic decrease in hepatic inhibitor of kappa B (I κ B) levels and NF- κ B pathway activation, leading to IL-6 and TNF α secretion, which contributes to apoptosis, and ultimately to liver damage and animal death (60).

Lipopolysaccharide-induced ALF

In the study by Tsutsui et al. (62) KC were shown to be the main producers of IL-18, acting not in a caspase-1-dependent manner but upon Fas Ligand stimulation, thereby increasing IFN γ and TNF α production, which causes substantial acute liver damage. Furthermore, Yan et al. demonstrated a correlation between IL-1 and MMP9 expression implicated in extracellular matrix degradation, sinusoidal collapse, leading to parenchymal cell death, and loss of liver function after ALF induction (63). Inhibition of the IL-1 pathway (using adenovirus IL-1Ra) before the induction of ALF led to a significant reduction in plasma levels of hepatic enzymes and to animal survival improvement (64). Lastly, in humans, treatment by recombinant human IL-1Ra (Anakinra) improved survival of patients with acute liver injury in a post-septic situation (65).

Acetaminophen-induced ALF

In the acetaminophen (acetyl-*p*-Aminophenol, APAP)-induced acute liver injury model, KC were shown to be the




FIGURE 2 | IL-1 α / β and IL-18 members are involved in ALF. **(A)** Acetaminophen-induced ALF. NAPQI (N-acetyl-*p*-benzoquinone-imine) is a toxic acetaminophen metabolite generated by hepatic cytochrome Cyp2e1 and Cyp1a2. Under NAPQI action, injured and necrotic centrilobular hepatocytes secrete DAMP. Among them, ATP and TLR9 receptor ligand induce IL-1 β and IL-18 production by Kupfer cells. Both cytokines drive NK and iNKT cell killing function through FasL and IFN- γ production. This inflammatory process can be blocked either by endogenous IL-18BP production by non-damaged hepatocytes, Kupfer and stellate cells, or by recombinant IL-18BP treatment. **(B)** Critical role of IL-18 in HAV fulminant hepatitis. HAV-infected hepatocytes produce IL-15 as well as PAMP and DAMP which, in turn, activate IL-18 production by Kupfer cells. IL-18 activates IFN- γ production by NK cells and CD8 T-cells with innate functions, hence promoting cytotoxicity toward infected hepatocytes. In healthy infected subjects (left), endogenous IL-18BP blocks the deleterious IL-18 action and modulates NK cell and probably CD8 T-cell activation. In IL-18BP deficient subjects (right), there is no inhibition of the deleterious effects of NK cell and CD8 T-cell activation. ALF, acute liver failure; KC, Kupfer cells; NK, natural killer; Hep, hepatocyte; bounded arrow, presumably yes.

main producers of IL-1 β and IL-18, in a TLR9-dependent induction and NALP3-ASC-caspase-1-dependent manner increasing/inducing IFN- γ and TNF- α production by Th1 and NK cells, which causes substantial acute liver damage (30, 62, 66) (**Figure 2A**). Overexpression of IL-18, known to induce FasL expression on NK cells and CD4(+) T cells, also increased expression of Fas on hepatocytes, which were thereby sensitized to NK cytotoxicity. These Fas/Fas Ligand interactions induced hepatocyte apoptosis, massive periportal fibrosis, inflammation, and severe liver failure (67). Remarkably, IL-18-deficient mice were resistant to ALF induction by APAP, and blocking of IL-1 β by a neutralizing antibody reduced ALF severity (68). Moreover, using the same model, it was recently shown that treatment of mice by IL-18BP, which blocks the binding of IL-18 to its receptor, provided protection from induction of hepatic lesions, thereby confirming the critical role of IL-18/IL-18R-NK cells (30).

To further explore the mechanisms involving the IL-33/ST2 axis in ALF, Antunes et al. (69) used a mouse model of APAP-IL (-induced liver) failure and showed that liver necrosis was associated with massive IL-33 and chemokine release. Non-parenchymal liver cells were the main sensors of IL-33, and the inflammatory response triggered by IL-33 was amplified by liver PMN infiltration. In this model, IL-33 may have acted in synergy with IL-18 in the recruitment of NK cells.

IL-36 is induced after treatment with acetaminophen, presumably in hepatocytes, and up-regulates chemokine ligand 20, a chemokine implicated in tissue protection and repair. Administration of the antagonist IL-36Ra aggravates liver damage and disturbs tissue recovery (70).

Poly-I:C-induced ALF

In a mouse model of ALF induced by poly(I:C), IL-33 expression was up-regulated and correlated with severe liver injury. Interestingly, iNKT cell-deficient mice exhibited protection against poly(I:C)-induced hepatitis accompanied by an increased number of IL-33-positive hepatocytes compared with wild-type (WT) controls (52, 71).

Liver IR injury

Ischemia, followed by reperfusion at the time of liver graft implantation, leads to what is called IR injury. Understanding of the mechanisms involved could help to identify novel therapeutic targets in view of improving organ survival. In mice and rats, warm ischemia followed by reperfusion represents the most common model to study IR injury.

On the one hand, IL-1 β and the NLRP3 inflammasome are implicated in the lesions of warm IR injury *via* high-mobility group box 1, NF- κ B and toll-like receptor (TLR)4 (72). Reactive oxygen species (ROS) mediate inflammasome activation in KC (73). On the other hand, it has been suggested that NLRP3 could be involved in IR injury independently of inflammasomes through PMN recruitment (74). Sadatomo et al. (75) showed in a murine model that macrophages secreted pro-IL-1 β , which is activated by neutrophils-derived proteinases.

Interaction between neutrophils and macrophages promoted IL-1 β maturation and causes IL-1 β -driven inflammation in the IR liver.

Regarding the role of IL-33 and its receptor ST2 in warm IR injury, Yazdani et al. (76) demonstrated in a warm IR injury murine model (confirmed in human liver resection specimens) that IL-33 release from LSEC increases sterile inflammation and results in PMN extracellular trap formation, while administration of recombinant IL-33 during IR exacerbates hepatotoxicity and inflammation. One may note that iNKT cells mediate hepatic IR injury by promotion of intrahepatic PMN influx (77), similarly to what has been shown in models of IR injury involving other organs, such as the kidney (43). Interestingly, preconditioning with intra-peritoneal injection of IL-33 (55) showed that IL-33 has a protective effect on hepatocytes, with decreased liver IR injury *via* the activation of NF- κ B, p38 MAPK, cyclinD1, and Bcl-2. Sakai et al. (78) used a mouse model of IR with injection of recombinant IL-37 at the time of reperfusion. They demonstrated that IL-37 protects against IR injury by reducing pro-inflammatory cytokine and chemokine production by hepatocytes and KC, and suppression of PMN activity.

Observational Clinical Studies

The pro-inflammatory/anti-inflammatory balance of IL-1 family members determines the clinical outcome

Clinical observations first revealed the contribution of the IL-1 family of cytokines to the pathophysiology of ALF. Significant increase in serum levels of the pro-inflammatory cytokine IL-1 α (79), IL-1 β (80), IL-18 and its activator Caspase-1 (81), IL-33 (82) positively correlates with acute hepatitis in humans. Both highly significant elevation in IL-1 α levels (79) and reduction in the IL-1Ra over IL-1 β ratio (IL-1Ra/IL-1 β) (80) have been detected in ALF patients with a fatal outcome. These findings show that the IL-1 family members brought into play are systematically associated in the event of a fatal outcome with loss of balance in the liver between inflammatory signals IL-1 α / β /IL-18 and anti-inflammatory IL-1Ra.

Upregulation of IL-33 and sST2 in serum has been shown to exist in ALF and acute-on-chronic liver failure in patients (82), correlating with the intensity of necrosis as assessed by transaminase activity. Interestingly, sST2 was elevated in acute and acute-on-chronic liver failure, but not in chronic liver failure, suggesting that sST2 could be a tool to monitor the course of the disease.

Post-viral hepatitis A ALF

Less than 1% of acute hepatitis A virus (HAV) infections result in acute fulminant hepatitis. A few cases of post-AHV ALF family isolates have been reported, suggesting the role of a genetic factor implicated in the determinism of this pathology. The study by Belkaya et al. (31) on an 11-year-old patient having died from fulminant HAV hepatitis, with no previous personal or familial medical history, identified a deletion of 40 nucleotides of the gene coding IL-18BP, leading to instability at the mRNA level, and loss of the normal structure, or the absence of IL-18BP.

During the acute phases of viral hepatitis, particularly HAV, massive secretion in patients' serum of IL-18 and IL-15 has

been observed (83). Interestingly, the secretion is accompanied by the appearance of a cell population consisting in innate CD8 T lymphocytes (whose expansion and cytotoxic function are independent of TCR engagement) in the blood and the liver (83). It is highly likely that acquisition of this “innate memory” phenotype depends, as has been described above (section Cytokines of the IL-1 Superfamily and Control of T Helper 1 Effectors and NK Cells), on the joint action of IL-18 and IL-15. Belkaya et al. (31) went on to show that in the patient having succumbed to post-HAV ALF, the aforementioned absence of IL-18BP was accompanied by excessive activation of NK cells, activation that depended on IL-18. On the basis of their analysis of this clinical case, the authors hypothesized that the IL-18BP deficiency could partially explain the fulminant viral hepatitis (**Figure 2B**).

To sum up, the different findings from experimental models and from clinical situations involving ALF underline the importance of the implication of NK/T CD8 cells by means of IL-18 and IL-1 and confirm the decisive role of “negative” regulatory elements (IL-1R8, IL-18BP, IL-37) whose appearance attenuates or reduces the severity of acute hepatic inflammation (**Figure 2**).

Alcoholic Liver Diseases

Alcoholic liver diseases (ALD) includes liver manifestations due to alcohol overconsumption: fatty liver, alcoholic hepatitis, and chronic hepatitis with liver fibrosis or cirrhosis.

Alcohol acts as an “exogenous signal” on KC through TLR with activation of the inflammasome NLRP3-caspase 1 and production of IL-1 β . A second “endogenous signal” leading to activation of the same pathway, is the release of two DAMP: ATP and uric acid by damaged hepatocytes

because of alcohol damage (84). Upregulation of IL-1 β activity leads to inflammation, steatosis, and additional damage (85, 86) (**Figure 3**).

Recently, Heo et al. (87) demonstrated in human and murine liver samples that alcohol decreases microRNA (miR)-148a through the transcriptional regulator forkhead box protein O1. TXNIP (thioredoxin-interacting protein) is a direct target of miR-148a and is overexpressed during alcoholic liver disease, activating NLRP3 inflammasome in hepatocytes, and caspase 1-mediated pyroptosis. Hepatocyte-specific delivery of miR-148a to mice abrogates alcohol-induced TXNIP overexpression and inflammasome activation (**Figure 3**).

In human patients with alcoholic liver disease, sST2 but not IL-33 was correlated to the severity of the disease (88). In the same fashion, Wang et al. (89) showed with IL-33- and ST2- deficient mice that ST2 decreases the inflammatory activation of hepatic macrophages by inhibiting NF- κ B in alcoholic liver disease, in an IL-33-independent manner. However, during severe liver injury, massive cell death, and release of IL-33 triggers IL33/ST2 signaling and increases tissue damage, presumably by local activation of different cells expressing ST2, namely hepatic NK cells, iNKT cells, ILC2, and Treg lymphocytes.

During ALD, one target of IL-1 β , and possibly IL-33, is the iNKT cell population (86). From this standpoint, it is interesting to note that mice genetically deficient in α 18 T cells or CD1d, and consequently without iNKT lymphocytes, are at least partially protected from ALD (90, 91). In these models, iNKT lymphocytes are responsible for PMN influx during ALD progression.




FIGURE 3 | Alcohol acts as an “exogenous signal” on Kupffer cells to activate the inflammasome NLRP3-caspase 1 and production of IL-1 β in ALD. Alcohol decreases microRNA miR-148a, a negative regulator of TXNIP (thioredoxin-interacting protein), which is overexpressed during ALD. TXNIP activates NLRP3 inflammasome in hepatocytes and caspase 1-mediated pyroptosis, leading to IL-1 β and IL-18 production. ATP and uric acid release by damaged hepatocytes after alcohol damage is a second “endogenous signal” leading directly and/or through Kupffer cell activation to inflammatory cytokine production. IL-1 β and IL-18 recruit and activate iNKT cells that, in turn, induce PMN afflux and hepatic injury. IL-33 released by damaged and necrotic hepatocytes exacerbates iNKT cells and PMN recruitment and activation. ALD, alcoholic liver disease; KC, Kupffer cells; iNKT, invariant natural killer T-cells; PMN, polymorphonuclear neutrophils; Hep, hepatocyte.

Non-alcoholic Fatty Liver Disease


With prevalence of 30% for adults and 10% for children in Western countries, non-alcoholic fatty liver disease (NAFLD) is a growing public health issue frequently associated with morbid obesity, type 2 diabetes, and metabolic syndrome (92, 93). NAFLD is characterized by excessive fat accumulation in the liver associated with insulin resistance, and is defined by the presence of steatosis in hepatocytes. NAFLD ranges from simple steatosis or non-alcoholic fatty liver to a more serious form called non-alcoholic steatohepatitis, which is characterized by liver inflammation and tissue damage that can lead to fibrosis, cirrhosis, and hepatocellular carcinoma (94).

There is increasing evidence that the induction of inflammation and production of inflammatory mediators released from the adipose tissue of obese subjects, such as adipocytokines and classical cytokines contribute to

obesity-induced NAFLD, in which IL-1 α / β plays a key role (Figure 4) (95, 96). In fact, NAFLD is the last step of low-grade inflammation depending on the bringing into play of “trained immunity”.

Free fatty acids, acting as DAMP, are capable together with TLR ligands present in gut microbiota of activating NLRP3 and NLRP6 inflammasomes, thereby inducing caspase-1 activation and the release of IL-1 α , IL-1 β , and IL-18 in hepatocytes (97, 98). IL-1 α / β , in turn, activates inflammasome, TNF α and IL1 β and IL-18 release in KC (98). Other extra-hepatic cell types such as adipocytes could also be important sources of IL-1 α and IL-1 β during NAFLD associated with obesity (99).

IL-1 β promotes hepatic steatosis by stimulating triglyceride and cholesterol accumulation in primary liver hepatocytes and lipid droplet formation (96) and acts on LSEC to promote liver inflammation by upregulating intercellular adhesion molecule



1 expression, which stimulates neutrophil recruitment in the liver (100). Moreover, IL-1 β produced by the liver, together with pro-inflammatory IL-6 and TNF α , contributes to the activation of resident immune cells and the recruitment of other leucocytes to the damaged liver, leading to chronic inflammation (101). Accordingly, IL-1 β and/or IL-1 α -deficient mice have demonstrated less diet-induced inflammation and liver fibrosis, as attested by lower serum transaminases and serum amyloid alpha concentrations, as well as decreased expression of inflammatory and fibrosis transcripts such as *IL-6*, *TNF α* , *P-Selectin*, *Cxcl1*, and *TGF β* , as compared to WT controls (102).

IL-1 α/β are upregulated in the adipose tissue of obese and insulin-resistant mice. Together with TNF α and IL-6, IL-1 α/β , in turn, contributes to the development of insulin resistance in adipocytes through downregulation of insulin receptor substrate-1 expression, thereby facilitating accumulation of lipids in the liver (103). Moreover, adipocyte-derived IL-1 β has been shown to directly induce, alone or in combination with TNF α , insulin resistance in hepatocytes *via* AMPK-ROS signaling and autophagy (104, 105). Accordingly, IL-1 α and IL-1 β -deficient mice have lower fasting glucose and insulin levels and improved insulin sensitivity (106).

Of note, IL-1 α deficiency increased both hepatic and systemic cholesterol levels, suggesting that liver fat storage and inflammation are not necessarily parallel events and that some hepatic lipids might support anti-inflammatory functions (102). IL-1Ra-deficient mice exhibited severe steatosis and pericellular fibrosis containing many inflammatory cells, as observed in steatohepatitis histological lesions in humans, following 20 weeks of feeding with an atherogenic diet (107). These different observations, particularly as regards the role of IL-1Ra in NAFLD, raise the question of the interest of therapeutic utilization of human recombinant IL-1Ra (Anakinra) in the natural history of NAFLD.

Henao-Mejia et al. (108) have reported that changes of gut microbiota linked to NLRP3 and NLRP6 inflammasome deficiency were associated with increased hepatic steatosis and inflammation. Interestingly, IL-18 production negatively regulates non-alcoholic steatohepatitis progression *via* modulation of the gut microbiota. This observation strongly highlights the role of the maintenance of digestive microbiome homeostasis in the genesis of “metabolic syndrome” and NAFLD (108).

Regarding new members of the IL-1 superfamily, IL-33 has not been widely explored. In a murine model, a high fat diet (HFD)-induced steatohepatitis was associated with upregulation of IL-33 but IL-33 deficiency did not affect severity of liver inflammation or liver fibrosis, suggesting that endogenous IL-33 has no effect on the progression of fibrosis during experimental steatohepatitis (109). Treatment with recombinant IL-33 in mice attenuated diet-induced hepatic steatosis but aggravated hepatic fibrosis in a ST2-dependent manner (110). The pro-fibrotic effect of pharmacological IL-33 treatment in HFD mice was confirmed in another study (111), where galectin-3 enabled upregulation of the IL-33/ST2 pathway and production of IL-13 by peritoneal macrophages.

Autoimmune Hepatitis

Autoimmune hepatitis (AIH) is a chronic inflammatory disease of the liver of which the pathogenic mechanisms have yet to be unraveled. Treatment relies mainly on lifelong immunosuppressive therapy and liver transplantation in the final stage. Acute hepatitis induced with ConA is the most widely used mouse model of AIH (**Figure 5**). In this model, massive activation of immune cells induces massive local production of cytokines such as IL-4, IFN γ , and TNF α (112–114), which are directly implicated in hepatocyte death (113). iNKT lymphocytes are of major importance in acute hepatitis induced with ConA, as is attested in works showing that iNKT lymphocyte-deficient mice resist hepatitis induction (115). Aside from their cytokine production, the pathogenic effect of iNKT lymphocytes depends on their cytotoxic functions, which bring into play perforin and granzymes as well as FasL/Fas and TRAIL/DR5 interactions (115, 116). Depletion of KC prior to ConA treatment attenuates hepatic injury (117). Activation of iNKT lymphocytes may consequently depend on KC, thereby ensuring a role of auto-antigen presentation. Another pathogenic mechanism may consist in activation of KC by iNKT lymphocytes, leading to production by the KC of TNF α , ROS and IL-1 β depending on the activation by ROS of the NLRP3 inflammasome (118). On this subject, it is interesting to note that in this model, treatment of mice by human recombinant IL-1Ra (Anakinra) at least partially prevents hepatic lesions (118).

IL-33 is rapidly neosynthesized (6 hours) in the hepatocytes after treatment by ConA (116, 119) (**Figure 5**). Interestingly, this expression of IL-33 depends on the presence of iNKT lymphocytes and possibly on TRAIL/DR5 interactions between iNKT lymphocytes and hepatocytes (116). IL-33 production seems to moderate hepatitis severity, because: (i) IL-33-deficient mice have more heavily damaged livers than WT mice (52, 71, 116, 119, 120); (ii) injection of recombinant IL-33 renders less severe the hepatitis induced by ConA (120). Biologically, the increased severity of hepatitis lesions in IL-33-deficient mice is associated with pronounced expression of TNF α and IL-1 β and is accompanied by accumulation in the liver of activated NK cells (119).

The protective action of IL-33 in the model of ConA-induced hepatitis may be attributed to the induction of anti-apoptotic factors such as BAX or Bcl2 (120). Another element associated with the protective action of IL-33 may be the recruitment in the liver of Treg lymphocytes [TCR- $\alpha\beta$ (+)CD4(+)CD25(+)FoxP3(+)] ST2(+) (119). One is tempted to suggest that these Treg ST2(+) cells contribute to hepatic repair by expression of trophic factors such as Areg, as previously described in other tissues (36, 121).

Viral Hepatitis B, C

According to the World Health Organization, 325,000,000 persons are currently living with a chronic infection of either hepatitis virus B (HBV) or hepatitis virus C (HCV). Infection with HBV rarely leads to acute liver failure. Chronic infections with both HBV and HCV are responsible for progressive fibrosis and cirrhosis, as well as development of hepatocellular carcinoma.




FIGURE 5 | The ConA model of hepatitis. The mitogen Concanavalin A (ConA) activates iNKT cells in a KC and iNKT cell-dependent mode. IL-4, IFN γ , and TNF α produced by hepatic activated iNKT cells activate the NLRP3 inflammasome leading to toxic IL-1 β production by hepatocytes. iNKT cells could also induce IL-33 production by hepatocytes through TRAIL/DR5 interactions. Secreted IL-33 acts as a retro-negative regulator of iNKT cells by recruiting Treg. Indeed, Treg can restrain T-cell cytotoxicity and promote repair through signals like Areg. ConA, concanavalin A; Areg, Amphiregulin; KC, Kupffer cell; Hep, hepatocyte.

IL-1 β and IL-18 may play a role in liver inflammation in the setting of HCV infection. Monocyte-derived human macrophages and Kupffer cells (122) produce IL-1 β and IL-18 in response to HCV infection: HCV induces pro-IL-1 β and pro-IL-18 in macrophages *via* the NF- κ B signaling pathway. It has been shown that viral RNA can trigger MyD88/TLR7 pathway, thereby inducing IL-1 β expression. HCV concomitantly activates the NLRP3 inflammasome, leading to IL-1 β secretion (123).

It has been shown in humans that with their IFN γ production, iNKT lymphocytes inhibit the replication of HCV in hepatocytes (124) and that over the course of HBV infection, recruitment of iNKT lymphocytes occurs at the level of the liver (125–127). Lastly, in a model of HBV-transgenic mice, the authors showed that IL-18 induces repression of viral replication in a context depending on local recruitment of NK cells and iNKT lymphocytes (128). Interestingly, when infection becomes chronic, hepatic iNKT lymphocytes are progressively

reduced in number (127), conceivably resulting in a loss of sensitivity of iNKT lymphocytes to IL-18 and/or IL-33.

From this standpoint, IL-33 expression in LSEC significantly increases in patients with chronic HBV (CHB) infection, and is likewise highlighted in the hepatocytes of lesions/inflammatory foci in patients with the most severe form of hepatitis (129). Seric IL-33 and ST2s concentrations are highly elevated compared to those in healthy donors and, interestingly enough, they are correlated with the severity of hepatic cytolysis (129, 130). In HBV acute-on-chronic liver failure (ACLF) (129), seric IL-33 and ST2s concentrations are pronouncedly higher than the concentrations observed in CHB patients. The circulating monocytes in ACLF patients present a highly activated and pro-inflammatory phenotype, which is correlated with IL-33 production. For that reason, it has been suggested that IL-33 increases inflammation and disease severity through monocyte activation, TNF α , IL-6, and IL-1 β secretion. Similarly, IL-33

levels in human patients correlate with HCV RNA and liver damage (131). Such IL-33 production, particularly in the early phases of infection, can act in synergy with IL-18 and IL-1 α/β in the recruitment of NK, iNKT, and T cytotoxic cells. And given the chronicity of its production, IL-33 is also liable to induce a state of systemic inflammation and, locally, a phenomenon of immunosubversion through recruitment of Treg lymphocytes and ILC2 as opposed to recruitment and activation of NK and iNKT cells.

Fibrosis

Liver fibrosis, the final stage of most types of chronic liver diseases, characterized by the excessive accumulation of extracellular matrix proteins, including collagen, is a consequence of extracellular matrix hepatic stem cell activation. Gieling et al. (132) established that IL-1 activates hepatic stem cells that produce MMP-9 and control the progression from liver injury to fibrogenesis. Advanced liver fibrosis results in cirrhosis with progressive liver failure and development of portal hypertension.

Marvie et al. (133) showed that in chronic hepatitis, in the model induced by CCl₄, IL-33 and ST2 expression correlates with fibrosis (133). In humans suffering from chronic hepatitis, there also exists at the level of the liver a correlation between IL-33 cellular expression and fibrosis severity (133). In models of fibrosis induced by CCl₄, thioacetamide, and bile duct ligation, increased hepatic IL-33 expression in activated HSC has been confirmed. Remarkably, fibrosis lesions are absent or non-severe in IL-33-deficient mice, thereby underscoring the major role of IL-33 in the manifestations of fibrosis (134). Lastly, this study suggests that the profibrosis function of IL-33 depends on its action in the activation and expansion of liver ILC2 (134). In the absence of IL-13 produced by ILC2, CCl₄ does not induce hepatic fibrosis.

The fibrosis lesions described in the aforementioned works depend on the impact of environmental, chemical or physical stress. It may be the case that in the induction of hepatic fibrosis coming about during NAFLD, IL-33 is non-indispensable. This hypothesis is premised on the fact that in a model of hepatic fibrosis induced by HFD, there is no difference in fibrosis severity between WT and genetically IL-33-deficient mice (109).

Hepatocellular Carcinoma (See Figure 6)

As described in section Cytokines of the IL-1 Superfamily and Control of T Helper 1 Effectors and NK Cells, the IL-18/IL-18R axis is a checkpoint driven by the immunological components controlling carcinogenesis and hepatic metastases. In addition to NK lymphocytes, iNKT lymphocytes are of key importance in the censoring of hepatic metastases (135, 136). As a result, iNKT lymphocytes strongly express IL-18 and IL-33 receptors (38, 39, 137) and may consequently be implicated in the anti-tumoral effect of the IL-18/IL-18R axis. Moreover, influx in the liver of iNKT lymphocytes depends on the expression of CXCL16 by LSEC, that is to say cells of which another characteristic is nuclear expression of IL-33. The study by Ma et al. (135) shows that recruitment of anti-tumoral iNKT lymphocytes in the liver is indirectly controlled by the digestive microbiota and by the metabolites produced by the latter, metabolites that will induce


CXCL16 expression by LSEC. From this standpoint, it should be pointed out that there exists an accumulation of iNKT CD4(+) cells with a type Th2 profile in HCC liver (138). Enriched with iNKT cells, its infiltration is correlated with HCC progression, attesting to the loss of effectiveness of the anti-tumoral activity of iNKT lymphocytes in a tumor niche during the natural history of HCC.

Several works in mouse models have shown that endogenous IL-33 is an adjuvant factor in anti-tumoral and anti-metastatic immunity, including at the level of the liver (139–141). The limitation of these studies resides in the utilization of metastasis models, some of which use HCC cell lines but do not reproduce the natural history of HCC.

Even though HCC may develop in healthy liver parenchyma, in most cases it develops in chronic inflammatory liver parenchyma in pathology secondary to the induction of trained immunity depending on IL-1 β production (NAFLD, ALD, Viral chronic hepatitis, hepatic fibrosis). As we previously discussed, the liver possesses its own means of induction and control of oncogenesis. Consequently, the process of transformation/oncogenesis at the hepatic level is partially dependent on IL-1 β and will avail itself of amplification paths.

Recently, Zhang et al. (142) demonstrated in cell cultures, mouse models, and human samples the existence of a hypoxia-inducible factor (HIF)-1 α /IL-1 β signaling loop between cancer cells and tumor-associated macrophages (TAM) in a hypoxic microenvironment, resulting in cancer cell epithelial-mesenchymal transition and metastasis. When exposed to moderate hypoxia, HIF-1 α accumulation up-regulates IL-1 β secretion in TAM. IL-1 β increases inflammatory signaling and enhances epithelial-mesenchymal transition in cancer cells through the IL-1 β /HIF-1 α /COX2 axis, which promotes the invasive capacities of tumoral cells in a hypoxic microenvironment.

IL-37 is an anti-inflammatory cytokine, which in numerous models seems to block the deleterious effects of IL-1 α/β and IL-18. Schematically, this cytokine counteracts the induction of trained immunity. From this standpoint, it is altogether remarkable that hepatic expression of IL-37 decreases during HCC evolution and that it does so both in the tumor itself and the non-tumoral tissue adjacent to the tumor. Factors suggesting that IL-37 may possess an anti-tumoral effect are corroborated by the following points: (i) persistence of hepatic expression of IL-37 is associated during HCC with infiltration of NK CD57(+) cells. *In vitro*, the HCC cell lines expressing IL-37 attract NK CD57(+) cells which, in turn, are cytotoxic with regard to the HCC lines (143); (ii) the profusion of NK CD57(+) cells and the elevated level of hepatic expression of IL-37 are associated with improved overall survival (143); (iii) IL-37 has an anti-tumoral effect during HCC. While the JNK/pSmad3L/c-Myc pathway is associated with tumor growth during HCC, the signaling depending on IL-37 diverts pSmad3L from that platform and associates it with the suppressor of tumor p21 in a pSmad3C/p21 complex. The presence of pSmad3L is inversely correlated with overall survival (144); (iv) IL-37 is liable to induce autophagia in the HCC cell lines by inhibiting the PI3K/Akt/mTOR signaling pathway, thereby sensitizing HCC cells to apoptosis induction (145).



As regards IL-33, data from the natural history of HCC are available in humans. Yang et al. (146) detected IL-33 protein expression by immunohistochemistry in patients with HCC, liver cirrhosis, hepatitis, and normal livers. They showed that expression of IL-33 is increased in HCC and LC patients, as compared to healthy donors. In HCC patients, IL-33 is expressed by both tumoral cells and peri-tumoral tissue. Immunohistology results in this work are corroborated by quantitative RT-PCR data. A second study (147) likewise underlines IL-33 expression at the level of the tumor during HCC. IL-33 seric expression is particularly pronounced when the disease has evolved, whereas high seric concentrations of ST2s have been found to be associated with reduced overall survival of patients with HCC (148).

High expression of IL-33 could be a factor contributing to deregulation of local response, possibly increased by desensitization of iNKT and NK lymphocytes as well as the recruitment of Treg lymphocytes and ILC2 producers of AReg, a cytokine of which the function in transformation and progression of HCC has been documented (149).

THERAPEUTIC PERSPECTIVES

Although the role of some members of the IL-1 cytokine family in liver disease has been extensively studied (IL-1 α , IL-1 β , IL-33), leading to strong arguments favoring these molecules as potential therapeutic targets, the role of the other members (IL-36, IL-37, IL-38) remains to be elucidated. Furthermore, some members of the IL-1 family are pro-inflammatory and enhance tissue damage and inflammation, while other members are more protective/anti-inflammatory by promoting tissue regeneration and preventing tissue damage and inflammation. Hence, these cytokines represent crucial targets for liver disease therapies and could open new perspectives of potentially innovative therapeutic approaches aimed at controlling local immune response and at limiting liver injury.

Members of the IL-1 superfamily have multiple cellular sources and targets, as well as numerous natural inducers and inhibitors. The pharmacological agents that can either suppress cytokine production or block their biological actions may have potential therapeutic value against a wide variety of liver diseases.

Moreover, several members of the IL-1 cytokine family can activate (or inhibit) the same receptors. For example, IL-1RAcP is shared by IL-1 α , IL-1 β , IL-33, and IL-36, highlighting redundancy and compensatory actions. Since IL-1 α , IL-1 β , IL-33, and IL-36 are implicated *in vivo* during liver diseases, the efficiency of the therapeutic approaches based on targeting a cytokine by the other members may be affected. In other words, when targeting the shared receptors, in this case IL-1RAcP may be more effective. In addition to redundancy, some cytokines in the IL-1 superfamily are known for their duality of function. For example, IL-33 has shown both pathological and hepatoprotective functions and rIL-33-ST2 therapeutic possibilities seem to depend on the presence of different inflammatory conditions, which may require either activation or inhibition of this pathway.

Modulation of IL-1 β and IL-18 by Interference With Inflammasome

While some pharmacological agents against the NLRP3–caspase-1 pathway have been developed, their efficiency regarding cytokine inhibition and liver inflammation is moderate (106, 150, 151). This may be explained by the implication of additional mechanisms of extracellular activation of the cytokines. In addition to the classical NLRP3 inflammasome–caspase-1 cytokine activation pathway, serine proteases from immune cells (Figure 1) are also potent cytokine activators contributing to liver disease progression. The use of agents acting against those serine proteases, combined or not with anti-inflammasome–caspase-1 therapy, can be considered as a valuable therapeutic strategy and merits further investigation (152).

Manipulation of the IL-1/IL-1R Pathway in Chronic Inflammatory Hepatopathies

IL-1 β is a critical mediator of trained immunity and chronic inflammatory hepatopathies such as NASH and NAFLD, raising the question of the therapeutic targeting of the IL-1/IL-1R axis.

Anakinra is the recombinant form of IL-1Ra used in treatment of chronic or acute inflammatory pathologies (Muckle-Wells, adult-onset Still's disease). It effectively treats macrophage activation syndrome (MAS), as regards both MAS associated with inflammatory pathologies such as Still's disease and MAS associated with septic shock (for a review, see 5, 63). Altogether remarkably, in the latter indication, Anakinra clearly improves survival of patients with disseminated intravascular coagulation or hepatobiliary injury (65).

Several studies have suggested that utilization of Anakinra in precancerous states slows evolution of the disease, particularly in cases of smoldering myeloma (associated with Dexametason) (153). Moreover, it is a treatment that seems beneficial in management of some cancers, particularly metastatic colorectal cancer (154). It is also remarkably well-tolerated, with relatively few adverse effects, especially infectious [for a review, see (8)].

Blockage of the IL-1/IL-1R pathway can also be achieved with a neutralizing antibody directed against IL-1 β , Canakinumab, which seems beneficial in atherosclerosis prevention in high-risk patients (155).

Most of these diseases involve inflammatory pathways depending on the bringing into play of trained immunity, of which we previously noted the role/implication in the genesis of different hepatopathies. While blockage of the IL-1/IL-R pathway is a therapeutic option that may be envisioned in cases of chronic inflammatory hepatopathies, conclusive demonstration of its effectiveness has yet to be provided.

IL-18BP in ALF and Acute Viral Hepatitis

Recent demonstration of the role of the IL-18/NK cell axis in ALF genesis, particularly in cases of viral origin (31), has underscored the critical role of negative regulatory signals characterizing the pathway: IL-1R8 and IL-18BP. From this standpoint, in terms of therapeutic manipulation, IL-18BP is an excellent candidate. A human recombinant form (Tradekinig Alfa, AB2 Bioltd) is currently under evaluation in treatment of Still's disease and hemophagocytic lymphohistiocytosis syndrome, that is to say a pathology in which the IL-18/IL-18R axis seems preponderant. It has also been suggested that IL-18BP can prevent acetaminophen-mediated hepatotoxicity (30, 68). These different studies raise the issue of pharmacological targeting of IL-18 in a wide range of ALF cases.

Recombinant IL-233

In situations of acute suffering due to IR injury, targeting of the IL-33/ST2 pathway is an option of genuine interest that nonetheless raises questions of intervention timing. Its application in organ preconditioning could represent a protective strategy, given that during an IR injury phase IL-33 is likely to be implicated in the appearance of lesions.

Another strategy aimed at favoring the recruitment of Treg lymphocytes during an IR injury sequence consists in the utilization of recombinant IL-233. This chimeric IL associates IL-2 and IL-33 functions: the N-terminal residues 21–169 of the IL-2 receptor are linked by 15 amino-acyl residues to C-terminal residues 109–266 of the IL-33 receptor (156). Recombinant IL-233 enables preferential recruitment of T lymphocytes and ILC2 at the expense of iNKT lymphocytes, NK cells and ILC1. Stremaska et al. (156) have shown that in a renal IR injury model, recombinant IL-233 is “formidably effective” as a means of protecting mice at post-sequence IR injury. In addition to a modified balance of recruitment between pro-inflammatory effectors (iNKT/T/NK/PMN) and anti-inflammatory effectors (Treg lymphocytes/ILC2), it is altogether probable that in this model, IL-233 promotes rapid, and effective repair signals. It is consequently relevant to ask whether this molecule may be beneficially utilized in cases of acute hepatic inflammation, especially those associated with liver transplantation.

Even though it currently remains speculative, future therapeutic utilization of recombinant IL-233 in cases of chronic inflammatory hepatopathies merits consideration. A recent study highlighted the beneficial effects of IL-233 in prevention of diabetic nephropathy in an Ob type 2 diabetes (DT2) mouse model. In this mouse model of predisposition to obesity and DT2 on account of an Ob mutation of leptin rendering the latter non-functional, IL233 treatment during or after the appearance of DT2 provides protection from diabetic nephropathy (46).

These results give rise to the question of the possible interest of a “humanized” form of this molecule in the prevention of long-term consequences of trained immunity, especially as regards the liver. However, other questions arise: Might there not ensue an effect stemming from loss of cancer censoring and, eventually, promotion of tumor growth depending on the Treg/ILC2 lymphocytes expressing Areg?

Conclusion Concerning Therapeutic Manipulation of Cytokines in the IL-1 Family

Further *in vivo* studies are needed to understand the potential benefits of targeting these shared receptors or simultaneously blocking multiple members of the IL-1 cytokine family.

Furthermore, these cytokines require enzymatic or proteolytic processing to become active, and their activators contribute to the process of liver inflammation and can therefore be potential therapeutic targets.

In conclusion, acute and chronic inflammations of the liver have various etiologies, ranging from toxic aggression (alcohol, fat diet, drugs) to viruses. Chronic liver inflammation leads to fibrosis and end-stage cirrhosis, and predisposes to the development of neoplasms such as hepatocellular carcinoma. The different members of the IL-1 superfamily of cytokines, which contribute to control of tissue homeostasis and help the liver to respond to damage and disease by promoting immune responses, have a major role in most

liver inflammatory diseases and may represent potential clinical targets.

AUTHOR CONTRIBUTIONS

AR, CS, and ES contributed to literature search and editing of the review. LB, MF, AB, J-MG, and AH contributed to literature search for the review and provided writing and editing of the review.

FUNDING

This study was supported by INSERM, CHU de Poitiers, Université de Poitiers, Ligue contre le Cancer (Comités départementaux de la Vienne, de la Charente, de la Charente Maritime et des Deux-Sèvres), and the Association pour la Recherche en Immunologie-Poitou-Charentes (ARIM-PC). MF and AB were supported by fellowships provided by INSERM, Région Poitou-Charentes, Octapharma, Fondation Bristol-Meyers Squibb, and Région Nouvelle Aquitaine, respectively.

ACKNOWLEDGMENTS

The authors would like to acknowledge Jeffrey Arsham, a professional American medical translator, who reviewed the manuscript. Figures used images from Sevier Medical Art (<https://smart.servier.com/>).

REFERENCES

- Menkin V. Studies on the isolation of the factor responsible for tissue injury in inflammation. *Science*. (1943) 97:165–7. doi: 10.1126/science.97.2511.165
- Beeson PB. Temperature-elevating effect of a substance obtained from polymorphonuclear leucocytes. *J Clin Invest*. (1948) 27:524.
- Atkins E, Wood WB Jr. Studies on the pathogenesis of fever. II. Identification of an endogenous pyrogen in the blood stream following the injection of typhoid vaccine. *J Exp Med*. (1955) 102:499–516. doi: 10.1084/jem.102.5.499
- Dayer JM. From supernatants to cytokines: a personal view on the early history of IL-1, IL-1Ra, TNF and its inhibitor in rheumatology. *Arthritis Res Ther*. (2018) 20:1–6. doi: 10.1186/s13075-018-1607-y
- Dinarello CA. Interleukin-1. *Cytokine Growth Factor Rev*. (1997) 8:253–65. doi: 10.1016/S1359-6101(97)00023-3
- Dinarello CA. IL-1: discoveries, controversies and future directions. *Eur J Immunol*. (2010) 40:599–606. doi: 10.1002/eji.201040319
- Tsutsui H, Cai X, Hayashi S. Interleukin-1 family cytokines in liver diseases. *Mediators Inflamm*. (2015) 2015:630265. doi: 10.1155/2015/630265
- Dinarello CA. Overview of the IL-1 family in innate inflammation and acquired immunity. *Immunol Rev*. (2018) 281:8–27. doi: 10.1111/imr.12621
- Mantovani A, Dinarello CA, Molgora M, Garlanda C. Interleukin-1 and related cytokines in the regulation of inflammation and immunity. *Immunity*. (2019) 50:778–95. doi: 10.1016/j.immuni.2019.03.012
- Molgora M, Bonavita E, Ponzetta A, Riva F, Barbagallo M, Jaillon S, et al. IL-1R8 is a checkpoint in NK cells regulating anti-tumour and anti-viral activity. *Nature*. (2017) 551:110–4. doi: 10.1038/nature24293
- Molgora M, Supino D, Mantovani A, Garlanda C. Tuning inflammation and immunity by the negative regulators IL-1R2 and IL-1R8. *Immunol Rev*. (2018) 281:233–47. doi: 10.1111/imr.12609
- Moussion C, Ortega N, Girard JP. The IL-1-like cytokine IL-33 is constitutively expressed in the nucleus of endothelial cells and epithelial cells *in vivo*: a novel ‘alarmin’? *PLoS ONE*. (2008) 3:1–8. doi: 10.1371/journal.pone.0003331
- Cayrol C, Girard J-P. Interleukin-33 (IL-33): a nuclear cytokine from the IL-1 family. *Immunol Rev*. (2018) 281:154–68. doi: 10.1111/imr.12619
- Bandara G, Beaven MA, Olivera A, Gilfillan AM, Metcalfe DD. Activated mast cells synthesize and release soluble ST2-a decoy receptor for IL-33. *Eur J Immunol*. (2015) 45:3034–44. doi: 10.1002/eji.201545501
- Hayakawa H, Hayakawa M, Kume A, Tominaga S. Soluble ST2 blocks interleukin-33 signaling in allergic airway inflammation. *J Biol Chem*. (2007) 282:26369–80. doi: 10.1074/jbc.M704916200
- Sanada S, Hakuno D, Higgins LJ, Schreiter ER, McKenzie ANJ, Lee RT. IL-33 and ST2 comprise a critical biomechanically induced and cardioprotective signaling system. *J Clin Invest*. (2007) 117:1538–49. doi: 10.1172/JCI30634
- Bessa J, Meyer CA, de Vera Mudry MC, Schlicht S, Smith SH, Iglesias A, et al. Altered subcellular localization of IL-33 leads to non-resolving lethal inflammation. *J Autoimmun*. (2014) 55:33–41. doi: 10.1016/j.jaut.2014.02.012
- Cohen I, Rider P, Carmi Y, Braiman A, Dotan S, White MR, et al. Differential release of chromatin-bound IL-1 α discriminates between necrotic and apoptotic cell death by the ability to induce sterile inflammation. *Proc Natl Acad Sci USA*. (2010) 107:2574–9. doi: 10.1073/pnas.0915018107
- Bekkering S, Arts RJW, Novakovic B, Kourtellos I, van der Heijden CDCC, Li Y, et al. Metabolic induction of trained immunity through the mevalonate pathway. *Cell*. (2018) 172:135–46.e9. doi: 10.1016/j.cell.2017.11.025
- Mitroulis I, Ruppova K, Wang B, Chen L-S, Grzybek M, Grinenko T, et al. Modulation of myelopoiesis progenitors is an integral component of trained immunity. *Cell*. (2018) 172:147–61.e12. doi: 10.1016/j.cell.2017.11.034
- Moorlag SJCFM, Röhring RJ, Joosten LAB, Netea MG. The role of the interleukin-1 family in trained immunity. *Immunol Rev*. (2018) 281:28–39. doi: 10.1111/imr.12617
- Dupaul-Chicoine J, Arabzadeh A, Dagenais M, Douglas T, Champagne C, Morizot A, et al. The Nlrp3 inflammasome suppresses colorectal cancer

- metastatic growth in the liver by promoting natural killer cell tumoricidal activity. *Immunity*. (2015) 43:751–63. doi: 10.1016/j.immuni.2015.08.013
23. Adams NM, O'Sullivan TE, Geary CD, Karo JM, Amezcua RA, Joshi NS, et al. NK cell responses redefine immunological memory. *J Immunol*. (2016) 197:2963–70. doi: 10.4049/jimmunol.1600973
 24. Madera S, Sun JC. Cutting edge: stage-specific requirement of IL-18 for antiviral NK cell expansion. *J Immunol*. (2015) 194:1408–12. doi: 10.4049/jimmunol.1402001
 25. Nabekura T, Girard J-P, Lanier LL. IL-33 receptor ST2 amplifies the expansion of NK cells and enhances host defense during mouse cytomegalovirus infection. *J Immunol*. (2015) 194:5948–52. doi: 10.4049/jimmunol.1500424
 26. Romee R, Schneider SE, Leong JW, Chase JM, Keppel CR, Sullivan RP, et al. Cytokine activation induces human memory-like NK cells. *Blood*. (2012) 120:4751–60. doi: 10.1182/blood-2012-04-419283
 27. Huang X, Hazlett LD, Du W, Barrett RP. SIGIRR promotes resistance against *Pseudomonas aeruginosa* keratitis by down-regulating type-1 immunity and IL-1R1 and TLR4 signaling. *J Immunol*. (2006) 177:548–56. doi: 10.4049/jimmunol.177.1.548
 28. Véliz Rodríguez T, Moalli F, Polentarutti N, Paroni M, Bonavita E, Anselmo A, et al. Role of Toll interleukin-1 receptor (IL-1R) 8, a negative regulator of IL-1R/Toll-like receptor signaling, in resistance to acute *Pseudomonas aeruginosa* lung infection. *Infect Immun*. (2012) 80:100–9. doi: 10.1128/IAI.05695-11
 29. Campesato LF, Silva APM, Cordeiro L, Correa BR, Navarro FCP, Zanin RF, et al. High IL-1R8 expression in breast tumors promotes tumor growth and contributes to impaired antitumor immunity. *Oncotarget*. (2017) 8:49470–83. doi: 10.18632/oncotarget.17713
 30. Bachmann M, Pfeilschifter J, Mühl H. A prominent role of interleukin-18 in acetaminophen-induced liver injury advocates its blockage for therapy of hepatic necroinflammation. *Front Immunol*. (2018) 9:1–7. doi: 10.3389/fimmu.2018.00161
 31. Belkaya S, Michailidis E, Korol CB, Kabbani M, Cobat A, Bastard P, et al. Inherited IL-18BP deficiency in human fulminant viral hepatitis. *J Exp Med*. (2019) 216:1777–90. doi: 10.1084/jem.20190669
 32. Langrish CL, Chen Y, Blumenschein WM, Mattson J, Basham B, Sedgwick JD, et al. IL-23 drives a pathogenic T cell population that induces autoimmune inflammation. *J Exp Med*. (2005) 201:233–40. doi: 10.1084/jem.20041257
 33. Mills KHG, Dungan LS, Jones SA, Harris J. The role of inflammasome-derived IL-1 in driving IL-17 responses. *J Leukoc Biol*. (2013) 93:489–97. doi: 10.1189/jlb.1012543
 34. Sutton C, Brereton C, Keogh B, Mills KHG, Lavelle EC. A crucial role for interleukin (IL)-1 in the induction of IL-17-producing T cells that mediate autoimmune encephalomyelitis. *J Exp Med*. (2006) 203:1685–91. doi: 10.1084/jem.20060285
 35. Sutton CE, Lalor SJ, Sweeney CM, Brereton CF, Lavelle EC, Mills KHG. Interleukin-1 and IL-23 induce innate IL-17 production from gammadelta T cells, amplifying Th17 responses and autoimmunity. *Immunity*. (2009) 31:331–41. doi: 10.1016/j.immuni.2009.08.001
 36. Arpaia N, Green JA, Molteni B, Arvey A, Hemmers S, Yuan S, et al. A distinct function of regulatory T cells in tissue protection. *Cell*. (2015) 162:1078–89. doi: 10.1016/j.cell.2015.08.021
 37. Monticelli LA, Osborne LC, Noti M, Tran SV, Zaiss DMW, Artis D. IL-33 promotes an innate immune pathway of intestinal tissue protection dependent on amphiregulin-EGFR interactions. *Proc Natl Acad Sci USA*. (2015) 112:10762–7. doi: 10.1073/pnas.1509070112
 38. Bourgeois E, Van LP, Samson M, Diem S, Barra A, Roga S, et al. The pro-Th2 cytokine IL-33 directly interacts with invariant NKT and NK cells to induce IFN-gamma production. *Eur J Immunol*. (2009) 39:1046–55. doi: 10.1002/eji.200838575
 39. Smithgall MD, Comeau MR, Yoon B-RP, Kaufman D, Armitage R, Smith DE. IL-33 amplifies both Th1- and Th2-type responses through its activity on human basophils, allergen-reactive Th2 cells, iNKT and NK cells. *Int Immunol*. (2008) 20:1019–30. doi: 10.1093/intimm/dxn060
 40. Kearley J, Silver JS, Sanden C, Liu Z, Berlin AA, White N, et al. Cigarette smoke silences innate lymphoid cell function and facilitates an exacerbated type I interleukin-33-dependent response to infection. *Immunity*. (2015) 42:566–79. doi: 10.1016/j.immuni.2015.02.011
 41. Bonilla WV, Fröhlich A, Senn K, Kallert S, Fernandez M, Johnson S, et al. The alarmin interleukin-33 drives protective antiviral CD8+ T cell responses. *Science*. (2012) 335:984–9. doi: 10.1126/science.1215418
 42. Bourgeois EA, Levescot A, Diem S, Chauvineau A, Bergès H, Milpied P, et al. A natural protective function of invariant NKT cells in a mouse model of innate-cell-driven lung inflammation. *Eur J Immunol*. (2011) 41:299–305. doi: 10.1002/eji.201040647
 43. Ferhat M, Robin A, Giraud S, Sena S, Goujon J-M, Touchard G, et al. Endogenous IL-33 contributes to kidney ischemia-reperfusion injury as an alarmin. *J Am Soc Nephrol*. (2018) 29:1272–88. doi: 10.1681/ASN.2017060650
 44. Ferhat MH, Robin A, Barbier L, Thierry A, Gombert J-M, Barbarin A, et al. The impact of invariant NKT cells in sterile inflammation: the possible contribution of the alarmin/cytokine IL-33. *Front Immunol*. (2018) 9:2308. doi: 10.3389/fimmu.2018.02308
 45. Thierry A, Giraud S, Robin A, Barra A, Bridoux F, Ametieu V, et al. The alarmin concept applied to human renal transplantation: evidence for a differential implication of HMGB1 and IL-33. *PLoS ONE*. (2014) 9:e88742. doi: 10.1371/journal.pone.0088742
 46. Sabapathy V, Stremeska ME, Mohammad S, Corey RL, Sharma PR, Sharma R. Novel immunomodulatory cytokine regulates inflammation, diabetes, and obesity to protect from diabetic nephropathy. *Front Pharmacol*. (2019) 10:572. doi: 10.3389/fphar.2019.00572
 47. Mahlaköiv T, Flamar A-L, Johnston LK, Moriyama S, Putzel GG, Bryce PJ, et al. Stromal cells maintain immune cell homeostasis in adipose tissue via production of interleukin-33. *Sci Immunol*. (2019) 4:eaa0416. doi: 10.1126/sciimmunol.aax0416
 48. Spallanzani RG, Zemmour D, Xiao T, Jayewickreme T, Li C, Bryce PJ, et al. Distinct immunocyte-promoting and adipocyte-generating stromal components coordinate adipose tissue immune and metabolic tenors. *Sci Immunol*. (2019) 4:eaaw3658. doi: 10.1126/sciimmunol.aaw3658
 49. Green JA, Arpaia N, Schizas M, Dobrin A, Rudensky AY. A nonimmune function of T cells in promoting lung tumor progression. *J Exp Med*. (2017) 214:3565–75. doi: 10.1084/jem.20170356
 50. Christ A, Günther P, Lauterbach MAR, Duestel P, Biswas D, Pelka K, et al. Western diet triggers NLRP3-dependent innate immune reprogramming. *Cell*. (2018) 172:162–75.e14. doi: 10.1016/j.cell.2017.12.013
 51. Kaufmann E, Sanz J, Dunn JL, Khan N, Mendonça LE, Pacis A, et al. BCG educates hematopoietic stem cells to generate protective innate immunity against tuberculosis. *Cell*. (2018) 172:176–90.e19. doi: 10.1016/j.cell.2017.12.031
 52. Arshad MI, Rauch M, L'Helgoualc'h A, Julia V, Leite-de-Moraes MC, Lucas-Clerc C, et al. NKT cells are required to induce high IL-33 expression in hepatocytes during ConA-induced acute hepatitis. *Eur J Immunol*. (2011) 41:2341–8. doi: 10.1002/eji.201041332
 53. Matsukawa A, Fukumoto T, Maeda T, Ohkawara S, Yoshinaga M. Detection and characterization of IL-1 receptor antagonist in tissues from healthy rabbits: IL-1 receptor antagonist is probably involved in health. *Cytokine*. (1997) 9:307–15. doi: 10.1006/cyto.1996.0170
 54. Dinarello CA. Immunological and inflammatory functions of the interleukin-1 family. *Annu Rev Immunol*. (2009) 27:519–50. doi: 10.1146/annurev.immunol.021908.132612
 55. Sakai N, Van Sweringen HL, Quillin RC, Schuster R, Blanchard J, Burns JM, et al. Interleukin-33 is hepatoprotective during liver ischemia/reperfusion in mice. *Hepatology*. (2012) 56:1468–78. doi: 10.1002/hep.25768
 56. Bernal W, Wendon J. Acute liver failure. *N Engl J Med*. (2013) 369:2525–34. doi: 10.1056/NEJMra1208937
 57. Gotthardt D, Riediger C, Weiss KH, Encke J, Schemmer P, Schmidt J, et al. Fulminant hepatic failure: etiology and indications for liver transplantation. *Nephrol Dial Transplant*. (2007) 22(Suppl. 8):viii5–8. doi: 10.1093/ndt/gfm650
 58. Sakurai T, He G, Matsuzawa A, Yu G, Maeda S, Hardiman G, et al. Hepatocyte necrosis induced by oxidative stress and IL-1 α release mediate carcinogen-induced compensatory proliferation and liver

- tumorigenesis. *Cancer Cell*. (2008) 14:156–65. doi: 10.1016/j.ccr.2008.06.016
59. Chen C-J, Kono H, Golenbock D, Reed G, Akira S, Rock KL. Identification of a key pathway required for the sterile inflammatory response triggered by dying cells. *Nat Med*. (2007) 13:851–6. doi: 10.1038/nm1603
 60. Sultan M, Ben-Ari Z, Masoud R, Pappo O, Harats D, Kamari Y, et al. Interleukin-1 α and interleukin-1 β play a central role in the pathogenesis of fulminant hepatic failure in mice. *PLoS ONE*. (2017) 12:e0184084. doi: 10.1371/journal.pone.0184084
 61. Gabay C, Smith MF, Eidlén D, Arend WP. Interleukin 1 receptor antagonist (IL-1Ra) is an acute-phase protein. *J Clin Invest*. (1997) 99:2930–40. doi: 10.1172/JCI119488
 62. Tsutsui H, Kayagaki N, Kuida K, Nakano H, Hayashi N, Takeda K, et al. Caspase-1-independent, Fas/Fas ligand-mediated IL-18 secretion from macrophages causes acute liver injury in mice. *Immunity*. (1999) 11:359–67. doi: 10.1016/S1074-7613(00)80111-9
 63. Yan C, Zhou L, Han Y-P. Contribution of hepatic stellate cells and matrix metalloproteinase 9 in acute liver failure. *Liver Int*. (2008) 28:959–71. doi: 10.1111/j.1478-3231.2008.01775.x
 64. Shinoda M, Tilles AW, Kobayashi N, Wakabayashi G, Takayanagi A, Totsugawa T, et al. A bioartificial liver device secreting interleukin-1 receptor antagonist for the treatment of hepatic failure in rats. *J Surg Res*. (2007) 137:130–40. doi: 10.1016/j.jss.2006.08.009
 65. Shakoory B, Carcillo JA, Chatham WW, Amdur RL, Zhao H, Dinarello CA, et al. Interleukin-1 receptor blockade is associated with reduced mortality in sepsis patients with features of macrophage activation syndrome: reanalysis of a prior phase III trial*. *Crit Care Med*. (2016) 44:275–81. doi: 10.1097/CCM.0000000000001402
 66. Tsutsui H, Matsui K, Kawada N, Hyodo Y, Hayashi N, Okamura H, et al. IL-18 accounts for both TNF- α - and Fas ligand-mediated hepatotoxic pathways in endotoxin-induced liver injury in mice. *J Immunol*. (1997) 159:3961–7.
 67. Finotto S, Siebler J, Hausding M, Schipp M, Wirtz S, Klein S, et al. Severe hepatic injury in interleukin 18 (IL-18) transgenic mice: a key role for IL-18 in regulating hepatocyte apoptosis *in vivo*. *Gut*. (2004) 53:392–400. doi: 10.1136/gut.2003.018572
 68. Imaeda AB, Watanabe A, Sohail MA, Mahmood S, Mohammadnejad M, Sutterwala FS, et al. Acetaminophen-induced hepatotoxicity in mice is dependent on Tlr9 and the Nalp3 inflammasome. *J Clin Invest*. (2009) 119:305–14. doi: 10.1172/JCI35958
 69. Antunes MM, Araújo AM, Diniz AB, Pereira RVS, Alvarenga DM, David BA, et al. IL-33 signalling in liver immune cells enhances drug-induced liver injury and inflammation. *Inflamm Res*. (2018) 67:77–88. doi: 10.1007/s00011-017-1098-3
 70. Scheiermann P, Bachmann M, Hårdle L, Pleli T, Piiper A, Zwissler B, et al. Application of IL-36 receptor antagonist weakens CCL20 expression and impairs recovery in the late phase of murine acetaminophen-induced liver injury. *Sci Rep*. (2015) 5:1–7. doi: 10.1038/srep08521
 71. Arshad MI, Patrat-Delon S, Piquet-Pellorce C, L'Helgoualc'h A, Rauch M, Genet V, et al. Pathogenic mouse hepatitis virus or poly(I:C) induce IL-33 in hepatocytes in murine models of hepatitis. *PLoS ONE*. (2013) 8:1–10. doi: 10.1371/journal.pone.0074278
 72. Kamo N, Ke B, Ghaffari AA, Shen X, Busuttill RW, Cheng G, et al. ASC/caspase-1/IL-1 β signaling triggers inflammatory responses by promoting HMGB1 induction in liver ischemia/reperfusion injury. *Hepatology*. (2013) 58:351–62. doi: 10.1002/hep.26320
 73. Kim HY, Kim SJ, Lee SM. Activation of NLRP3 and AIM2 inflammasomes in Kupffer cells in hepatic ischemia/reperfusion. *FEBS J*. (2015) 282:259–70. doi: 10.1111/febs.13123
 74. Inoue Y, Shirasuna K, Kimura H, Usui F, Kawashima A, Karasawa T, et al. NLRP3 regulates neutrophil functions and contributes to hepatic ischemia-reperfusion injury independently of inflammasomes. *J Immunol*. (2014) 192:4342–51. doi: 10.4049/jimmunol.1302039
 75. Sadatomo A, Inoue Y, Ito H, Karasawa T, Kimura H, Watanabe S, et al. Interaction of neutrophils with macrophages promotes IL-1 β maturation and contributes to hepatic ischemia-reperfusion injury. *J Immunol*. (2017) 199:3306–15. doi: 10.4049/jimmunol.1700717
 76. Yazdani HO, Chen H-W, Tohme S, Tai S, van der Windt DJ, Loughran P, et al. IL-33 exacerbates liver sterile inflammation by amplifying neutrophil extracellular trap formation. *J Hepatol*. (2017) 68:130–9. doi: 10.1016/j.jhep.2017.09.010
 77. Arrenberg P, Maricic I, Kumar V. Sulfatide-mediated activation of type II natural killer T cells prevents hepatic ischemic reperfusion injury in mice. *Gastroenterology*. (2011) 140:646–55. doi: 10.1053/j.gastro.2010.10.003
 78. Sakai N, Van Sweringen HL, Belzaira RM, Quillin RC, Schuster R, Blanchard J, et al. Interleukin-37 reduces liver inflammatory injury via effects on hepatocytes and non-parenchymal cells. *J Gastroenterol Hepatol*. (2012) 27:1609–16. doi: 10.1111/j.1440-1746.2012.07187.x
 79. Muto Y, Nouri-Aria KT, Meager A, Alexander GJ, Eddleston AL, Williams R. Enhanced tumour necrosis factor and interleukin-1 in fulminant hepatic failure. *Lancet*. (1988) 2:72–4. doi: 10.1016/S0140-6736(88)90006-2
 80. Sekiyama KD, Yoshida M, Thomson AW. Circulating proinflammatory cytokines (IL-1 β , TNF- α , and IL-6) and IL-1 receptor antagonist (IL-1Ra) in fulminant hepatic failure and acute hepatitis. *Clin Exp Immunol*. (1994) 98:71–7. doi: 10.1111/j.1365-2249.1994.tb06609.x
 81. Roth GA, Faybik P, Hetz H, Hacker S, Ankersmit HJ, Bacher A, et al. Pro-inflammatory interleukin-18 and Caspase-1 serum levels in liver failure are unaffected by MARS treatment. *Dig Liver Dis*. (2009) 41:417–23. doi: 10.1016/j.dld.2008.09.021
 82. Roth GA, Zimmermann M, Lubczyk BA, Pilz J, Faybik P, Hetz H, et al. Up-regulation of interleukin 33 and soluble ST2 serum levels in liver failure. *J Surg Res*. (2010) 163:e79–83. doi: 10.1016/j.jss.2010.04.004
 83. Kim J, Chang D-Y, Lee HW, Lee H, Kim JH, Sung PS, et al. Innate-like cytotoxic function of bystander-activated CD8+ T cells is associated with liver injury in acute hepatitis A. *Immunity*. (2018) 48:161–73.e5. doi: 10.1016/j.immuni.2017.11.025
 84. Iracheta-Vellve A, Petrasek J, Satishchandran A, Gyongyosi B, Saha B, Kodys K, et al. Inhibition of sterile danger signals, uric acid and ATP, prevents inflammasome activation and protects from alcoholic steatohepatitis in mice. *J Hepatol*. (2015) 63:1147–55. doi: 10.1016/j.jhep.2015.06.013
 85. Petrasek J, Bala S, Csak T, Lippai D, Kodys K, Menashy V, et al. IL-1 receptor antagonist ameliorates inflammasome-dependent alcoholic steatohepatitis in mice. *J Clin Invest*. (2012) 122:3476–89. doi: 10.1172/JCI60777
 86. Cui K, Yan G, Xu C, Chen Y, Wang J, Zhou R, et al. Invariant NKT cells promote alcohol-induced steatohepatitis through interleukin-1 β in mice. *J Hepatol*. (2015) 62:1311–8. doi: 10.1016/j.jhep.2014.12.027
 87. Heo MJ, Kim TH, You JS, Blaya D, Sancho-Bru P, Kim SG. Alcohol dysregulates miR-148a in hepatocytes through FoxO1, facilitating pyroptosis via TXNIP overexpression. *Gut*. (2019) 68:708–20. doi: 10.1136/gutjnl-2017-315123
 88. Sun Z, Chang B, Huang A, Hao S, Gao M, Sun Y, et al. Plasma levels of soluble ST2, but not IL-33, correlate with the severity of alcoholic liver disease. *J Cell Mol Med*. (2018) 23:887–97. doi: 10.1111/jcmm.13990
 89. Wang M, Shen G, Xu L, Liu X, Brown JM, Feng D, et al. IL-1 receptor like 1 protects against alcoholic liver injury by limiting NF- κ B activation in hepatic macrophages. *J Hepatol*. (2018) 68:109–17. doi: 10.1016/j.jhep.2017.08.023
 90. Maricic I, Sheng H, Marrero I, Seki E, Kisseleva T, Chaturvedi S, et al. Inhibition of type I natural killer T cells by retinoids or following sulfatide-mediated activation of type II natural killer T cells attenuates alcoholic liver disease in mice. *Hepatology*. (2015) 61:1357–69. doi: 10.1002/hep.27632
 91. Mathews S, Feng D, Maricic I, Ju C, Kumar V, Gao B. Invariant natural killer T cells contribute to chronic-plus-binge ethanol-mediated liver injury by promoting hepatic neutrophil infiltration. *Cell Mol Immunol*. (2016) 13:206–16. doi: 10.1038/cmi.2015.06
 92. Bugianesi E, Rosso C, Cortez-Pinto H. How to diagnose NAFLD in 2016. *J Hepatol*. (2016) 65:643–4. doi: 10.1016/j.jhep.2016.05.038
 93. Bellentani S. The epidemiology of non-alcoholic fatty liver disease. *Liver Int*. (2017) 37:81–4. doi: 10.1111/liv.13299
 94. European Association for the Study of the Liver (EASL), European Association for the Study of Diabetes (EASD), European Association for the Study of Obesity (EASO). EASL–EASD–EASO Clinical Practice Guidelines for the management of non-alcoholic fatty liver disease. *Diabetologia*. (2016) 59:1121–40. doi: 10.1007/s00125-016-3902-y

95. Yu J, Marsh S, Hu J, Feng W, Wu C. The pathogenesis of nonalcoholic fatty liver disease: interplay between diet, gut microbiota, and genetic background. *Gastroenterol Res Pract.* (2016) 2016:2862173. doi: 10.1155/2016/2862173
96. Negrin KA, Flach RJR, DiStefano MT, Matevossian A, Friedline RH, Jung D, et al. IL-1 Signaling in obesity-induced hepatic lipogenesis and steatosis. *PLoS ONE.* (2014) 9:e107265. doi: 10.1371/journal.pone.0107265
97. Miura K, Yang L, van Rooijen N, Brenner DA, Ohnishi H, Seki E. Toll-like receptor 2 and palmitic acid cooperatively contribute to the development of nonalcoholic steatohepatitis through inflammasome activation in mice. *Hepatology.* (2013) 57:577–89. doi: 10.1002/hep.26081
98. Csak T, Ganz M, Pespisa J, Kodys K, Dolganiuc A, Szabo G. Fatty acid and endotoxin activate inflammasomes in mouse hepatocytes that release danger signals to stimulate immune cells. *Hepatology.* (2011) 54:133–44. doi: 10.1002/hep.24341
99. Moschen AR, Molnar C, Enrich B, Geiger S, Ebenbichler CF, Tilg H. Adipose and liver expression of interleukin (IL)-1 family members in morbid obesity and effects of weight loss. *Mol Med Camb Mass.* (2011) 17:840–5. doi: 10.2119/molmed.2010.00108
100. Patrick AL, Rullo J, Beaudin S, Liaw P, Fox-Robichaud AE. Hepatic leukocyte recruitment in response to time-limited expression of TNF- α and IL-1 β . *Am J Physiol-Gastrointest Liver Physiol.* (2007) 293:G663–72. doi: 10.1152/ajpgi.00070.2007
101. Tilg H, Moschen AR, Szabo G. Interleukin-1 and inflammasomes in alcoholic liver disease/acute alcoholic hepatitis and nonalcoholic fatty liver disease/nonalcoholic steatohepatitis. *Hepatology.* (2016) 64:955–65. doi: 10.1002/hep.28456
102. Kamari Y, Shaish A, Vax E, Shemesh S, Kandel-Kfir M, Arbel Y, et al. Lack of interleukin-1 α or interleukin-1 β inhibits transformation of steatosis to steatohepatitis and liver fibrosis in hypercholesterolemic mice. *J Hepatol.* (2011) 55:1086–94. doi: 10.1016/j.jhep.2011.01.048
103. Jager J, Grémeaux T, Cormont M, Le Marchand-Brustel Y, Tanti J-F. Interleukin-1 β -induced insulin resistance in adipocytes through down-regulation of insulin receptor substrate-1 expression. *Endocrinology.* (2007) 148:241–51. doi: 10.1210/en.2006-0692
104. Nov O, Kohl A, Lewis EC, Bashan N, Dvir I, Ben-Shlomo S, et al. Interleukin-1 β may mediate insulin resistance in liver-derived cells in response to adipocyte inflammation. *Endocrinology.* (2010) 151:4247–56. doi: 10.1210/en.2010-0340
105. Wen H, Gris D, Lei Y, Jha S, Zhang L, Huang MT-H, et al. Fatty acid-induced NLRP3-ASC inflammasome activation interferes with insulin signaling. *Nat Immunol.* (2011) 12:408–15. doi: 10.1038/ni.2022
106. Matsuki T, Horai R, Sudo K, Iwakura Y. IL-1 plays an important role in lipid metabolism by regulating insulin levels under physiological conditions. *J Exp Med.* (2003) 198:877–88. doi: 10.1084/jem.20030299
107. Isoda K, Sawada S, Ayaori M, Matsuki T, Horai R, Kagata Y, et al. Deficiency of interleukin-1 receptor antagonist deteriorates fatty liver and cholesterol metabolism in hypercholesterolemic mice. *J Biol Chem.* (2005) 280:7002–9. doi: 10.1074/jbc.M412220200
108. Henao-Mejia J, Elinav E, Jin C, Hao L, Mehal WZ, Strowig T, et al. Inflammasome-mediated dysbiosis regulates progression of NAFLD and obesity. *Nature.* (2012) 482:179–85. doi: 10.1038/nature10809
109. Vasseur P, Dion S, Filliol A, Genet V, Jean-philippe G, Silvain C, et al. Endogenous IL-33 has no effect on the progression of fibrosis during experimental steatohepatitis. *Oncotarget.* (2017) 8:48563–74. doi: 10.18632/oncotarget.18335
110. Gao Y, Liu Y, Yang M, Guo X, Zhang M, Li H, et al. IL-33 treatment attenuated diet-induced hepatic steatosis but aggravated hepatic fibrosis. *Oncotarget.* (2016) 7:1–13. doi: 10.18632/oncotarget.9259
111. Jetic I, Jovicic N, Pantic J, Arsenijevic N, Lukic ML, Pejinovic N. Galectin-3 ablation enhances liver steatosis, but attenuates inflammation and IL-33-dependent fibrosis in obesogenic mouse model of nonalcoholic steatohepatitis. *Mol Med Camb Mass.* (2015) 21:453–65. doi: 10.2119/molmed.2014.00178
112. Jaruga B, Hong F, Sun R, Radaeva S, Gao B. Crucial role of IL-4/STAT6 in T cell-mediated hepatitis: up-regulating eotaxins and IL-5 and recruiting leukocytes. *J Immunol.* (2003) 171:3233–44. doi: 10.4049/jimmunol.171.6.3233
113. Tagawa Y, Sekikawa K, Iwakura Y. Suppression of concanavalin A-induced hepatitis in IFN- γ (-/-) mice, but not in TNF- α (-/-) mice: role for IFN- γ in activating apoptosis of hepatocytes. *J Immunol.* (1997) 159:1418–28.
114. Zhu R, Diem S, Araujo LM, Aumeunier A, Denizeau J, Philadelphie E, et al. The Pro-Th1 cytokine IL-12 enhances IL-4 production by invariant NKT cells: relevance for T cell-mediated hepatitis. *J Immunol.* (2007) 178:5435–42. doi: 10.4049/jimmunol.178.9.5435
115. Takeda K, Hayakawa Y, Van Kaer L, Matsuda H, Yagita H, Okumura K. Critical contribution of liver natural killer T cells to a murine model of hepatitis. *Proc Natl Acad Sci USA.* (2000) 97:5498–503. doi: 10.1073/pnas.040566697
116. Arshad MI, Piquet-Pellorce C, L'Helgoual'h A, Rauch M, Patrat-Delon S, Ezan F, et al. TRAIL but not FasL and TNF α , regulates IL-33 expression in murine hepatocytes during acute hepatitis. *Hepatology.* (2012) 56:2353–62. doi: 10.1002/hep.25893
117. Schümann J, Wolf D, Pahl A, Brune K, Papadopoulos T, van Rooijen N, et al. Importance of Kupffer cells for T-cell-dependent liver injury in mice. *Am J Pathol.* (2000) 157:1671–83. doi: 10.1016/S0002-9440(10)64804-3
118. Luan J, Zhang X, Wang S, Li Y, Fan J, Chen W, et al. NOD-like receptor protein 3 inflammasome-dependent IL-1 β accelerated ConA-induced Hepatitis. *Front Immunol.* (2018) 9:758. doi: 10.3389/fimmu.2018.00758
119. Noel G, Arshad MI, Filliol A, Genet V, Rauch M, Lucas-Clerc C, et al. Ablation of interaction between IL-33 and ST2+ regulatory T cells increases immune cell-mediated hepatitis and activated NK cell liver infiltration. *Am J Physiol Gastrointest Liver Physiol.* (2016) 311:G313–23. doi: 10.1152/ajpgi.00097.2016
120. Volarevic V, Mitrovic M, Milovanovic M, Zelen I, Nikolic I, Mitrovic S, et al. Protective role of IL-33/ST2 axis in Con A-induced hepatitis. *J Hepatol.* (2012) 56:26–33. doi: 10.1016/j.jhep.2011.03.022
121. Burzyn D, Kuswanto W, Kolodin D, Shadrach JL, Cerletti M, Jang Y, et al. A special population of regulatory T cells potentiates muscle repair. *Cell.* (2013) 155:1282–95. doi: 10.1016/j.cell.2013.10.054
122. Shrivastava S, Mukherjee A, Ray R, Ray RB. Hepatitis C virus induces interleukin-1 (IL-1)/IL-18 in circulatory and resident liver macrophages. *J Virol.* (2013) 87:12284–90. doi: 10.1128/JVI.01962-13
123. Negash AA, Ramos HJ, Crochet N, Lau DTY, Doehle B, Papp N, et al. IL-1 β production through the NLRP3 inflammasome by hepatic macrophages links hepatitis C virus infection with liver inflammation and disease. *PLoS Pathog.* (2013) 9:1–13. doi: 10.1371/journal.ppat.1003330
124. Yanagisawa K, Yue S, van der Vliet HJ, Wang R, Alatrakchi N, Golden-Mason L, et al. *Ex vivo* analysis of resident hepatic pro-inflammatory CD1d-reactive T cells and hepatocyte surface CD1d expression in hepatitis C. *J Viral Hepat.* (2013) 20:556–65. doi: 10.1111/jvh.12081
125. Fasicaro P, Valdatta C, Massari M, Loggi E, Biasini E, Sacchelli L, et al. Antiviral intrahepatic T-cell responses can be restored by blocking programmed death-1 pathway in chronic hepatitis B. *Gastroenterology.* (2010) 138:682–93, 693.e1–4. doi: 10.1053/j.gastro.2009.09.052
126. Hofmann M, Thimme R. iNKT cells in chronic HBV: a balancing act. *Hepatology Int.* (2016) 10:535–7. doi: 10.1007/s12072-016-9739-0
127. Li M, Zhou Z-H, Sun X-H, Zhang X, Zhu X-J, Jin S-G, et al. The dynamic changes of circulating invariant natural killer T cells during chronic hepatitis B virus infection. *Hepatology Int.* (2016) 10:594–601. doi: 10.1007/s12072-015-9650-0
128. Kimura K, Kakimi K, Wieland S, Guidotti LG, Chisari FV. Interleukin-18 inhibits hepatitis B virus replication in the livers of transgenic mice. *J Virol.* (2002) 76:10702–7. doi: 10.1128/JVI.76.21.10702-10707.2002
129. Du XX, Shi Y, Yang Y, Yu Y, Lou HG, Lv FF, et al. DAMP molecular IL-33 augments monocyte inflammatory storm in hepatitis B-precipitated acute-on-chronic liver failure. *Liver Int.* (2018) 38:229–38. doi: 10.1111/liv.13503
130. Huan S-L, Zhao J-G, Wang Z-L, Gao S, Wang K. Relevance of serum interleukin-33 and ST2 levels and the natural course of chronic hepatitis B virus infection. *BMC Infect Dis.* (2016) 16:200. doi: 10.1186/s12879-016-1543-x
131. Wang J, Zhao P, Guo H, Sun X, Jiang Z, Xu L, et al. Serum IL-33 levels are associated with liver damage in patients with chronic hepatitis C. *Mediators Inflamm.* (2012) 2012:819636. doi: 10.1155/2012/819636

132. Gieling RG, Wallace K, Han Y-P. Interleukin-1 participates in the progression from liver injury to fibrosis. *Am J Physiol-Gastrointest Liver Physiol.* (2009) 296:G1324–31. doi: 10.1152/ajpgi.90564.2008
133. Marvie P, Lisbonne M, L'helgoualc'h A, Rauch M, Turlin B, Preisser L, et al. Interleukin-33 overexpression is associated with liver fibrosis in mice and humans. *J Cell Mol Med.* (2010) 14:1726–39. doi: 10.1111/j.1582-4934.2009.00801.x
134. Mchedlidze T, Waldner M, Zopf S, Walker J, Rankin AL, Schuchmann M, et al. Interleukin-33-dependent innate lymphoid cells mediate hepatic fibrosis. *Immunity.* (2013) 39:357–71. doi: 10.1016/j.immuni.2013.07.018
135. Ma C, Han M, Heinrich B, Fu Q, Zhang Q, Sandhu M, et al. Gut microbiome-mediated bile acid metabolism regulates liver cancer via NKT cells. *Science.* (2018) 360:eaan5931. doi: 10.1126/science.aan5931
136. Miyagi T, Takehara T, Tatsumi T, Kanto T, Suzuki T, Jinushi M, et al. CD1d-mediated stimulation of natural killer T cells selectively activates hepatic natural killer cells to eliminate experimentally disseminated hepatoma cells in murine liver. *Int J Cancer.* (2003) 106:81–9. doi: 10.1002/ijc.11163
137. Leite-De-Moraes MC, Hameg A, Arnould A, Machavoine F, Koezuka Y, Schneider E, et al. A distinct IL-18-induced pathway to fully activate NK T lymphocytes independently from TCR engagement. *J Immunol.* (1999) 163:5871–6.
138. Bricard G, Cesson V, Devevre E, Bouzourene H, Barbey C, Rufer N, et al. Enrichment of human CD4⁺ V α 24/V β 11 invariant NKT cells in intrahepatic malignant tumors. *J Immunol.* (2009) 182:5140–51. doi: 10.4049/jimmunol.0711086
139. Gao X, Wang X, Yang Q, Zhao X, Wen W, Li G, et al. Tumoral expression of IL-33 inhibits tumor growth and modifies the tumor microenvironment through CD8⁺ T and NK cells. *J Immunol.* (2015) 194:438–45. doi: 10.4049/jimmunol.1401344
140. Hollande C, Boussier J, Ziai J, Nozawa T, Bondet V, Phung W, et al. Inhibition of the dipeptidyl peptidase DPP4 (CD26) reveals IL-33-dependent eosinophil-mediated control of tumor growth. *Nat Immunol.* (2019) 20:257–64. doi: 10.1038/s41590-019-0321-5
141. Villarreal DO, Wise MC, Walters JN, Reuschel EL, Choi MJ, Obeng-Adjei N, et al. Alarmin IL-33 acts as an immunoadjuvant to enhance antigen-specific tumor immunity. *Cancer Res.* (2014) 74:1789–800. doi: 10.1158/0008-5472.CAN-13-2729
142. Zhang J, Zhang Q, Lou Y, Fu Q, Chen Q, Wei T, et al. Hypoxia-inducible factor-1 α /interleukin-1 β signaling enhances hepatoma epithelial-mesenchymal transition through macrophages in a hypoxic-inflammatory microenvironment. *Hepatology.* (2018). 67:1872–89. doi: 10.1002/hep.29681
143. Zhao J-J, Pan Q-Z, Pan K, Weng D-S, Wang Q-J, Li J-J, et al. Interleukin-37 mediates the antitumor activity in hepatocellular carcinoma: role for CD57+ NK cells. *Sci Rep.* (2015) 4:5177. doi: 10.1038/srep05177
144. Liu R, Tang C, Shen A, Luo H, Wei X, Zheng D, et al. IL-37 suppresses hepatocellular carcinoma growth by converting pSmad3 signaling from JNK/pSmad3L/c-Myc oncogenic signaling to pSmad3C/P21 tumor-suppressive signaling. *Oncotarget.* (2016) 7:85079–96. doi: 10.18632/oncotarget.13196
145. Li T-T, Zhu D, Mou T, Guo Z, Pu J-L, Chen Q-S, et al. IL-37 induces autophagy in hepatocellular carcinoma cells by inhibiting the PI3K/AKT/mTOR pathway. *Mol Immunol.* (2017) 87:132–40. doi: 10.1016/j.molimm.2017.04.010
146. Yang Y, Wang JB, Li YM, Zhao Y, Wang R, Wu Q, et al. Role of IL-33 expression in oncogenesis and development of human hepatocellular carcinoma. *Oncol Lett.* (2016) 12:429–36. doi: 10.3892/ol.2016.4622
147. Zhang P, Liu X-K, Chu Z, Ye J, Li K-L, Zhuang W-L, et al. Detection of interleukin-33 in serum and carcinoma tissue from patients with hepatocellular carcinoma and its clinical implications. *J Int Med Res.* (2012) 40:1654–61. doi: 10.1177/030006051204000504
148. Bergis D, Kassis V, Ranglack A, Koeberle V, Piiper A, Kronenberger B, et al. High serum levels of the interleukin-33 receptor soluble ST2 as a negative prognostic factor in hepatocellular carcinoma. *Transl Oncol.* (2013) 6:311–8. doi: 10.1593/tlo.12418
149. Berasain C, Castillo J, Perugorria MJ, Latasa MU, Prieto J, Avila MA. Inflammation and liver cancer: new molecular links. *Ann N Y Acad Sci.* (2009) 1155:206–21. doi: 10.1111/j.1749-6632.2009.03704.x
150. Wu X, Dong L, Lin X, Li J. Relevance of the NLRP3 inflammasome in the pathogenesis of chronic liver disease. *Front Immunol.* (2017) 8:1728. doi: 10.3389/fimmu.2017.01728
151. Yang Y, Wang H, Kouadir M, Song H, Shi F. Recent advances in the mechanisms of NLRP3 inflammasome activation and its inhibitors. *Cell Death Dis.* (2019) 10:128. doi: 10.1038/s41419-019-1413-8
152. Adkison AM, Raptis SZ, Kelley DG, Pham CTN. Dipeptidyl peptidase I activates neutrophil-derived serine proteases and regulates the development of acute experimental arthritis. *J Clin Invest.* (2002) 109:363–71. doi: 10.1172/JCI0213462
153. Lust JA, Lacy MQ, Zeldenrust SR, Witzig TE, Moon-Tasson LL, Dinarello CA, et al. Reduction in C-reactive protein indicates successful targeting of the IL-1/IL-6 axis resulting in improved survival in early stage multiple myeloma. *Am J Hematol.* (2016) 91:571–4. doi: 10.1002/ajh.24352
154. Wu T-C, Xu K, Martinek J, Young RR, Banchereau R, George J, et al. IL1 receptor antagonist controls transcriptional signature of inflammation in patients with metastatic breast cancer. *Cancer Res.* (2018) 78:5243–58. doi: 10.1158/0008-5472.CAN-18-0413
155. Ridker PM, Everett BM, Thuren T, MacFadyen JG, Chang WH, Ballantyne C, et al. Antiinflammatory therapy with canakinumab for atherosclerotic disease. *N Engl J Med.* (2017) 377:1119–31. doi: 10.1056/NEJMoa1707914
156. Stremaska ME, Jose S, Sabapathy V, Huang L, Bajwa A, Kinsey GR, et al. IL233, A novel IL-2 and IL-33 hybrid cytokine, ameliorates renal injury. *J Am Soc Nephrol.* (2017) 28:2681–93. doi: 10.1681/ASN.2016121272

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Barbier, Ferhat, Salamé, Robin, Herbelin, Gombert, Silvain and Barbarin. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.