



HAL
open science

Adapting palliative radiation therapy for bone metastases during the Covid-19 pandemic: GEMO position paper

Sébastien Thureau, Jean Faivre, Richard Assaker, Emmanuel Biver, Cyrille Confavreux, Françoise Debais, Martine Duterque-Coquillaud, Fransesco Giammarile, Dominique Heymann, Frédéric Lecouvet, et al.

► To cite this version:

Sébastien Thureau, Jean Faivre, Richard Assaker, Emmanuel Biver, Cyrille Confavreux, et al.. Adapting palliative radiation therapy for bone metastases during the Covid-19 pandemic: GEMO position paper. Journal of Bone Oncology, Amsterdam: Elsevier, c2012-, 2020, 22, pp.100291. 10.1016/j.jbo.2020.100291 . inserm-02529599v2

HAL Id: inserm-02529599

<https://www.hal.inserm.fr/inserm-02529599v2>


Submitted on 23 Aug 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives | 4.0 International License



Adapting palliative radiation therapy for bone metastases during the Covid-19 pandemic: GEMO position paper

Sébastien Thureau^{a,*}, Jean Christophe Faivre^b, Richard Assaker^c, Emmanuel Biver^d,
Cyrille B. Confavreux^e, Françoise Debiais^f, Martine Duterque-Coquillaud^g,
Francesco Giammarile^h, Dominique Heymannⁱ, Frédéric E. Lecouvet^j, Laetitia Morardet^k,
Frederic Paycha^l, Jean-Jacques Body^m, Marie-Hélène Vieillardⁿ

^a Radiation department, Center Henri Becquerel, Quantif-Litis EA 4108, University of Rouen, Rouen, France.

^b Radiation department, Institut de Cancérologie de Lorraine - Alexis-Vautrin, Vandœuvre-lès-Nancy, France.

^c University Department of neurosurgery, Lille University Hospital, Lille, France.

^d Division of Bone Diseases, Geneva University Hospitals and Faculty of Medicine, University of Geneva, Geneva, Switzerland.

^e INSERM UMR 1033 – University of Lyon, Bone Metastasis Expert Center (CEMOS) Cancer Institute of Hospices Civils de Lyon, Rheumatology Department, Centre Hospitalier Lyon Sud, Pierre Bénite, F- 69310 France.

^f Rheumatology department, CHU Poitiers, Poitiers, France

^g CNRS UMR9020, INSERM UMR1277, University of Lille, Institut Pasteur de Lille, CHU Lille, France.

^h Nuclear Medicine LUMEN, Centre Leon Berard, Lyon, France

ⁱ Institut de Cancérologie de l'Ouest, University of Nantes, Inserm, CRCINA, Saint-Herblain, France.

^j Institut du Cancer Roi Albert 2, Cliniques Universitaires Saint-Luc, Service de Radiologie et d'Imagerie Médicale, Institut de Recherche Expérimentale et Clinique (IREC), Université Catholique de Louvain (UCLouvain), Brussels, Belgium.

^k Oncology Department, Pitié Salpêtrière Hospital, Paris, France

^l Nuclear Medicine Department, Lariboisière University Hospital, Assistance Publique-Hôpitaux de Paris, Paris, France.

^m Department of Medicine, CHU Brugmann, Université Libre de Bruxelles (ULB), Brussels, Belgium.

ⁿ University department of Rheumatology, Lille University hospital; Oscar Lambret Center; CNRS UMR9020, INSERM UMR1277, University of Lille, Institut Pasteur, Lille.

A B S T R A C T

The current health crisis caused by COVID-19 is a challenge for oncology treatment, especially when it comes to radiotherapy. Cancer patients are already known to be very fragile and COVID-19 brings about the risk of severe respiratory complications. In order to treat patients safely while protecting medical teams, the entire health care system must optimize the way it approaches prevention and treatment at a time when social distancing is key to stemming this pandemic. All indications and treatment modalities must be re-discussed. This is particularly the case for radiotherapy of bone metastases for which it is possible to reduce the number of sessions, the frequency of transport and the complexity of treatments. These changes will have to be discussed according to the organization of each radiotherapy department and the health situation, while medical teams must remain vigilant about the risks of complications of bone metastases, particularly spinal metastases. In this short piece, the members of the GEMO (the European Study Group of Bone Metastases) offer a number of recommendations to achieve the above objectives, both in general and in relation to five of the most common situations on radiation therapy for bone metastases.

The current health crisis caused by COVID-19 is making it difficult to organize oncology departments, particularly radiotherapy departments [1]. While it continues to be paramount to avoid the transmission of COVID-19 and to ensure the safety of health and medical professionals, it is equally critical to allow essential treatments to continue in order to maximise palliation and quality of life of patients with advanced cancers. We can already learn from the Chinese and Italian experience with regard to the risk to cancer patients and the organization of radiotherapy departments [2,3].

GEMO is the European Multidisciplinary Francophone Society for the study of bone metastases. It consists of scientists and physicians on the frontline of the prevention, treatment, and research on bone metastases. And it is in this framework and on the back of our collective experience that this group sets out recommendations for palliative radiotherapy to bone metastases during this crisis period of the COVID-19 pandemic.

As this pandemic expands, the management of patients with bone metastases, and the organization of radiotherapy departments, may not

* Corresponding author: Academic Department of Radiation Therapy & Medical Physics, Centre Henri Becquerel [Cancer Center] – Unicancer, Rue d'Amiens, Rouen cedex, F-76038, France.

E-mail address: sebastien.thureau@chb.unicancer.fr (S. Thureau).

<https://doi.org/10.1016/j.jbo.2020.100291>

Received 31 March 2020; Received in revised form 1 April 2020; Accepted 2 April 2020

Available online 13 April 2020

2212-1374/ © 2020 Published by Elsevier GmbH. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

immediately appear to be as a priority. Yet, it is an important activity of our departments that can dramatically reduce pain and improve the quality of life of our patients while reducing the need for oral or intravenous analgesic treatments and their associated side-effects

In this time of crisis we must bear in mind that it is possible to adapt our approach to minimize the risk incurred by patients while optimizing their care. These recommendations are made in the context of the COVID-19 epidemic in France and are likely to evolve depending on the health situation.

During this pandemic we must first and foremost be more vigilant than ever to the symptoms reported by patients with bone metastases. This is particularly the case for spinal cord compression (SCC), which requires rapid diagnosis and treatment in order to minimise neurologic complications.

Overall, the treatment of bone metastases by radiotherapy is based on existing, wide-ranging and well-established evidence-based literature and recommendations [4,5].

As a starting point in this context, we would like to make several general recommendations:

- Firstly, and whenever possible, medical teams should carry out the patient's CT scan planning and treatment at the same time as the consultation in order to limit transport and close contact with other potential COVID-19 patients.
- Secondly, always consider carrying out teleconsultations if possible and appropriate.
- Thirdly, there are well-known criteria to assess bone neoplastic instability, and proper stabilization techniques must be used. In this context, try to opt for the least invasive techniques.

In addition to the above, here are recommendations in relation to five of the most common situations on radiation therapy for bone metastases:

1. Painful bone metastasis

This metastatic population is extremely frail and should be kept away from the risk of COVID-19 infection. It is essential to adapt the medical treatment as much as possible and to avoid palliative radiotherapy in patients controlled by level 1 to 3 oral analgesics.

- Statement 1: Palliative radiotherapy remains an important option for patients experiencing significant pain, diminished quality of life and reduced autonomy as a result of bone metastases. This is especially so if it enables a reduction in the need for daily nursing care (patients at home with intravenous pain medication), in opioid-resistant pain or in case of poor tolerance of opioids (strong opioids).
- Statement 2: We recommend the use of monofractionated radiotherapy (8Gy in 1 fraction). Fractionated treatment should be avoided (eg., 30Gy / 10 fractions, 20Gy / 4 or 5 fractions).
- Statement 3: The simplest conformal radiation therapy techniques should be used.

2. Retreatment of painful bone metastasis

- Statement 1: We recommend waiting a minimum of 6 weeks after completion of the initial radiotherapy (possibility of delayed analgesic response) before following the same procedure as for the initial treatment.
- Statement 2: One fraction of 8 Gy is also the reference schedule for retreatment [6].
- Statement 3: The simplest conformal radiation therapy techniques should be used.

3. Metastatic Epidural Spinal Cord Compression (MESCC)

Surgical treatment should theoretically be preferred whenever possible and for all patients with a life expectancy of more than a few months. In clinical practice, the dramatic reduction in places available in recovery units because of the COVID-19 crisis means that surgery should be restricted to interventions not requiring resuscitation stays. This is in part linked to the high risk of COVID-19 contamination during inpatient stay.

- Statement 1: Adjuvant radiotherapy after surgery for MESCC can be postponed for 4 to 12 weeks.
- Statement 2: In cases where surgical treatment is contraindicated or not appropriate, radiotherapy should be arranged without delay [7,8].
- Statement 3: For exclusive radiotherapy, systematically perform monofractionated radiotherapy (8Gy in 1 fraction) [9].
- Statement 4: The simplest conformal radiation therapy techniques should be used.
- Statement 5: MESCC is likely the only instance justifying urgent management of a COVID+ patient.
- Statement 6: There is a limited evidence in the literature of the efficacy of corticosteroid therapy on spinal cord compression, particularly during radiotherapy. If there is no suspicion of COVID-19 and if there is a significant inflammatory risk during radiotherapy, a short course of corticosteroid therapy (16mg dexamethasone per os daily) may be prescribed [10,11].

4. Adjuvant bone metastasis radiotherapy

The level of evidence is insufficient to justify systematic adjuvant radiotherapy in either the spine or peripheral bones.

- Statement 1: radiotherapy may be postponed or performed secondarily in case of progressive post-operative signs [12].
- Statement 2: Fractionated treatment is recommended (30Gy / 10 fractions, 20Gy / 4 or 5 fractions).

5. Bone oligometastases and other SBRT (Stereotactic Body Radiation Therapy) indications

Stereotactic radiotherapy has some limited use for bone metastases (irradiation of radioresistant lesions, oligometastatic patients or radiotherapy in the irradiated area). However it is a more complex technique that requires additional examinations (spinal MRI in particular) as well as additional medical and physicist time for planning.

- Statement 1: The level of evidence for using SBRT in oligometastatic disease is too low for this treatment option to be considered in the current situation. Moreover, it is often possible to postpone this treatment for a few weeks, especially for hormone sensitive tumours.
- Statement 2: Symptomatic SCC in an irradiated area can be viewed as an indication for SBRT if conventional irradiation is not an option. Single fraction treatment should be preferred (16 to 24Gy), although there is a slightly higher fracture risk than with fractionated SBRT (the alternative is a fractionated treatment in 3 to 5 fractions). It is essential to take into account the initial radiotherapy and in particular the dose to the spinal cord (Equivalent dose in 2-Gy fractions; interval with the first treatment) [13,14]. Alternative local treatments utilising interventional radiology techniques may also be discussed if they are not overly demanding in time, resources and personnel.

In conclusion, as COVID-19 takes its toll on our healthcare systems, the optimization of supportive care is necessary.

The adaptation of our practices for radiotherapy of bone metastases should be aimed at drastically reducing the number of radiotherapy sessions. This will contribute to minimizing the risk of infection in both patients and healthcare professionals, while optimizing the use of resources so that they can be best allocated to patients requiring radical cancer treatments and COVID-19 sufferers.

Treatment proposals must in all cases be tailored to each patient, with their agreement and in accordance with the local organisations, resources and health situation. The current context makes therapeutic decisions more difficult and creates a need for therapeutic alternatives, or for treatments to be modified or postponed. These decisions should be validated in multidisciplinary oncology meetings dedicated to bone pathology, which must continue to take place remotely by video-conference if at all possible.

CRedit authorship contribution statement

Sébastien Thureau: Writing - original draft. **Jean Christophe Faivre:** Writing - original draft. **Richard Assaker:** Writing - review & editing. **Emmanuel Biver:** Writing - review & editing. **Cyrille B. Confavreux:** Writing - review & editing. **Françoise Debiais:** Writing - review & editing. **Martine Duterque-Coquillaud:** Writing - review & editing. **Francesco Giammarile:** Writing - review & editing. **Dominique Heymann:** Writing - review & editing. **Frédéric E. Lecouvet:** Writing - review & editing. **Laetitia Morardet:** Writing - review & editing. **Frederic Paycha:** Writing - review & editing. **Jean-Jacques Body:** Writing - review & editing. **Marie-Hélène Vieillard:** Writing - original draft.

Declaration of Competing Interest

All authors declare no conflict of interest.

Acknowledgements

The authors are also grateful to Alexandre Duterque for reviewing the English style.

References

- [1] M. Ueda, R. Martins, P.C. Hendrie, T. McDonnell, J.R. Crews, T.L. Wong, B. McCreery, B. Jagels, A. Crane, D.R. Byrd, S.A. Pergam, N.E. Davidson, C. Liu, F.M. Stewart, Managing cancer care during the COVID-19 pandemic: agility and collaboration toward a common goal, *J. Natl. Compr. Canc. Netw.* (2020 Mar 20) 1–4.
- [2] H. Wang, L. Zhang, Risk of COVID-19 for patients with cancer, *Lancet Oncol.* 21 (4) (2020 Apr) e181.
- [3] A.R. Filippi, E. Russi, S.M. Magrini, R. Corvò, COVID-19 outbreak in northern Italy: first practical indications for radiotherapy departments, *Int. J. Radiat. Oncol. Biol. Phys.* (2020 Mar 19) pii: S0360-3016(20)30930-5.
- [4] S. Lutz, T. Balboni, J. Jones, S. Lo, J. Petit, S.E. Rich, R. Wong, C. Hahn, Palliative radiation therapy for bone metastases: Update of an ASTRO Evidence-Based Guideline, *Pract. Radiat. Oncol.* 7 (1) (2017) 4–12.
- [5] S. Lutz, L. Berk, E. Chang, E. Chow, C. Hahn, P. Hoskin, D. Howell, A. Konski, L. Kachnic, S. Lo, A. Sahgal, L. Silverman, C. von Gunten, E. Mendel, A. Vassil, D.W. Bruner, W. Hartsell, American Society for Radiation Oncology (ASTRO), Palliative radiotherapy for bone metastases: an ASTRO evidence-based guideline. *Int. J. Radiat. Oncol. Biol. Phys.* 79 (4) (2011) 965–976.
- [6] Y.M. van der Linden, Lok, J.J. Steenland, E. Martijn, H. van Houwelingen, H. Marijnen, C.A. Leer, J.W. Dutch Bone Metastasis Study Group, Single fraction radiotherapy is efficacious: a further analysis of the Dutch Bone Metastasis Study controlling for the influence of retreatment, *Int. J. Radiat. Oncol. Biol. Phys.* 59 (2004) 528–537.
- [7] D. Rades, A.J. Conde-Moreno, J. Cacicedo, T. Veninga, B. Segedin, K. Stanic, L. Rudat, S.E. Schild, 1x8 Gy versus 5x4 Gy for metastatic epidural spinal cord compression: a matched-pair study of three prognostic patient subgroups, *Radiat. Oncol.* 13 (2018) 21.
- [8] L. Bollen, S.P.D. Dijkstra, R.H.M.A. Bartels, A. de Graeff, D.L.H. Poelma, T. Brouwer, P.R. Algra, J.M.A. Kuijlen, M.C. Minnema, C. Nijboer, C. Rolf, T. Sluis, M.A.M.B. Terheggen, A.C.M. van der Togt-van Leeuwen, Y.M. van der Linden, W. Taal, Clinical management of spinal metastases-The Dutch national guideline, *Eur. J. Cancer* 104 (2018) 81–90.
- [9] P.J. Hoskin, K. Hopkins, V. Misra, T. Holt, R. McMenemin, D. Dubois, F. McKinna, B. Foran, K. Madhavan, C. MacGregor, A. Bates, N. O'Rourke, J.F. Lester, T. Sevitt, D. Roos, S. Dixit, G. Brown, S. Arnott, S.S. Thomas, S. Forsyth, S. Beare, K. Reczko, A. Hackshaw, A. Lopes, Effect of single-fraction vs multifraction radiotherapy on ambulatory status among patients with spinal canal compression from metastatic cancer: the SCORAD randomized clinical trial, *JAMA* 322 (21) (2019 Dec 3) 2084–2094, <https://doi.org/10.1001/jama.2019.17913>.
- [10] A. Kumar, M.H. Weber, Z. Gokaslan, J.P. Wolinsky, M. Schmidt, L. Rhines, M.G. Fehlings, I. Laufer, D.M. Sciubba, M.J. Clarke, N. Sundaresan, J.J. Verlaan, A. Sahgal, D. Chou, C.G. Fisher, Metastatic spinal cord compression and steroid treatment: a systematic review, *Clin. Spine Surg.* 30 (4) (2017 May) 156–163, <https://doi.org/10.1097/BSD.0000000000000528>.
- [11] G.D. Skeoch, M.K. Tobin, S. Khan, A.A. Linninger, A.I. Mehta, Corticosteroid treatment for metastatic spinal cord compression: a review, *Global Spine J.* 7 (3) (2017 May) 272–279, <https://doi.org/10.1177/2192568217699189> Epub 2017 Apr 20.
- [12] J.J. Willeumier, Y.M. van der Linden, P.D. Dijkstra, Lack of clinical evidence for postoperative radiotherapy after surgical fixation of impending or actual pathologic fractures in the long bones in patients with cancer; a systematic review, *Radiother. Oncol.* 121 (2016) 138–142.
- [13] Z.A. Husain, A. Sahgal, A. De Salles, M. Funaro, J. Glover, M. Hayashi, M. Hiraoka, M. Levivier, L. Ma, R. Martínez-Alvarez, J.I. Paddick, J. Régis, B.J. Slotman, S. Ryu, Stereotactic body radiotherapy for de novo spinal metastases: systematic review, *J. Neurosurg. Spine* 27 (3) (2017) 295–302.
- [14] A. Hashmi, M. Guckenberger, R. Kersh, P.C. Gerszten, F. Mantel, I.S. Grills, J.C. Flickinger, J.H. Shin, D.K. Fahim, B. Winey, K. Oh, B.C. John Cho, D. Létourneau, J. Sheehan, A. Sahgal, Re-irradiation stereotactic body radiotherapy for spinal metastases: a multi-institutional outcome analysis, *J. Neurosurg. Spine* 25 (5) (2016 Nov) 646–653 Epub 2016 Jun 24.