

Excess cases of influenza-like illnesses synchronous with coronavirus disease (COVID-19) epidemic, France, March 2020

Pierre-Yves Boëlle, Cécile Souty, Titouan Launay, Caroline Guerrisi, Clément Turbelin, Sylvie Behillil, Vincent Enouf, Chiara Poletto, Bruno Lina, Sylvie van der Werf, et al.

► **To cite this version:**

Pierre-Yves Boëlle, Cécile Souty, Titouan Launay, Caroline Guerrisi, Clément Turbelin, et al.. Excess cases of influenza-like illnesses synchronous with coronavirus disease (COVID-19) epidemic, France, March 2020. *Eurosurveillance*, European Centre for Disease Prevention and Control, 2020, 25 (14), 10.2807/1560-7917.ES.2020.25.14.2000326 . inserm-02545745

HAL Id: inserm-02545745

<https://www.hal.inserm.fr/inserm-02545745>

Submitted on 17 Apr 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Excess cases of influenza-like illnesses synchronous with coronavirus disease (COVID-19) epidemic, France, March 2020

Pierre-Yves Boëlle¹, Cécile Souty¹, Titouan Launay¹, Caroline Guerrisi¹, Clément Turbelin¹, Sylvie Behillil⁵, Vincent Enouf⁵, Chiara Poletto², Bruno Lina^{8,9}, Sylvie van der Werf^{4,5,6,7}, Daniel Lévy-Bruhl³, Vittoria Colizza², Thomas Hanslik¹, Thierry Blanchon¹

1. Sorbonne Université, Institut Pierre Louis d'Epidémiologie et de Santé Publique, Paris, France

2. INSERM, Institut Pierre Louis d'Epidémiologie et de Santé Publique, Paris, France

3. Santé Publique France, Saint Maurice, France

4. Institut Pasteur, Unité de Génétique Moléculaire des Virus à ARN, Paris, France

5. Institut Pasteur, Centre Coordonnateur du Centre National de Référence des virus des infections respiratoires (dont la grippe), Paris, France

6. UMR CNRS 3569, Paris, France

7. Université Paris Diderot, Sorbonne Paris Cité, Unité de Génétique Moléculaire des Virus à ARN, Paris, France

8. Laboratoire de Virologie, Hospices Civils de Lyon, Institut des Agents Infectieux (IAI), Centre National de Référence des virus respiratoires (dont la grippe), Centre de Biologie et de Pathologie Nord, Groupement Hospitalier Nord, Lyon, France

9. Université de Lyon, Virpath, CIRI, INSERM U1111, CNRS UMR5308, ENS Lyon, Université Claude Bernard Lyon 1, Lyon, France

Correspondence: Pierre-Yves Boelle (pierre-yves.boelle@upmc.fr)

Citation style for this article:

Boëlle Pierre-Yves, Souty Cécile, Launay Titouan, Guerrisi Caroline, Turbelin Clément, Behillil Sylvie, Enouf Vincent, Poletto Chiara, Lina Bruno, van der Werf Sylvie, Lévy-Bruhl Daniel, Colizza Vittoria, Hanslik Thomas, Blanchon Thierry. Excess cases of influenza-like illnesses synchronous with coronavirus disease (COVID-19) epidemic, France, March 2020. *Euro Surveill.* 2020;25(14):pii=2000326. <https://doi.org/10.2807/1560-7917.ES.2020.25.14.2000326>

Article submitted on 15 Mar 2020 / accepted on 08 Apr 2020 / published on 09 Apr 2020

Several French regions where coronavirus disease (COVID-19) has been reported currently show a renewed increase in ILI cases in the general practice-based *Sentinelles* network. We computed the number of excess cases by region from 24 February to 8 March 2020 and found a correlation with the number of reported COVID-19 cases so far. The data suggest larger circulation of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in the French population than apparent from confirmed cases.

The *Sentinelles* network monitors influenza-like illnesses (ILI) and acute respiratory infections (ARI) in general practice in France [1]. The 2019/20 influenza epidemic reached its peak in mid-February in France [2] and it was expected that ILI incidence would decrease thereafter. However, in the first week of March 2020, we observed a renewed increase in ILI cases in some regions. Several nasopharyngeal swabs collected in that period by *Sentinelles* general practitioners (GPs) tested positive for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). Here, we quantified the number of consultations for ILI in France in early March 2020 in excess of what was expected to be caused by the influenza epidemic. As coronavirus disease (COVID-19) cases were first identified in France at the end of January 2020 [3] we examined the relationship between ILI reports and reported cases of COVID-19.


Sentinelles reporting of influenza-like illness

Cases of ILI and ARI are reported in real time by participating GPs (ca 600 GPs) all over France. The case definition for ILI is fever of sudden onset ($>39^{\circ}\text{C}$) with respiratory signs (cough, running nose) and myalgia in a person of any age. Cases of ARI include any disease with respiratory signs and are only monitored in people older than 65 years, unlike ILI. In addition, *Sentinelles* GPs take nasopharyngeal swabs in one ILI case and up to two ARI cases per week for viral characterisation [4]. Influenza virus (see Supplement), respiratory syncytial virus, human rhinovirus and human metapneumovirus are routinely looked for, and SARS-CoV-2 was added at the end of February 2020. In the first week of testing, week 9 (the week starting 24 February 2020), none of 119 submitted swabs tested positive for SARS-CoV-2. In week 10 of 2020 (the week starting 2 March 2020), 93 swabs were collected in ILI patients and 23 in ARI patients, and two swabs in each category were positive for SARS-CoV-2.

For ILI, one of the two positive ILI cases had been collected because the patient reported a direct link with an existing cluster in the east of France. We computed the expected number of ILI consultations using the superposition of a seasonal [5] and an epidemic influenza component [6], as detailed in the Supplement.

FIGURE

Number of consultations for influenza-like illness (per 100,000) in France and 11 French regions, with expected number of consultations fitted on the first eight weeks, week 1–10, 2020 (n = 11 regions)



ARA: Auvergne-Rhone-Alpes; BRE: Brittany; CVL: Centre-Val de Loire; COR: Corsica; GRE: Grand-Est; HDF: Hauts de France; IDF: Ile de France; ILI: influenza-like illness; NAQ: Nouvelle Aquitaine; OCC: Occitanie; PACA: Provence-Alpes-Cote d'Azur; PDL: Pays de la Loire.

Blue: excess consultations, computed by difference between observations (squares) and expected counts. Green: expected number of consultations fitted on the first eight weeks of 2020 (circles).

Excess cases were computed as the difference between observed cases and expected numbers. This modelling approach allowed computing excess cases in 11 of 13 regions but did not provide a good fit in the remaining two regions where a marked influenza peak was not present (see Supplement). The overall ILI incidence showed renewed increase with 33 (95% credible interval (CrI): -8 to 64) consultations per 100,000 in excess during week 9 and with 84 (95% CrI: 447 to 108) consultations per 100,000 in excess in week 10. Four of the

11 regions displayed positive excess (CrI excluding 0) in week 9 and seven regions in week 10 (Figure).

Confirmed COVID-19 case counts were obtained from the Santé Publique France website [7]. The number of excess cases correlated with the cumulated number of COVID-19 cases reported in the same regions ($r=0.59$; $p<0.05$) in week 10. Assuming exponential growth (typical of early epidemics) from week 8 to 10, 2020, we found that the excess number of cases had

TABLE

Excess consultations per 100,000 by region and cumulative number of confirmed coronavirus disease (COVID-19) cases in the corresponding weeks, France, week 9–10, 2020 (n = 11 regions)

Region	Consultations in excess/100,000		Confirmed COVID-19 cases (cumulative)		Population (in millions)
	Week 9	Week 10	Week 9	Week 10	
Auvergne-Rhone-Alpes (ARA)	82 (17 to 147)	115 (36 to 174)	28	146	8
Brittany (BRE)	9 (-32 to 44)	40 (-7 to 73)	5	59	3.3
Centre-Val de Loire (CVL)	72 (43 to 96)	117 (97 to 130)	0	17	2.6
Corsica (COR)	35 (32 to 37)	21 (20 to 23)	0	5	0.3
Grand-Est (GRE)	-55 (-109 to -13)	107 (56 to 138)	5	259	5.5
Hauts de France (HDF)	51 (-3 to 94)	172 (111 to 208)	34	163	6.0
Ile de France (IDF)	40 (32 to 46)	107 (103 to 110)	18	191	12.3
Nouvelle Aquitaine (NAQ)	-3 (-110 to 32)	-9 (-44 to 26)	4	28	6.0
Occitanie (OCC)	54 (30 to 75)	99 (79 to 112)	4	37	5.9
Pays de la Loire (PDL)	-25 (-70 to 5)	18 (-16 to 38)	4	25	3.5
Provence-Alpes-Cote d'Azur (PACA)	36 (-4 to 80)	94 (47 to 140)	8	52	5.0
France	33 (-8 to 64)	84 (44 to 108)	110	982	64.9^a

^a Total population of all 13 regions of mainland France.

an exponential growth rate per week of 0.69 (95% CrI: 0.55–0.86) in Grand-Est (GRE), 0.67 (95% CrI: 0.55 to 0.83) in Ile de France (IDF), 0.68 (95% CrI: 0.56 to 0.83) in Hauts de France (HDF), 0.61 (95% CrI: 0.48 to 0.75) in Occitanie (OCC) and 0.56 (95% CrI: -1.6 to 0.8) in France overall (Table).

For ARI, the two positive swabs had been collected in the regions Bourgogne-Franche Comté (BFC) and GRE, from a total of seven swabs in these two regions. We estimated that 760 (95% CrI: 219–1,706) ARI consultations in those older than 65 years in these two regions (BFC and GRE) could have been caused by COVID-19 during week 10 (2 in 7 for a total of 2,600 ARI visits). We did not pursue modelling excess cases for ARI as the time series (shown in the Supplement) were too noisy to properly break down the data into an expected seasonal part plus excess cases.

Discussion

This is the first time that an increase in ILI cases was observed simultaneously in several regions after the peak of the annual influenza epidemic in the last 30 seasons of routine surveillance in France with the *Sentinelles* network [9]. Several processes may have contributed to this observation, including, among others, characteristics of the influenza season, change in population behaviour or increase in COVID-19 incidence.

Influenza A (H1N1)pdm09 and influenza B viruses co-circulated in 2019/20 in France [10] (see Supplement). Seasons with type B virus circulation can lead to a slow decay in incidence but a renewed increase is atypical [1]. The 2-week-long normal school holidays started between mid-February and early March depending on the region. While holidays may change the dynamics of influenza [9], this generally has an effect in the ascending phase of the epidemic rather than after the peak.

The COVID-19 pandemic may have changed the health-seeking behaviour of patients and, to some extent, the reporting of the contributing GPs although they conform to a case definition. In the crowdsourced surveillance system grippenet.fr [10], there was no evidence of increasing consultations: 30% consulted with a GP in for ILI during week 10, compared with 38% in week 9. As information regarding the coronavirus risk is widespread, a uniform increase in all regions could have been expected in this scenario rather than only in a few regions. We however acknowledge that increased consultation rates in regions where COVID-19 is the most reported may be possible.

The similarity between COVID-19 and influenza symptoms makes it possible that the excess ILI cases were due to COVID-19 cases. The presence of SARS-CoV-2-positive swabs in the patients supports this possibility. The correlation between the number of confirmed cases and the computed excess is also consistent with this scenario. In this case, the excess in cases is compatible with an exponential growth at an estimated rate of ca 0.7 per week. Interestingly, this estimate was very similar in the four most affected regions (GRE, IDF, OCC, HDF). It may prove useful to help calibrate models for studying the impact of the COVID-19 pandemic.

Estimating the true number of COVID-19 cases can only be tentative at this early stage of the epidemic with data from a non-dedicated surveillance system. It is likely that the case definition of ILI does not allow identifying all COVID-19 cases. In addition, only a few cases were virologically tested, adding uncertainty in the estimates. Yet, the rate of excess ILI consultations applied to the French regions yielded a number of cases far in excess of confirmed cases, with 12,840 cases in IDF and around 10,000 cases in the regions GRE and HDF. Likewise, the extrapolated number of COVID-19 presenting as ARI was 760 in those older

than 65 years in the two eastern regions (GRE and BFC): this was already twice the number of reported cases in these regions at the time ($n = 379$), all ages combined.

As we enter a period of generalised circulation of SARS-CoV-2, surveillance based on clinical description and swabbing by GPs will prove essential in assessing the situation.

Acknowledgements

We thank all GPs participating to the Sentinelles network.

Funding: Santé Publique France.

Conflict of interest

None declared.

Authors' contributions

PYB, VC, CP, CS, TB designed the study. PYB, CS, TL analysed the data. CG, CT, TH contributed data on incidence. SB, VE, BL, SVDW characterised viruses. PYB drafted the manuscript. BL, SVDW, DLB contributed to the manuscript. All authors approved the last version.

References

1. Souty C, Amoros P, Falchi A, Capai L, Bonmarin I, van der Werf S, et al. Influenza epidemics observed in primary care from 1984 to 2017 in France: A decrease in epidemic size over time. *Influenza Other Respir Viruses*. 2019;13(2):148-57. <https://doi.org/10.1111/irv.12620> PMID: 30428158
2. Sentinelles R. French GPs Sentinelles network > Public Health Surveillance > Database. Paris: Sorbonne Université. [Accessed: 7 Apr 2020]. Available from: <http://www.sentiweb.fr/france/en/?page=database>
3. Bernard Stoecklin S, Rolland P, Silue Y, Mailles A, Campese C, Simondon A, et al. First cases of coronavirus disease 2019 (COVID-19) in France: surveillance, investigations and control measures, January 2020. *Euro Surveill*. 2020;25(6):2000094. <https://doi.org/10.2807/1560-7917.ES.2020.25.6.2000094> PMID: 32070465
4. Souty C, Masse S, Valette M, Behillil S, Bonmarin I, Pino C, et al. Baseline characteristics and clinical symptoms related to respiratory viruses identified among patients presenting with influenza-like illness in primary care. *Clin Microbiol Infect*. 2019;25(9):1147-53. <https://doi.org/10.1016/j.cmi.2019.01.014> PMID: 30703528
5. Costagliola D, Flahault A, Galinec D, Garnerin P, Menares J, Valleron AJ. A routine tool for detection and assessment of epidemics of influenza-like syndromes in France. *Am J Public Health*. 1991;81(1):97-9. <https://doi.org/10.2105/AJPH.81.1.97> PMID: 1983924
6. Hsieh Y-H, Ma S, Velasco Hernandez JX, Lee VJ, Lim WY. Early outbreak of 2009 influenza A (H1N1) in Mexico prior to identification of pH1N1 virus. *PLoS One*. 2011;6(8):e23853. <https://doi.org/10.1371/journal.pone.0023853> PMID: 21909366
7. Santé Publique France. COVID-19 : point épidémiologique du 7 mars 2020. [COVID-19: daily report 7 March 2020]. Saint-Maurice: Sante Publique France; 7 Mar 2020. French. Available from: <https://www.santepubliquefrance.fr/maladies-et-traumatismes/maladies-et-infections-respiratoires/infection-a-coronavirus/documents/bulletin-national/covid-19-point-epidemiologique-du-7-mars-2020>
8. Sentinelles R. Weekly report on 03/11/2020, 2020w10 (from 03/02/2020 to 03/08/2020). Weekly epidemiological bulletin for influenza, week 10. Paris: Sorbonne Université. [Accessed:

7 Apr 2020]. Available from <http://www.sentiweb.fr/document.php?doc=4844>

9. Cauchemez S, Valleron A-J, Boëlle P-Y, Flahault A, Ferguson NM. Estimating the impact of school closure on influenza transmission from Sentinel data. *Nature*. 2008;452(7188):750-4. <https://doi.org/10.1038/nature06732> PMID: 18401408
10. Guerrisi C, Turbelin C, Blanchon T, Hanslik T, Bonmarin I, Levy-Bruhl D, et al. Participatory syndromic surveillance of influenza in Europe. *J Infect Dis*. 2016;214(suppl_4):S386-92. <https://doi.org/10.1093/infdis/jiw280> PMID: 28830105

License, supplementary material and copyright

This is an open-access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0) Licence. You may share and adapt the material, but must give appropriate credit to the source, provide a link to the licence and indicate if changes were made.

Any supplementary material referenced in the article can be found in the online version.

This article is copyright of the authors or their affiliated institutions, 2020.