



COVID-19: home poisoning throughout the containment period

Gaël Le Roux, Sandra Sinno-Tellier, Alexis Descatha

► **To cite this version:**

Gaël Le Roux, Sandra Sinno-Tellier, Alexis Descatha. COVID-19: home poisoning throughout the containment period. *Lancet Public Health*, Elsevier, 2020, 10.1016/S2468-2667(20)30095-5 . inserm-02565564

HAL Id: inserm-02565564

<https://www.hal.inserm.fr/inserm-02565564>

Submitted on 6 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

COVID-19: home poisoning throughout the containment period

The world is facing a major pandemic situation, with more than 2 million confirmed cases and 139 000 related deaths, as of April 18, 2020.¹ Although health systems have to deal with such disasters to decrease massive contaminations, several countries, such as France, have implemented containment measures.^{2,3} Poison control centres are—perhaps unexpectedly—being involved too in the COVID-19 outbreak as they are observing a change in the typology of the calls they receive, which is directly related to containment measures.

Indeed, disproportionate fear of the coronavirus leads to dramatic behaviour modification, such as excessive house cleaning and misuse of cleaning products for personal hygiene or for food cleaning. Consequently, we are witnessing an increase in the number of exposures to the chlorine that results from mixing bleach with other cleaning products.⁴

With school closures, young children at home are thus more frequently exposed to dangerous—although common—substances including parent medication, hydroalcoholic solution, potentially poisonous plants, and homemade products. Moreover, hazardous self-medication attempts to fight coronavirus also lead to their fair share of domestic accidents, such as essential oil poisoning, daily treatment cutoff, or, conversely, unproven medication use.

Lastly, and in addition to the containment measures, we must also acknowledge that fear of being contaminated in the emergency room is another obstacle for many people who need health care after poisoning.

Worldwide, prevention of such home poisonings is urgently needed to avoid unnecessary emergency medical system use in such critical times. Once this crisis ends, it will be important

that poison centres, and health and safety agencies, provide exact figures and consider what lessons can be drawn for the future.

We declare no competing interests. The authors would like to thank Juliette Bloch from the French Agency for Food, Environmental and Occupational Health and Safety, Maisons-Alfort, France and the French Poison Control Center members Emmanuel Puskarczyk and Christine Tournoud (Nancy), Dominique Vodovar, Jerome Langrand, and Hervé Laborde-Casterot (Paris), Fanny Pelissier and Nicolas Franchitto (Toulouse), Magalie Labadie and Camille Paradis (Bordeaux), Nicolas Simon (Marseille), Patrick Nisse (Lille), Nathalie Paret and Anne-Marie Patat (Lyon), Marie Deguigne and Marion Legeay (Angers).

Copyright © 2020 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC-BY-NC-ND 4.0 license.

Gael Le Roux, Sandra Sinno-Tellier, French Poison Control Centre members, *Alexis Descatha
alexis.descatha@inserm.fr

Poison Control Center, Academic Hospital CHU Angers, Angers, France (GLR, AD); French Agency for Food, Environmental and Occupational Health and Safety, Maisons-Alfort, France (SS-T); University of Angers, Centre Hospitalier Universitaire d'Angers, Université de Rennes, INSERM, École des hautes études en santé publique, Institut de recherche en santé, environnement et travail UMR_S 1085, F49000 Angers, France (AD)

- 1 WHO. Coronavirus WHO novel coronavirus. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019> (accessed April 18, 2020).
- 2 Hellewell J, Abbott S, Gimma A, et al. Feasibility of controlling COVID-19 outbreaks by isolation of cases and contacts. *Lancet Glob Health* 2020; **8**: e488–96.
- 3 Ng Y, Li Z, Chua YX, et al. Evaluation of the effectiveness of surveillance and containment measures for the first 100 patients with COVID-19 in Singapore—January 2–February 29, 2020. *MMWR Morb Mortal Wkly Rep* 2020; **69**: 307–11.
- 4 Kampf G, Todt D, Pfaender S, Steinmann E. Persistence of coronaviruses on inanimate surfaces and their inactivation with biocidal agents. *J Hosp Infect* 2020; **104**: 246–51.



Lancet Public Health 2020

Published Online
April 24, 2020
[https://doi.org/10.1016/S2468-2667\(20\)30095-5](https://doi.org/10.1016/S2468-2667(20)30095-5)

This online publication has been corrected. The corrected version first appeared at thelancet.com/public-health on May 4, 2020