

**Maternal and neonatal outcomes associated with
induction of labor after one previous cesarean delivery:
A French retrospective study**

Emma Vecchioli, Anne-Gaël Cordier, Anne Chantry, Alexandra Benachi,
Isabelle Monier

► **To cite this version:**

Emma Vecchioli, Anne-Gaël Cordier, Anne Chantry, Alexandra Benachi, Isabelle Monier. Maternal and neonatal outcomes associated with induction of labor after one previous cesarean delivery: A French retrospective study. PLoS ONE, Public Library of Science, 2020, 15 (8), pp.e0237132. 10.1371/journal.pone.0237132 . inserm-02973922

HAL Id: inserm-02973922

<https://www.hal.inserm.fr/inserm-02973922>

Submitted on 21 Oct 2020




HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



RESEARCH ARTICLE

Maternal and neonatal outcomes associated with induction of labor after one previous cesarean delivery: A French retrospective study

Emma Vecchioli^{1,2}, Anne-Gaël Cordier^{1‡}, Anne Chantry^{2,3‡}, Alexandra Benachi¹, Isabelle Monier^{1,3}*

1 Department of Obstetrics and Gynaecology, AP-HP, Antoine Bécclère Hospital, University Paris Saclay, Clamart, France, **2** Midwifery School of Baudelocque, Paris-Descartes University, AP-HP, DHU Risks in Pregnancy, Paris, France, **3** Obstetrical, Perinatal and Pediatric Epidemiology Research Team (EPOPé), Université de Paris, Epidemiology and Statistics Research Center (CRESS), INSERM, INRA, Paris, France

 These authors contributed equally to this work.

‡ These authors also contributed equally to this work.

* isabelle.monier@inserm.fr



Abstract

Background

The safety of methods of labor induction in women with previous cesarean deliveries is still debated. We investigated perinatal outcomes associated with labor induction among women with a trial of labor after one cesarean delivery.

Methods

This retrospective study included 339 women with a trial of labor after one prior cesarean and a singleton term fetus in cephalic presentation in 2013–2016 in a French maternity unit. Labor induction was performed with oxytocin, artificial rupture of membranes and/or prostaglandin E2, according to the Bishop score. The primary outcome was a composite of uterine rupture, low Apgar score, neonatal resuscitation or admission to a neonatal unit. The secondary outcomes included cesarean delivery after onset of labor, postpartum hemorrhage and maternal hospital stay after delivery. We used logistic regression to estimate odds ratios adjusted (aOR) for potential confounders.

Results

In our sample, 67.3% of women had spontaneous labor and 32.7% were induced. More than half of the women received oxytocin during labor regardless of the mode of labor. The proportions of the composite outcome and of cesarean after onset of labor were higher in the induced group compared to the spontaneous group (26.1% vs 15.8%, $p = 0.02$ and 45.0% vs 27.6%, $p < 0.01$, respectively). There were 9 uterine ruptures (2.6%) and this proportion was higher in the induced group compared to the spontaneous group, although this difference was not statistically significant (3.6% vs 2.2%, $p = 0.48$). After adjustment, labor

OPEN ACCESS

Citation: Vecchioli E, Cordier A-G, Chantry A, Benachi A, Monier I (2020) Maternal and neonatal outcomes associated with induction of labor after one previous cesarean delivery: A French retrospective study. PLoS ONE 15(8): e0237132. <https://doi.org/10.1371/journal.pone.0237132>

Editor: Ricardo Q. Gurgel, Federal University of Sergipe, BRAZIL

Received: February 11, 2020

Accepted: June 21, 2020

Published: August 7, 2020

Copyright: © 2020 Vecchioli et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: Authors did not have financial disclosures and no specific funding was received for this work.

Competing interests: The authors have declared that no competing interests exist.

induction was associated with higher risks of the composite outcome (aOR = 2.45, 95% CI: 1.29–4.65), cesarean after onset of labor (aOR = 2.06, 95% CI: 1.15–3.68) and maternal hospital stay after delivery ≥ 6 days (aOR = 6.20, 95% CI: 3.25–11.81). No association was found with postpartum hemorrhage.

Conclusion

Labor induction after one prior cesarean was associated with a higher risk of adverse perinatal outcome. Nevertheless, the higher proportion of uterine rupture did not differ significantly from that in the spontaneous labor group.

Introduction

Cesarean section rates increased worldwide from 12.1% in 2000 to 21.1% in 2015 [1]. Consequently, the number of pregnant women with previous cesarean deliveries has also increased. For instance, the proportion of pregnant women with a history of cesarean delivery increased in France from 8% to 11% between 2003 and 2016 [2, 3]. Cesarean births are an important concern because the presence of a uterine scar increases the risk of uterine rupture, which is associated with major risks of perinatal mortality and maternal and infant morbidity [4–8]. A previous cesarean delivery is also associated with higher risks of abnormal placentation [9, 10], hysterectomy [10, 11] and blood transfusion [10] for the subsequent pregnancies.

There are no adequately randomized controlled trials that have assessed the safety and effectiveness of methods of induction in women with a prior cesarean delivery [12]. Whether labor can be induced in women with a previous cesarean section is still debated, in particular in the case of an unfavorable cervix [13, 14]. Nevertheless, several observational studies have shown that labor induction in such cases is associated with a higher risk of uterine rupture when oxytocin is used compared with no oxytocin or spontaneous labor [5, 8, 15–18]. For induction with vaginal prostaglandin E2, the results are inconsistent: some studies found that prostaglandin E2 is associated with a higher risk of uterine rupture compared to other means of induction or spontaneous labor [5–7], while others did not [8, 19–21].

This study assessed perinatal outcomes associated with labor induction versus spontaneous labor among women with one previous cesarean delivery.

Methods

Study design and population

We conducted an observational retrospective study in one tertiary university hospital in France from January 1st, 2013 to October 26th, 2016. Data were obtained from the computerized database of the maternity unit recording all live births and stillbirths at 22 weeks of gestational age (GA) or more. This database includes information on maternal sociodemographic characteristics, medical and obstetrical history, pregnancy complications, delivery and infant outcomes at birth, as well as surgical reports.

This observational study used anonymized data from medical records and was approved by the CNIL (Commission Nationale de l'Informatique et des Libertés) under the notification number 2020 0117175609.

The study population included women with a history of one cesarean delivery and a singleton live fetus in cephalic presentation born at 37 weeks GA or more. Exclusion criteria were:

preterm births <37 weeks GA, multiple pregnancies, stillbirths and terminations of pregnancy, non-cephalic presentation, no history of cesarean delivery, a history of uterine surgery. Women with an elective cesarean delivery for the current pregnancy were also excluded.

Definition of outcomes

The primary outcome was a composite criterion of adverse perinatal outcomes including any of the following: uterine rupture, Apgar score <7 at 5 minutes and neonatal resuscitation in the delivery room or admission to a neonatal unit. The diagnosis of uterine rupture was based on surgical reports and defined as a complete disruption of the prior uterine scar including the peritoneum. Neonatal resuscitation was defined as oxygen administration or intubation in the delivery room. At the time of the study period, cord blood pH and lactate were determined only in cases of risk of fetal distress that included all emergency situations (cesarean section after onset of labor, placental abruption, uterine rupture, vasa previa, etc.). Because these tests were not systematically performed, we did not include cord blood pH and lactate in the composite criteria.

The secondary outcomes included cesarean section after onset of labor, postpartum hemorrhage and duration of maternal stay after delivery above 6 days. Based on French guidelines, postpartum hemorrhage was defined as a blood loss of at least 500 mL estimated by clinicians regardless of the mode of delivery [22]. We considered maternal hospital stay after delivery ≥ 6 days as an adverse outcome because the average length of hospital stay after a cesarean section in France in 2010 was 5.4 days [2].

Covariables

Maternal characteristics included maternal age, parity, body mass index (BMI) before pregnancy and geographical origin. We also reported information on previous deliveries and current pregnancy complications. Neonatal characteristics included gestational age at delivery, birthweight and the infant's sex. Birthweight was expressed as a continuous variable and in percentiles defined using an intrauterine growth reference adapted to the French population [23]. In France, gestational age is determined from the crown-rump length at the first ultrasound scan between 11⁺⁰ and 13⁺⁶ weeks [24].

Indications for labor induction were categorized into 5 classes: prolonged pregnancy, premature rupture of membranes (PROM), maternal medical cause (hypertensive diseases, diabetes, etc.), fetal cause (antenatal suspicion of growth restriction or macrosomia) and other indications. Indications for cesarean section after onset of labor were classified as follows: failure to progress, non-reassuring fetal monitoring, maternal cause and other indications. We also reported information on obstetrical management during labor: the type of analgesia, the use of intravenous oxytocin for labor augmentation and the method used for the induction of labor. In the maternity unit, the protocol for induction of labor in women who had one previous cesarean section and a favorable cervix (Bishop score ≥ 6) consisted of an amniotomy, secondarily associated with intravenous oxytocin. In contrast, vaginal prostaglandin E2 was used in women with an unfavorable cervix (Bishop score <6). Prostaglandin was kept for up to 24 hours with regular fetal heart monitoring. In the case of uterine contractions with cervical modifications or a favorable cervix, labor was continued with an amniotomy associated, if necessary, with oxytocin. A cesarean section was performed if the cervix was unchanged or unfavorable after 24 hours with vaginal prostaglandin E2. At the time of the study period, the Foley catheter and the double-balloon catheter were not used in the maternity unit.

Statistical analysis

Maternal, pregnancy and neonatal characteristics as well as obstetrical management were compared between women with spontaneous labor and those with induced labor in univariate analysis using Student's t-tests for continuous variables and chi-squared or Fisher exact tests for categorical variables. Statistical significance was defined for a p-value <0.05. The association between mode of labor and perinatal outcomes was assessed using a multivariable logistic regression to estimate odds ratios adjusted for potential confounding factors (aOR). We used directed acyclic graphs (DAGs) in order to select co-variables included in the multivariable models based on the literature and clinical assumptions. Interactions between labor induction and several co-variables (parity, pregnancy complications) were tested and were not significant. Because of the small number of uterine ruptures and neonates with a low Apgar score, we did not perform multivariable analysis separately for these two outcomes. However, these outcomes were included in the composite criteria of adverse perinatal outcomes. Finally, we described clinical characteristics and neonatal outcomes including cord blood pH and lactate related to cases with uterine rupture. Analyses were performed using Stata 13.0 software (StataCorp LP, College Station, TX, USA).

Results

Among the 11,843 births at 22 weeks GA or more recorded during the study period, 339 women had a history of one cesarean delivery and a singleton live fetus in cephalic presentation born at 37 weeks GA or more (Fig 1). Among the 339 women, 67.3% had spontaneous




Fig 1. Flow chart.

<https://doi.org/10.1371/journal.pone.0237132.g001>

labor and 32.7% were induced. Maternal pre-pregnancy BMI was higher in the group of women with labor induction compared to women with spontaneous labor (26.1 kg/m² vs. 24.3 kg/m², $p < 0.01$). (Table 1) None of the other maternal characteristics differed between the groups. More than half of the women received oxytocin during labor regardless of the mode of onset of labor.

Among the 111 women who had induction of labor, the primary indication was maternal medical causes (34.2%) followed by prolonged pregnancy (26.1%). (Table 2) Labor was mainly induced with prostaglandin E2 (41.8%), then with both prostaglandin E2 and oxytocin (31.8%) and finally with oxytocin only (26.4%).

Maternal and neonatal outcomes are compared according to the mode of onset of labor in Table 3. The rates of the composite adverse perinatal outcome and cesarean section after onset of labor were higher in women with labor induction compared to those with spontaneous labor (respectively, 26.1% vs. 15.8%, $p = 0.02$ and 45.0% vs. 27.6%, $p < 0.01$). Overall, there were 9 uterine ruptures: 2.2% in the spontaneous group and 3.6% in the induced group ($p = 0.48$). There was a higher proportion of women who stayed more than 6 days after delivery in the induced group (39.1% vs. 9.0%, $p < 0.01$). No difference was found between groups for postpartum hemorrhage. For neonatal outcomes, resuscitation in the delivery room was more frequent in infants born to mothers who were induced than in mothers with spontaneous labor (18.9% vs 9.6%, $p = 0.02$) and there was a non-significant trend to a higher rate of admission to a neonatal unit (14.4% vs. 8.3%, $p = 0.08$).

After adjustment for covariables, the risk of the composite adverse perinatal outcome was increased compared to women with spontaneous labor (aOR = 2.45, (95% confidence interval (CI): 1.29–4.65)). (Table 4) Induction was associated with higher risks of cesarean section (aOR = 2.06, 95% CI: 1.15–3.68) and maternal stay after delivery ≥ 6 days (aOR = 6.20, 95% CI: 3.25–11.81). No association was found with postpartum hemorrhage.

Table 5 reports information on the mode and duration of labor, gestational age at birth, drugs used to induce labor and the results of cord blood tests for the 9 uterine ruptures. Oxytocin was used in 6 cases (including 5 cases in women with spontaneous labor) and rupture of membranes was artificial in 7 cases. Three infants had a cord blood pH < 7.15 and were transferred to a neonatal unit.

Discussion

Main findings

In our maternity unit, one-third of women with a prior cesarean birth and a trial of labor were induced. Uterine rupture was seen in 2.6% of cases and this proportion was higher in the induced group compared to the spontaneous group, although this difference did not reach significance. Oxytocin was administered to more than half of the women. Labor induction was associated with higher risks of the composite adverse perinatal outcome, cesarean section and longer maternal stay after delivery compared to spontaneous labor.

Limitations and strengths

Our study had several limitations. As we used an observational design, we were not able to establish a causal effect between mode of labor and observed perinatal outcomes. Our study was also retrospective, leading to a lower quality of data compared to a prospective study. However, adverse maternal outcomes as uterine rupture are rare and therefore many studies of the association between mode of labor and pregnancy outcomes in women with previous cesarean birth have also used a retrospective design [6, 8, 18, 19, 25–28]. In addition, the size of our sample was too small to compare methods of induction and therefore our analysis was

Table 1. Maternal, obstetrical and neonatal characteristics according to mode of onset of labor.

	Total n (%) or mean \pm sd	Spontaneous labor n (%) or mean \pm sd	Induction of labor n (%) or mean \pm sd	p-value
Total	339 (100)	228 (67.3)	111 (32.7)	
Maternal characteristics				
Maternal age (years)	33.0 \pm 4.4	33.0 \pm 4.4	33.1 \pm 4.6	0.89
Parity	1.3 \pm 0.7	1.3 \pm 0.7	1.4 \pm 0.8	0.43
Pre-pregnancy BMI (kg/m ²)	24.8 \pm 5.2	24.3 \pm 5.0	26.1 \pm 5.5	<0.01
Geographical origin				
France	167 (49.3)	113 (49.6)	54 (48.6)	0.57
Europe	22 (6.5)	16 (7.0)	6 (5.4)	
Africa	121 (35.7)	77 (33.8)	44 (39.6)	
Other countries	29 (8.5)	22 (9.6)	7 (6.3)	
Obstetrical history				
Previous vaginal delivery	74 (21.8)	50 (21.9)	24 (21.6)	0.95
Indication of previous cesarean delivery				
Failure to progress	78 (23.0)	59 (25.9)	19 (17.1)	0.31
Non-reassuring fetal status	150 (44.2)	95 (41.7)	55 (49.5)	
Maternal indication	26 (7.7)	15 (6.6)	11 (9.9)	
Breech presentation	37 (10.9)	26 (11.4)	11 (9.9)	
Other	48 (14.2)	33 (14.5)	15 (13.5)	
Interval between cesarean section and actual delivery (months)	54.6 \pm 36.7	53.2 \pm 34.5	57.6 \pm 40.9	0.31
Pregnancy complications				
None	206 (60.8)	156 (68.4)	50 (45.0)	<0.01 ^a
Diabetes	63 (18.6)	28 (12.3)	35 (31.5)	
Hypertensive diseases	10 (2.9)	4 (1.7)	6 (5.4)	
Antenatal suspicion of FGR or macrosomia	24 (7.1)	14 (6.1)	10 (9.0)	
Other	36 (10.6)	26 (11.4)	10 (9.0)	
Obstetrical management				
Artificial rupture of membranes	164 (48.4)	108 (47.4)	56 (50.4)	0.59
Oxytocin use	187 (55.5)	123 (54.4)	64 (57.7)	0.58
Type of analgesia				
Epidural analgesia	312 (92.0)	215 (94.3)	97 (87.4)	<0.01 ^a
Spinal anesthesia	16 (4.7)	5 (2.2)	11 (9.9)	
Other	4 (1.2)	2 (0.9)	2 (1.8)	
None	7 (2.1)	6 (2.6)	1 (0.9)	
Neonatal characteristics				
Gestational age at birth (weeks)				
37–38	62 (18.3)	37 (16.2)	25 (22.5)	<0.01
39–40	206 (60.8)	159 (69.7)	47 (42.3)	
41+	71 (20.9)	32 (14.0)	39 (35.1)	
Birthweight				
<10 th percentile ^b	34 (10.0)	22 (9.6)	12 (10.8)	0.88
10–90 th percentile ^b	280 (82.6)	190 (83.3)	90 (81.1)	
90 th percentile ^b	25 (7.4)	16 (7.0)	9 (8.1)	
Male sex	170 (50.1)	112 (49.1)	58 (52.2)	0.59

^a Fisher's exact test^b Based on an intrauterine growth curve adapted to the French population for gestational age and sex.<https://doi.org/10.1371/journal.pone.0237132.t001>

Table 2. Indications and methods used among women with induced labor.

	Induction of labor
	n (%)
Total N = 111	
Indications of induction of labor	
Maternal indications	38 (34.2)
Prolonged pregnancy	29 (26.2)
PROM	17 (15.3)
Fetal indications	10 (9.0)
Other	17 (15.3)
Induction with	
Prostaglandin E2 alone	46 (41.8)
Prostaglandin E2 + oxytocin	35 (31.8)
Oxytocin alone	29 (26.4)

<https://doi.org/10.1371/journal.pone.0237132.t002>

performed in women who had induced labor regardless of the method used, as in other studies [19]. The sample size was also not sufficient to detect differences in outcomes such as uterine rupture, however, this outcome was analyzed using a composite criterion of adverse perinatal outcomes. Finally, the results of our single-center study cannot be extrapolated to medical practices in France. Nevertheless, we retrospectively reviewed all surgical reports in medical records while several previous studies included uterine rupture based on a diagnosis code using hospital discharge databases [15, 26].

Interpretation

We found that labor induction was associated with an increased risk of cesarean section compared to spontaneous labor and this result is concordant with previous reports. In a large prospective study in the United States, labor induction was associated with a lower success rate of vaginal birth after a trial of labor [29]. One retrospective study also found a higher risk of cesarean section for women with a trial of labor compared to spontaneous labor [19]. In

Table 3. Maternal and neonatal outcomes according to the mode of onset of labor.

	Total	Spontaneous labor	Induction of labor	p-value
	n (%)	n (%)	n (%)	
Total	339 (100)	228 (67.3)	111 (32.7)	
Maternal outcomes				
Cesarean section after onset of labor	113 (33.3)	63 (27.6)	50 (45.0)	<0.01
Uterine rupture	9 (2.6)	5 (2.2)	4 (3.6)	0.48 ^b
Postpartum hemorrhage	32 (9.4)	24 (10.5)	8 (7.2)	0.33
Maternal hospital stay ≥6 days	63 (18.9)	20 (9.0)	43 (39.1)	<0.01
Neonatal outcomes				
Apgar score <7 at 5 minutes	4 (1.2)	3 (1.3)	1 (0.9)	1 ^b
Resuscitation	43 (12.7)	22 (9.6)	21 (18.9)	0.02
Admission to a neonatal unit	35 (10.3)	19 (8.3)	16 (14.4)	0.08
Composite adverse perinatal outcome ^a	65 (19.2)	36 (15.8)	29 (26.1)	0.02

^a Includes uterine rupture, Apgar score <7 at 5 minutes, neonatal resuscitation in the delivery room or admission to a neonatal unit

^b Fisher's exact test.

<https://doi.org/10.1371/journal.pone.0237132.t003>

Table 4. Association between maternal and neonatal outcomes and induction of labor.

	Induction of labor	
	Crude OR (95% CI)	Adjusted OR (95% CI)
Maternal outcomes		
Cesarean section ^a	2.14 (1.33–3.44)	2.06 (1.15–3.68)
Postpartum hemorrhage ^b	0.66 (0.28–1.52)	0.66 (0.27–1.63)
Maternal hospital stay ≥ 6 days ^c	6.51 (3.58–11.84)	6.20 (3.25–11.81)
Neonatal outcomes		
Resuscitation ^d	2.18 (1.14–4.17)	2.49 (1.25–4.97)
Admission to a neonatal unit ^e	1.85 (0.91–3.76)	2.19 (1.03–4.64)
Composite adverse perinatal outcomes ^e	1.89 (1.08–3.28)	2.45 (1.29–4.65)

Reference group: spontaneous labor.

CI: confidence interval.

^a Adjusted for maternal age, parity, pre-pregnancy BMI, geographical origin, pregnancy complications, previous vaginal delivery, gestational age and birthweight.

^b Adjusted for maternal age, parity, pre-pregnancy BMI, epidural analgesia, oxytocin administration during labor, mode of delivery, episiotomy, duration of labor and birthweight.

^c Adjusted for maternal age, parity, pregnancy complications and geographical origin

^d Adjusted for birthweight, sex and gestational age.

^e Adjusted for maternal age, parity, pre-pregnancy BMI, geographical origin, previous vaginal delivery, oxytocin administration during labor, birthweight, sex and gestational age.

<https://doi.org/10.1371/journal.pone.0237132.t004>

contrast, in a population of low-risk nulliparous women, a recent multicenter randomized trial found a lower risk of cesarean section after induction of labor [30] and this confirms that a first cesarean section has an impact on mode of delivery.

In previous retrospective population-based and single center studies, rates of uterine rupture were between 0.5% and 2% [5, 8, 17, 26]. In our series, this rate was 2.6% and was also high among women with spontaneous labor (2.2%). Several observational studies found an increased risk of uterine rupture related to labor induction after a prior cesarean birth [6, 8, 31], while other studies did not [19, 20]. In our study, almost one-third of women in the induced group received both prostaglandin and oxytocin; oxytocin was frequently used during

Table 5. Characteristics of the nine cases with uterine rupture.

Case	Labor and delivery characteristics				Neonatal outcomes		
	Type of induction	Gestational age at birth	Drugs used	Duration of labor (hours)	Cord blood pH	Cord blood lactate	Admission to a neonatal unit
1	Spontaneous	40+3	Oxytocin and ARM	4	6.99	8.6	No
2	Spontaneous	40+5	Oxytocin and ARM	10	7.29	4.2	No
3	Spontaneous	40+4	Oxytocin	4.5	7.16	8.7	No
4	Spontaneous	40+0	Oxytocin and ARM	8	7.19	5.9	No
5	Spontaneous	40+6	Oxytocin and ARM	5	7.18	3.1	No
6	Induction	40	PGE2	3	7.3	4.0	Yes
7	Induction	38+1	PGE2 and ARM	1	6.98	12.0	Yes
8	Induction	40+5	PGE2 and ARM	6.5	7.28	4.7	No
9	Induction	39+5	PGE2 and then oxytocin and ARM	2	7.13	7.7	Yes

ARM: artificial rupture of membranes; PGE2: prostaglandin E2.

<https://doi.org/10.1371/journal.pone.0237132.t005>

labor, regardless of mode of onset of labor, and this practice may have an effect on the risk of uterine rupture even in women with spontaneous labor. A large prospective study using the National Institute of Child Health and Human Development Maternal–Fetal Medicine Units Network showed that the risk of uterine rupture increased with the use of oxytocin alone and increased more with the use of prostaglandin with or without oxytocin [32]. A nested-control study also found that a maximum dose of oxytocin above 20 mU/min was associated with a four-fold increased risk of uterine rupture [18]. Unfortunately, this information was not available in our study.

We found that labor induction was related to a ≥ 6 -day maternal hospital stay after delivery. This result may be related to the higher proportion of cesarean section in the induced group as cesarean sections result in a longer hospital stay compared to vaginal deliveries. There was also a trend to a higher proportion of pregnancy complications and admission to a neonatal unit in the induced group and this could impact the duration of maternal stay after delivery. We did not find an association between labor induction and the risk of postpartum hemorrhage. This result is not concordant with previous studies, which found a higher risk of postpartum hemorrhage related to labor induction [33]. Many other factors such as duration of the second stage of labor could influence these results and were not investigated in our study.

After one previous cesarean birth, all professional societies recommend an assessment of individual risks to select candidates for a trial of labor [13, 34–37]. The selection of eligible women for a trial of labor after one previous cesarean birth would be relevant to the assessment of benefits and harms associated with the mode of birth. Compared to an elective repeat cesarean section, a trial of labor increases the risk of uterine rupture and is associated with a higher risk of neonatal hypoxic–ischemic encephalopathy [5, 7, 32]. In contrast, it is also important to avoid elective cesarean section because this increases the risk of maternal death and neonatal respiratory morbidity compared to vaginal delivery [38, 39]. However, a study using the 2010 French Perinatal Survey showed that 42% of women eligible for a trial of labor had an elective repeat cesarean delivery [40].

Prostaglandin E1 (i.e. misoprostol) was found to be associated with an increased risk of uterine rupture and there is a consensus not to use it in women with one previous cesarean delivery [13, 34, 41, 42]. In contrast, professional societies have not contraindicated the use of prostaglandin E2, but have recommended avoidance of its use in women with prior cesarean delivery [13, 34–36] in favor of the use of a Foley catheter, which seems to be associated with a lower risk of uterine rupture compared to induction with prostaglandins [6, 43]. However, a recent Cochrane review reported that randomized controlled studies of birth outcomes among women with prior cesarean section were underpowered to detect significant differences between methods of induction [12]. Large studies assessing the use of the Foley catheter to induce labor in women with a prior cesarean delivery are needed.

Conclusion

Labor induction among women with a previous cesarean delivery was associated with increased risk of adverse perinatal outcomes. We estimated that the risk of uterine rupture was high, even in the case of spontaneous labor, highlighting that oxytocin administration during labor in women with a uterine scar should be used with caution and only if necessary.

Supporting information

S1 Dataset.
(XLSX)

Author Contributions

Conceptualization: Emma Vecchioli, Alexandra Benachi, Isabelle Monier.

Data curation: Isabelle Monier.

Formal analysis: Isabelle Monier.

Investigation: Emma Vecchioli, Alexandra Benachi, Isabelle Monier.

Methodology: Emma Vecchioli, Alexandra Benachi, Isabelle Monier.

Validation: Anne-Gaël Cordier, Anne Chantry, Alexandra Benachi.

Writing – original draft: Isabelle Monier.

Writing – review & editing: Anne-Gaël Cordier, Anne Chantry, Alexandra Benachi, Isabelle Monier.

References

1. Boerma T, Ronsmans C, Melesse DY, Barros AJD, Barros FC, Juan L, et al. Global epidemiology of use of and disparities in caesarean sections. *Lancet*. 2018; 392(10155):1341–8. [https://doi.org/10.1016/S0140-6736\(18\)31928-7](https://doi.org/10.1016/S0140-6736(18)31928-7) PMID: 30322584.
2. Blondel B, Lelong N, Kermarrec M, Goffinet F, National Coordination Group of the National Perinatal S. Trends in perinatal health in France from 1995 to 2010. Results from the French National Perinatal Surveys. *Journal de gynécologie, obstétrique et biologie de la reproduction*. 2012; 41(4):e1–e15. <https://doi.org/10.1016/j.jgyn.2012.04.014> PMID: 22613118.
3. Blondel B, Coulm B, Bonnet C, Goffinet F, Le Ray C, National Coordination Group of the National Perinatal S. Trends in perinatal health in metropolitan France from 1995 to 2016: Results from the French National Perinatal Surveys. *Journal of gynecology obstetrics and human reproduction*. 2017; 46(10):701–13. <https://doi.org/10.1016/j.jogoh.2017.09.002> PMID: 29031048.
4. Chauhan SP, Martin JN Jr, Henrichs CE, Morrison JC, Magann EF. Maternal and perinatal complications with uterine rupture in 142,075 patients who attempted vaginal birth after cesarean delivery: A review of the literature. *American journal of obstetrics and gynecology*. 2003; 189(2):408–17. [https://doi.org/10.1067/s0002-9378\(03\)00675-6](https://doi.org/10.1067/s0002-9378(03)00675-6) PMID: 14520209.
5. Lydon-Rochelle M, Holt VL, Easterling TR, Martin DP. Risk of uterine rupture during labor among women with a prior cesarean delivery. *The New England journal of medicine*. 2001; 345(1):3–8. <https://doi.org/10.1056/NEJM200107053450101> PMID: 11439945.
6. Ravasia DJ, Wood SL, Pollard JK. Uterine rupture during induced trial of labor among women with previous cesarean delivery. *American journal of obstetrics and gynecology*. 2000; 183(5):1176–9. <https://doi.org/10.1067/mob.2000.109037> PMID: 11084562.
7. Smith GC, Pell JP, Pasupathy D, Dobbie R. Factors predisposing to perinatal death related to uterine rupture during attempted vaginal birth after caesarean section: retrospective cohort study. *Bmj*. 2004; 329(7462):375. <https://doi.org/10.1136/bmj.38160.634352.55> PMID: 15262772; PubMed Central PMCID: PMC509342.
8. Zelop CM, Shipp TD, Repke JT, Cohen A, Caughey AB, Lieberman E. Uterine rupture during induced or augmented labor in gravid women with one prior cesarean delivery. *American journal of obstetrics and gynecology*. 1999; 181(4):882–6. [https://doi.org/10.1016/s0002-9378\(99\)70319-4](https://doi.org/10.1016/s0002-9378(99)70319-4) PMID: 10521747.
9. National Institutes of H. National Institutes of Health Consensus Development Conference Statement vaginal birth after cesarean: new insights March 8–10, 2010. *Seminars in perinatology*. 2010; 34(5):351–65. <https://doi.org/10.1053/j.semperi.2010.06.002> PMID: 20869552.
10. Silver RM, Landon MB, Rouse DJ, Leveno KJ, Spong CY, Thom EA, et al. Maternal morbidity associated with multiple repeat cesarean deliveries. *Obstetrics and gynecology*. 2006; 107(6):1226–32. <https://doi.org/10.1097/01.AOG.0000219750.79480.84> PMID: 16738145.
11. Knight M, Kurinczuk JJ, Spark P, Brocklehurst P, United Kingdom Obstetric Surveillance System Steering C. Cesarean delivery and peripartum hysterectomy. *Obstetrics and gynecology*. 2008; 111(1):97–105. <https://doi.org/10.1097/01.AOG.0000296658.83240.6d> PMID: 18165397.
12. West HM, Jozwiak M, Dodd JM. Methods of term labour induction for women with a previous caesarean section. *The Cochrane database of systematic reviews*. 2017; 6:CD009792. <https://doi.org/10.1002/14651858.CD009792.pub3> PMID: 28599068.

13. Sentilhes L, Vayssiere C, Beucher G, Deneux-Tharaux C, Deruelle P, Diemunsch P, et al. Delivery for women with a previous cesarean: guidelines for clinical practice from the French College of Gynecologists and Obstetricians (CNGOF). *European journal of obstetrics, gynecology, and reproductive biology*. 2013; 170(1):25–32. <https://doi.org/10.1016/j.ejogrb.2013.05.015> PMID: 23810846.
14. Tsakiridis I, Mamopoulos A, Athanasiadis A, Dagklis T. Vaginal Birth After Previous Cesarean Birth: A Comparison of 3 National Guidelines. *Obstetrical & gynecological survey*. 2018; 73(9):537–43. <https://doi.org/10.1097/OGX.0000000000000596> PMID: 30265740.
15. Kaczmarczyk M, Sparen P, Terry P, Chantingius S. Risk factors for uterine rupture and neonatal consequences of uterine rupture: a population-based study of successive pregnancies in Sweden. *BJOG: an international journal of obstetrics and gynaecology*. 2007; 114(10):1208–14. <https://doi.org/10.1111/j.1471-0528.2007.01484.x> PMID: 17877673.
16. Kayani SI, Alfirevic Z. Uterine rupture after induction of labour in women with previous caesarean section. *BJOG: an international journal of obstetrics and gynaecology*. 2005; 112(4):451–5. <https://doi.org/10.1111/j.1471-0528.2004.00336.x> PMID: 15777443.
17. Dekker GA, Chan A, Luke CG, Priest K, Riley M, Halliday J, et al. Risk of uterine rupture in Australian women attempting vaginal birth after one prior caesarean section: a retrospective population-based cohort study. *BJOG: an international journal of obstetrics and gynaecology*. 2010; 117(11):1358–65. <https://doi.org/10.1111/j.1471-0528.2010.02688.x> PMID: 20716251.
18. Cahill AG, Waterman BM, Stamilio DM, Odibo AO, Allsworth JE, Evanoff B, et al. Higher maximum doses of oxytocin are associated with an unacceptably high risk for uterine rupture in patients attempting vaginal birth after cesarean delivery. *American journal of obstetrics and gynecology*. 2008; 199(1):32 e1–5. <https://doi.org/10.1016/j.ajog.2008.03.001> PMID: 18455132.
19. Delaney T, Young DC. Spontaneous versus induced labor after a previous cesarean delivery. *Obstetrics and gynecology*. 2003; 102(1):39–44. [https://doi.org/10.1016/s0029-7844\(03\)00355-7](https://doi.org/10.1016/s0029-7844(03)00355-7) PMID: 12850605.
20. Schmitz T, Pourcelot AG, Moutafoff C, Biran V, Sibony O, Oury JF. Cervical ripening with low-dose prostaglandins in planned vaginal birth after cesarean. *PloS one*. 2013; 8(11):e80903. <https://doi.org/10.1371/journal.pone.0080903> PMID: 24260505; PubMed Central PMCID: PMC3834249.
21. Yogev Y, Ben-Haroush A, Lahav E, Horowitz E, Hod M, Kaplan B. Induction of labor with prostaglandin E2 in women with previous cesarean section and unfavorable cervix. *European journal of obstetrics, gynecology, and reproductive biology*. 2004; 116(2):173–6. <https://doi.org/10.1016/j.ejogrb.2004.02.019> PMID: 15358459.
22. Sentilhes L, Vayssiere C, Mercier FJ, Aya AG, Bayoumeu F, Bonnet MP, et al. [Postpartum hemorrhage: Guidelines for clinical practice—Text of the Guidelines (short text)]. *Journal de gynécologie, obstétrique et biologie de la reproduction*. 2014; 43(10):1170–9. <https://doi.org/10.1016/j.jgyn.2014.10.009> PMID: 25453206.
23. Ego A, Prunet C, Lebreton E, Blondel B, Kaminski M, Goffinet F, et al. [Customized and non-customized French intrauterine growth curves. I—Methodology]. *Journal de gynécologie, obstétrique et biologie de la reproduction*. 2016; 45(2):155–64. <https://doi.org/10.1016/j.jgyn.2015.08.009> PMID: 26422365.
24. Vayssiere C, Haumonte JB, Chantry A, Coatleven F, Debord MP, Gomez C, et al. Prolonged and post-term pregnancies: guidelines for clinical practice from the French College of Gynecologists and Obstetricians (CNGOF). *European journal of obstetrics, gynecology, and reproductive biology*. 2013; 169(1):10–6. <https://doi.org/10.1016/j.ejogrb.2013.01.026> PMID: 23434325.
25. Agnew G, Turner MJ. Vaginal prostaglandin gel to induce labour in women with one previous caesarean section. *Journal of obstetrics and gynaecology: the journal of the Institute of Obstetrics and Gynaecology*. 2009; 29(3):209–11. <https://doi.org/10.1080/01443610902743789> PMID: 19358026.
26. Al-Zirqi I, Stray-Pedersen B, Forsen L, Vangen S. Uterine rupture after previous caesarean section. *BJOG: an international journal of obstetrics and gynaecology*. 2010; 117(7):809–20. <https://doi.org/10.1111/j.1471-0528.2010.02533.x> PMID: 20236103.
27. Caughey AB, Shipp TD, Repke JT, Zelop C, Cohen A, Lieberman E. Trial of labor after cesarean delivery: the effect of previous vaginal delivery. *American journal of obstetrics and gynecology*. 1998; 179(4):938–41. [https://doi.org/10.1016/s0002-9378\(98\)70192-9](https://doi.org/10.1016/s0002-9378(98)70192-9) PMID: 9790374.
28. Macones GA, Peipert J, Nelson DB, Odibo A, Stevens EJ, Stamilio DM, et al. Maternal complications with vaginal birth after cesarean delivery: a multicenter study. *American journal of obstetrics and gynecology*. 2005; 193(5):1656–62. <https://doi.org/10.1016/j.ajog.2005.04.002> PMID: 16260206.
29. Landon MB, Leindecker S, Spong CY, Hauth JC, Bloom S, Varner MW, et al. The MFMU Cesarean Registry: factors affecting the success of trial of labor after previous cesarean delivery. *American journal of obstetrics and gynecology*. 2005; 193(3 Pt 2):1016–23. <https://doi.org/10.1016/j.ajog.2005.05.066> PMID: 16157104.

30. Grobman WA, Rice MM, Reddy UM, Tita ATN, Silver RM, Mallett G, et al. Labor Induction versus Expectant Management in Low-Risk Nulliparous Women. *The New England journal of medicine*. 2018; 379(6):513–23. <https://doi.org/10.1056/NEJMoa1800566> PMID: 30089070; PubMed Central PMCID: PMC6186292.
31. Palatnik A, Grobman WA. Induction of labor versus expectant management for women with a prior cesarean delivery. *American journal of obstetrics and gynecology*. 2015; 212(3):358 e1–6. <https://doi.org/10.1016/j.ajog.2015.01.026> PMID: 25725658.
32. Landon MB, Hauth JC, Leveno KJ, Spong CY, Leindecker S, Varner MW, et al. Maternal and perinatal outcomes associated with a trial of labor after prior cesarean delivery. *The New England journal of medicine*. 2004; 351(25):2581–9. <https://doi.org/10.1056/NEJMoa040405> PMID: 15598960.
33. Khireddine I, Le Ray C, Dupont C, Rudigoz RC, Bouvier-Colle MH, Deneux-Tharaux C. Induction of labor and risk of postpartum hemorrhage in low risk parturients. *PloS one*. 2013; 8(1):e54858. <https://doi.org/10.1371/journal.pone.0054858> PMID: 23382990; PubMed Central PMCID: PMC3555986.
34. ACOG Practice Bulletin No. 205 Summary: Vaginal Birth After Cesarean Delivery. *Obstetrics and gynecology*. 2019; 133(2):393–5. <https://doi.org/10.1097/AOG.0000000000003079> PMID: 30681536.
35. Royal Australian and New Zealand College of Obstetricians and Gynaecologists. Birth after previous caesarean section. July 2018. Available at: [https://www.ranzcog.edu.au/RANZCOG_SITE/media/RANZCOG-MEDIA/Women%27s%20Health/Statement%20and%20guidelines/Clinical-Obstetrics/Birth-after-previous-Caesarean-Section-\(C-Obs-38\)-Re-write-July-2015.pdf?ext=.pdf](https://www.ranzcog.edu.au/RANZCOG_SITE/media/RANZCOG-MEDIA/Women%27s%20Health/Statement%20and%20guidelines/Clinical-Obstetrics/Birth-after-previous-Caesarean-Section-(C-Obs-38)-Re-write-July-2015.pdf?ext=.pdf).
36. Royal College of Obstetricians and Gynaecologists. Birth after previous caesarean birth. Green top bulletin No.45. London: RCOG; 2015.
37. Society of O, Gynaecologists of C. SOGC clinical practice guidelines. Guidelines for vaginal birth after previous caesarean birth. Number 155 (Replaces guideline Number 147), February 2005. *International journal of gynaecology and obstetrics: the official organ of the International Federation of Gynaecology and Obstetrics*. 2005; 89(3):319–31. <https://doi.org/10.1016/j.ijgo.2005.03.015> PMID: 16001462.
38. Deneux-Tharaux C, Carmona E, Bouvier-Colle MH, Breart G. Postpartum maternal mortality and cesarean delivery. *Obstetrics and gynecology*. 2006; 108(3 Pt 1):541–8. <https://doi.org/10.1097/01.AOG.0000233154.62729.24> PMID: 16946213.
39. Hansen AK, Wisborg K, Ulbjerg N, Henriksen TB. Risk of respiratory morbidity in term infants delivered by elective caesarean section: cohort study. *Bmj*. 2008; 336(7635):85–7. <https://doi.org/10.1136/bmj.39405.539282.BE> PMID: 18077440; PubMed Central PMCID: PMC2190264.
40. Bartolo S, Goffinet F, Blondel B, Deneux-Tharaux C. Why women with previous caesarean and eligible for a trial of labour have an elective repeat caesarean delivery? A national study in France. *BJOG: an international journal of obstetrics and gynaecology*. 2016; 123(10):1664–73. <https://doi.org/10.1111/1471-0528.14056> PMID: 27126956.
41. Blanchette HA, Nayak S, Erasmus S. Comparison of the safety and efficacy of intravaginal misoprostol (prostaglandin E1) with those of dinoprostone (prostaglandin E2) for cervical ripening and induction of labor in a community hospital. *American journal of obstetrics and gynecology*. 1999; 180(6 Pt 1):1551–9. [https://doi.org/10.1016/s0002-9378\(99\)70051-7](https://doi.org/10.1016/s0002-9378(99)70051-7) PMID: 10368503.
42. Aslan H, Unlu E, Agar M, Ceylan Y. Uterine rupture associated with misoprostol labor induction in women with previous cesarean delivery. *European journal of obstetrics, gynecology, and reproductive biology*. 2004; 113(1):45–8. [https://doi.org/10.1016/S0301-2115\(03\)00363-4](https://doi.org/10.1016/S0301-2115(03)00363-4) PMID: 15036710.
43. Jozwiak M, van de Lest HA, Burger NB, Dijksterhuis MG, De Leeuw JW. Cervical ripening with Foley catheter for induction of labor after cesarean section: a cohort study. *Acta obstetrica et gynecologica Scandinavica*. 2014; 93(3):296–301. <https://doi.org/10.1111/aogs.12320> PMID: 24354335.