



Treatment and Outcomes of Children with Febrile Urinary Tract Infection due to Extended Spectrum Beta-lactamase-producing Bacteria in Europe: TOO CUTE Study

Tina Plankar Srovin, Ana Isabel Valdivielso Martínez, Konstantinos Vazouras, Yingfen Hsia, Laura Folgori, Julia Bielicki, Elise Aguadisch, Alasdair Bamford, Ana Brett, Marion Caseris, et al.

► To cite this version:

Tina Plankar Srovin, Ana Isabel Valdivielso Martínez, Konstantinos Vazouras, Yingfen Hsia, Laura Folgori, et al.. Treatment and Outcomes of Children with Febrile Urinary Tract Infection due to Extended Spectrum Beta-lactamase-producing Bacteria in Europe: TOO CUTE Study. *Pediatric Infectious Disease Journal*, 2020, 39 (12), pp.1081-1087. 10.1097/INF.0000000000002838 . inserm-03226769

HAL Id: inserm-03226769

<https://inserm.hal.science/inserm-03226769>

Submitted on 15 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Treatment and Outcomes of Children with Febrile Urinary Tract Infection due to Extended Spectrum Beta-lactamase-producing Bacteria in Europe: TOO CUTE Study

Konstantinos Vazouras, MD, MPH (1,2)*, Yingfen Hsia, PhD (1,3), Laura Folgori, MD (1,4), Julia Bielicki, MD (1,5), Elise Aguadisch, MD (6), Alasdair Bamford, MD, PhD (7), Ana Brett, MD, MSc (8), Marion Caseris, MD (9), Rimante Cerkauskiene, MD, PhD (10), Maia De Luca, MD (11), Elias Iosifidis, MD, MSc, PhD (12), John Kopsidas, MD (2), Ángela Manzanares, MD (13), Tim Planche, MD (14), Andrew Riordan, MD, FRCPC (15), Tina Plankar Srovin, MD, PhD (16), Ana Isabel Valdivielso Martínez, MD (17), Eleni Vergadi, MD, MSc, PhD (18), Mike Sharland, MD (1), and Romain Basmaci, MD, PhD (6, 19)

Corresponding Author: Konstantinos Vazouras

Address: Collaborative Center of Clinical epidemiology and Outcomes Research

5, Chatzigianni Mexi, Post Code 11528, Athens

Tel: + 306981031132

E-mail: k.vazouras@cleoresearch.org

Abbreviated title: ESBL Febrile Urinary UTIs in Children: Treatment and Outcomes

Running Head: TOO CUTE Study

1. Paediatric Infectious Diseases Research Group, Institute for Infection and Immunity, St George's University of London, London, United Kingdom
2. Stavros Niarchos Foundation-Collaborative Center for Clinical Epidemiology and Outcomes Research and Division of Infectious Diseases (CLEO), Athens, Greece

3. School of Pharmacy, Queen's University Belfast, Belfast, UK
4. Paediatric Infectious Disease Unit, Department of Paediatrics, Luigi Sacco Hospital,
University of Milan, Milan, Italy
5. Paediatric Pharmacology, University Children's Hospital Basel, Basel, Switzerland
6. Service de Pédiatrie-Urgences, Hôpital Louis-Mourier, Assistance Publique – Hôpitaux
de Paris, F-92700 Colombes, France
7. Department of Paediatric Infectious Diseases, Great Ormond Street Hospital for Children
NHS Trust, London
8. Serviço de Urgência e Unidade de Infecçciologia, Hospital Pediátrico, Centro Hospitalar e
Universitário de Coimbra, Coimbra, Portugal
9. Service pédiatrie générale, Hôpital Robert Debré, F-75019 Paris, France
10. Clinic of Paediatrics, Faculty of Medicine, Vilnius University, Vilnius, Lithuania
11. Immunology and Infectious Diseases Unit, University Hospital Pediatric Department,
Bambino Gesù Children's Hospital, Rome, Italy
12. Third Department of Pediatrics, Aristotle University, Medical Faculty, School of Health
Sciences, Hippokration Hospital, Thessaloniki, Greece
13. Hospital 12 de Octubre, Facultad de Medicina, Universidad Complutense, Madrid, Spain
14. Medical Microbiology Department, South West London Pathology, Jenner Wing St
George's Hospital, London, UK
15. Paediatric Infectious Diseases and Immunology, Alder Hey Children's NHS Foundation
Trust, Liverpool, UK
16. Department of Infectious Diseases, University Medical Centre, Ljubljana, Slovenia
17. Department of Paediatrics, Regional University Hospital of Malaga, Spain

1
2
3
4 18. Department of Paediatrics, Heraklion University Hospital, University of Crete, Greece

5
6 19. Université de Paris, Infection, Antimicrobiens, Modélisation, Evolution, Unité Mixte de
7
8
9 Recherche 1137, Institut National de la Santé Et de la Recherche Médicale, F-75018 Paris,
10
11 France
12
13
14
15
16
17
18
19
20

21 **Key Words:** pyelonephritis, extended-spectrum beta lactamase, antibiotics, clinical outcomes,
22
23 children
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Objective

The prevalence of extended-spectrum beta-lactamase producing Enterobacteriaceae (ESBL-PE) is increasing globally. ESBL-PE are an important cause of urinary tract infections (UTIs) in children. We aimed to characterize the clinical presentation, treatment and outcomes of childhood UTI caused by ESBL-PE in Europe.

Design

Multi-center retrospective cohort study

Patients and Setting

Children 0-18 years of age with fever, positive urinalysis and positive urine culture for an ESBL-PE uropathogen, seen in a participating hospital from January 2016 to July 2017.

Main outcome measures

Primary outcome measure: day of defervescence was compared between a) initial microbiologically effective treatment (IET) versus initial microbiologically ineffective treatment (IIT), b) single initial antibiotic treatment versus combined initial antibiotic treatment.

Secondary outcome measures: clinical and microbiological failure of initial treatment

Results

We included 142 children from 14 hospitals in 8 countries. Sixty-one children had IET and 77 IIT. There was no statistical difference in time to defervescence for effective/ineffective groups ($p=0.722$) and single/combination therapy groups ($p=0.574$). Two out of 59 (3.4%) and 4/66 (6.1%) patients exhibited clinical failure during treatment ($p=0.683$) when receiving IET or IIT respectively. Eight of 51 (15.7%) receiving IET and 6/58 (10.3%) receiving IIT

patients ($p=0.568$) had recurring symptoms/signs suggestive of a UTI. Recurrence of a UTI occurred 15.5 days (IQR 9.0 – 19.0) after the end of treatment.

Conclusions

Time to defervescence and clinical failure did not differ between IET/IIT groups. Non-carbapenem beta-lactam antibiotics may be used for the empiric treatment of ESBL febrile UTIs, until susceptibility testing results become available.

INTRODUCTION

The extended-spectrum beta-lactamase producing enterobacteria (ESBL-PE) are recognized as a serious threat by the United States Center for Disease Control and Prevention and the World Health Organization^{1,2}. ESBL enzymes are capable of conferring bacterial resistance to commonly used antibiotics such as penicillins and cephalosporins³. Due to concern about high resistance rates to these antibiotics, carbapenems are often used as empiric therapy for severe infections caused by ESBL-PE's^{4,5}. Several studies have reported the growing prevalence of ESBL-PE infections worldwide⁶⁻⁸, including children with urinary tract infection (UTI)^{6,9,10}. A meta-analysis has shown that the estimated pooled prevalence of childhood UTIs caused by ESBL-PE was globally 14%¹⁰, for both community acquired and healthcare acquired infections. The prevalence of childhood ESBL-PE UTIs varies across regions with the lowest rate in the Americas (2%), 12% in Europe, 37% in South-East Asia, and the highest rate in Africa with 76%¹⁰. ESBL-PE UTIs are associated with longer hospital stays and higher costs in adults¹¹. International guidelines recommend a range of empiric options for UTI treatment in children, including amoxicillin/clavulanic acid (co-amoxiclav), cotrimoxazole, ciprofloxacin or 2nd or 3rd generation cephalosporins (2GC or 3GC) for oral and 3GC or aminoglycosides for intravenous treatment of febrile UTIs (fUTIs)¹²⁻¹⁴. However, ESBL-PEs often possess mechanisms which confer co-resistance to a number of the recommended first line antibiotics³. Despite international guidance, currently there are limited studies on evaluating clinical management of children with ESBL-PE UTIs. We conducted a European multicenter study to: a) characterize the clinical and microbiologic presentations of pediatric patients with ESBL-PE UTIs, b) describe treatment patterns, clinical and microbiologic outcomes.

METHODS

Study design and population

This was a multicenter, multinational, retrospective cohort study. Anonymous data were collected using a web-based electronic form in REDCap. Demographic data, diagnosis, comorbidity, medical history, antibiotic prescribing information (drug name, dose, route of administration and dosing), clinical and microbiologic outcomes were recorded.

Inclusion and Exclusion Criteria

Children 0-18 years of age were included in the study if they were seen within a hospital setting (inpatients, outpatients clinics or emergency departments) from 1st January 2016 to 31st July 2017. We included patients with fever $\geq 38^{\circ}\text{C}$ at or prior to admission and positive urine examination. Positive urine examination was defined by the presence of both “a” and “b”: a) isolation of ESBL-producing bacteria either from spontaneously voided urine / midstream clean-catch method with $\geq 10^5$ colony forming units (cfu) per 1 mL of urine OR suprapubic aspirate/urinary catheter with $\geq 10^4$ cfu/mL of urine, b) either abnormal dipstick test (leucocyte esterase $>1+$, or nitrite positive) or abnormal urine microscopy (WBCs > 5 per high-power field in centrifuged or $>10/\text{mm}^3$ in non-centrifuged samples). Patients were excluded from the study in the presence of any other bacterial species $\geq 10^4$ cfu/mL within the same urine sample. Patients were also excluded if they received antibiotics (other than prophylaxis) in the past 48 hours prior to clinical diagnosis or if they had another concurrent infection necessitating antibiotic treatment.

Definitions

1
2
3
4 Patients were classified into two groups according to their initial antibiotic treatment: those who
5
6 received microbiologically effective and those who received microbiologically ineffective
7
8 treatment. Microbiologic effectiveness was defined as *in vitro* susceptibility of the pathogen to at
9
10 least one of the initial choices of empiric antibiotic therapy. Microbiologic ineffectiveness was
11
12 defined as *in vitro* resistance or intermediate sensitivity of the pathogen to each of the initial
13
14 antibiotic(s) prescribed¹⁵. Switched treatment was defined as antibiotics given after initial
15
16 antibiotics were changed. Switched treatment could either be empiric or targeted. Early clinical
17
18 failure of the initial treatment was defined as the persistence/recurrence of fever or
19
20 signs/symptoms suggestive of UTI during treatment. Late clinical failure was defined as the
21
22 recurrence of signs/symptoms within one month after the end of treatment. Microbiologic failure
23
24 during treatment or after treatment was defined as a persistent or recurrent positive urine culture.
25
26
27
28
29
30
31
32

33 *Main Outcome Measures*

34
35 The primary outcome measure was to assess the day of defervescence compared between a)
36
37 initial microbiologically effective treatment versus initial microbiologically ineffective
38
39 treatment, and b) single initial antibiotic treatment versus combined initial antibiotic treatment.
40
41 The secondary outcome measures were clinical and microbiologic failure of the initial treatment.
42
43 Patients were excluded from early clinical outcomes if the assessment was done earlier than 2
44
45 days after starting treatment or if the timing of clinical assessment was uncertain.
46
47
48
49
50
51
52

53 *Statistical analysis*

54
55
56
57
58
59
60
61
62
63
64
65

We compared categorical variables with χ^2 or Fisher's exact test and continuous variables with Mann-Whitney *U* tests. A P value of <0.05 was considered significant for all tests. The analysis was performed using STATA version 14 (Stata Corp LP, College Station, TX, United States).

Ethical approval

Each site was responsible to clarify the type of permission needed to participate in the study. Ethical approvals were obtained- where needed- either from the local Research and Development Department or the Institutional Review Board /Ethics Committee.

RESULTS

There were 14 hospitals from 8 European countries (United Kingdom, 3; Greece, 3; France, 2; Spain, 2; Italy, 1; Portugal, 1; Slovenia, 1; Lithuania, 1) participating in this study (see Table, Supplemental Digital Content 1).

A total of 142 children were included (Figure 1), of which 57 (40.1%) were males. Median age was 1.1 years (interquartile range 0.5 – 4.0). Demographical, clinical and laboratory findings are shown in Table 1. Microbiologic effectiveness of the initial treatment was determined in 138/140 patients, as 2 patients were treated with an antibiotic not tested in the antibiogram. Early or late clinical outcomes could not be assessed in two patients (autonomic dysregulation and palliative care pathway), thus they were excluded from outcomes analysis. Baseline characteristics of “effective” (n=61) and “ineffective group” (n=77) were compared. Patients in the “ineffective group” were younger in age (p=0.034), had fewer urinary tract (UT)-related (p=0.013), or other comorbidities (p<0.001), and lower rates of antibiotic prophylaxis (p=0.008) (Table 1).

The most commonly identified ESBL-PE pathogen was *E. coli*, 122 (85.9%), followed by *Klebsiella* spp., 15 (10.6%), *Enterobacter* spp. 3 (2.1%), *Pseudomonas aeruginosa* 1 (0.7%) and *Morganella morganii* 1 (0.7%). ESBL screening was carried out using European Committee on Antimicrobial Susceptibility Testing (EUCAST) guidelines¹⁶ in 12/14 (85.7%) of participating centers and in 2/14 (14.3%) using Clinical and Laboratory Standards Institute (CLSI) guidelines¹⁷. Production of ESBL was confirmed with the combination disk synergy test in 10/14 (71.4%) of participating centers, in 2/14 (14.3%) using Vitek2 or in 2/14 (14.3%) using other tests (ABCD test, 1; MicroScan WalkAway, 1). Local phenotypic resistance of *E. coli* and *Klebsiella* spp. are shown in Table 2. No molecular typing data was available at any site. Thirty different initial regimens were identified in this population (see Table, Supplemental Digital Content 2). Initial treatment was empiric for 121 (85.2%) patients. A total of 124 (87.3%) patients were initially prescribed a single antibiotic, while 18 (12.7%) patients had a combination of two antibiotics. Seventy-six (53.5%) patients were initially prescribed a single, non-carbapenem beta-lactam agent, mostly co-amoxiclav (17, 12.0%) or cefotaxime (16, 11.3%), while 12 (8.5%) patients received a carbapenem alone. One hundred-one (71.1%) patients had initial intravenous treatment, 40 (28.2%) patients received only oral treatment, and 1 (0.7%) a combination of oral and intravenous antibiotics. Median duration of treatment was 10 days (IQR 7.0 – 11.0).

Time to defervescence

The day of defervescence was available for 99 patients. There was no statistically significant difference ($p=0.722$) in time to defervescence between the effective (median 1.0 days, IQR 0 – 2.0 days) and ineffective groups (median 1.0 days, IQR 0 – 2.0 days). Subgroup analysis revealed no difference in time to defervescence between patients with effective and ineffective

1
2
3
4 treatment regarding age-groups, sex, co-morbidities, pathogens and initial type of treatment
5
6 (Table 3).
7

8
9 No significant difference in time to defervescence ($p=0.574$) was detected between the single
10
11 therapy (median 1.0, IQR 0 – 2.0) and combination therapy groups (median 1.0, IQR 0 – 2.0).
12

13
14 The two groups (single versus combined antibiotics) were similar in terms of age ($p=0.055$), sex
15
16 ($p=0.199$), pathogens ($p=0.285$), UT-related ($p=0.129$) or other comorbidities ($p=0.797$), and
17
18 effectiveness of treatment ($p=0.183$).
19
20
21
22

23 *Clinical Failure*

24
25 Early clinical outcome was recorded for 59/61 (96.7%) and 66/77 (85.7%) patients receiving an
26
27 initial effective or ineffective regimen, respectively. Two patients (3.4%) receiving initial
28
29 effective treatment exhibited early clinical failure, whereas 4/66 (6.1%) with initial ineffective
30
31 therapy had early clinical failure ($p=0.683$) (Table 1). No significant trends were identified in
32
33 subgroup analysis among baseline characteristics between the two groups (Table 3). All
34
35 characteristics and outcomes of patients treated with ineffective treatments are shown in
36
37 Supplemental Digital Content (Table 3).
38
39
40
41

42
43 Fourteen (14/113, 12.4%) children had late clinical failure 15.5 days (IQR 9-19) after the end of
44
45 treatment. Among the 16 patients whose total treatment course included only ineffective
46
47 therapies (either initial or switched treatment) and had full clinical follow-up, 2/16 (12.5%) had
48
49 late clinical failure. No difference in late clinical failure ($p=1.000$) was found between patients
50
51 having only microbiologically ineffective antibiotics (2/16, 12.5%) and those having at least one
52
53 effective antibiotic (12/97, 12.4%) during their total course of treatment.
54
55
56
57
58
59
60
61
62
63
64
65

Microbiologic Failure

Among 43 patients having a repeated urine culture during treatment, 12 exhibited a microbiologic failure. A repeated urine culture was also collected after the end of treatment in 24 children. In 17/24 patients the urine culture was positive. Among these patients, 9 had the same pathogen and 8 had either a new pathogen or the same pathogen but with a different susceptibility pattern.

DISCUSSION

Main Findings

In our study, we did not observe any significant difference in time to defervescence and rates of clinical failure between patients receiving initial effective compared with ineffective treatment for ESBL UTIs. The great majority of patients (95.2%) had an adequate clinical and microbiologic response despite receiving antibiotics to which their UTI was phenotypically resistant to *in vitro*.

ESBL fUTIs pathogens and current treatment options

E. coli was the most common reported pathogen in our study. This is consistent with other studies^{15,18–20}. We observed high resistance rates (>50%) of *E. coli* to co-amoxiclav, cotrimoxazole and ciprofloxacin. Resistance rates were 10% for aminoglycosides and 50% for piperacillin-tazobactam. Resistance rates to fosfomycin, mecillinam, and temocillin were low (<10%). The resistance rates in our study are similar to other European cohort studies^{18,21–23}.

1
2
3
4 Parenteral carbapenems are considered as the treatment of choice for invasive ESBL
5
6 infections^{24,25}. However, due to the increased need for carbapenem-sparing strategies, the use of
7
8 alternative agents is important. In our study, a wide variety of oral and intravenous antibiotics
9
10 were used for the initial or targeted treatment of ESBL fUTIs, such as beta-lactam/beta-lactam
11
12 inhibitor combinations, cephalosporins, aminoglycosides and quinolones.
13
14

15
16 Piperacillin-tazobactam has been investigated for ESBL *E.coli* UTIs in adults¹⁹, while a recent
17
18 multi-center study of ESBL fUTIs has shown similar clinical efficacy for children receiving
19
20 amikacin or carbapenem monotherapy as initial empiric treatment¹⁸. Oral antibiotics, such as
21
22 ciprofloxacin or cotrimoxazole have also been suggested for targeted treatment of ESBL fUTIs
23
24 in children¹⁸. Finally, the combination of co-amoxiclav and cefixime²⁶ and older, off patent
25
26 antibiotics (pivmecillinam, temocillin and fosfomycin) have investigated as potential ESBL-PE
27
28 treatment options due to their favorable *in vitro* activity²¹. It should be noted that clinicians
29
30 should take into account a patient's clinical condition, type of ESBL-PE strain and MIC while
31
32 treating with a carbapenem-alternative⁴.
33
34
35
36
37
38
39
40

41 *Outcomes and microbiologically ineffective treatments*

42
43 The results of our study are similar to a French cohort study¹⁸, where children became afebrile in
44
45 1.8 days (1.0 day in our study). The time to defervescence and length of hospital stay did not
46
47 differ between effective or ineffective treatments¹⁸. In another pediatric cohort of non-resistant
48
49 UTIs, no difference in fever duration was detected between effective (48 hours, IQR 24-240) and
50
51 ineffective treatment groups (78 hours, IQR 48 - 132)¹⁵. A similar effect of ineffective initial
52
53 therapy on the time to resolution of symptoms has been found in adult studies²⁷. Inappropriate
54
55
56
57
58
59
60
61
62
63
64
65

empirical antibiotic therapy did not adversely affect clinical and microbiologic cure rate or UTI recurrence in a Korean cohort study of adults with acute ESBL pyelonephritis²⁸.

In a multicenter study in the United States, 316 children with a UTI (63% of which were febrile) who received initially discordant antibiotics, only 2.2% required an escalation of therapy.

However, patients with complicated backgrounds were excluded from this cohort, while the median of follow-up was 3 days. In our study, although most children became afebrile within 2 days, recurrence of a UTI occurred in 14/113 (12.4%) children 15.5 days (IQR 9.0 – 19.0) after the end of treatment²⁹. A relapse of a UTI has been documented in 4/146 (2.7%) within 2 weeks after treatment in another study³⁰. The lower rate of relapse in this previous study could be due to lower rates of UT-related (29.5%) or other comorbidities (10.3%), or the shorter follow-up period (2 weeks versus 1 month)³⁰. In another study¹⁸, no recurrence was observed within 10 days after the end of treatment. Whether this may be due to the short follow-up period is unclear. Furthermore, this difference may also be explained by a lower comorbidity rate (15.7%) in the French cohort than in our study (35.9%) and the more frequent use of combined antibiotics (40.2% versus 12.7%).

In our study, 16 patients had exclusively ineffective treatments during their treatment course.

These children had identical recurrence rates after the end of treatment with those treated with at least one effective antibiotic during their course. Our multi-center study supports the findings of a previous, smaller, single-center study³¹. The achieved concentration of antibiotics in the urinary tract may explain these results. Most antibiotics used to treat UTIs, such as beta-lactams and aminoglycosides, are renally excreted, thus antibiotic concentrations at the site of infection are higher than can be achieved in the serum or cerebrospinal fluid^{5,32}.

Strengths and limitations

To our knowledge, this is the first multicenter European study to explore ESBL fUTIs treatment patterns and clinical outcomes. Our study has shown the wide variation of antibiotics used as initial treatment for ESBL fUTIs. We stratified outcomes according to the microbiologic effectiveness of initial treatment given. We observed favorable initial clinical outcomes of ineffective antibiotics against ESBL fUTIs in children, as previously shown in childhood and adult studies^{15,28}.

However, there are a number of limitations in our study. The main limitation is the lack of any systematic sampling method, which does not allow for results to be generalized to other populations. We also need to note the potential sampling bias. Misclassification of uropathogens or coding error may have affected patient's inclusion in the study. Another limitation is the noted differences between initial "effective" and "ineffective" antibiotic groups at their baseline characteristics, namely age and comorbidities. In this context, the comparison of outcome measures between these groups has to be interpreted with caution. Furthermore, patients were included only if they were seen and followed-up within the same hospital. This may have affected the clinical failure rates if children were treated for a recurrence in another hospital or in the community. Finally, no imaging data was available for these children and any evidence of renal scarring associated with ineffective treatment could not be determined.

Implications for future studies

Our study has shown a low rate (6.1%) of clinical failures with initial microbiologically ineffective treatment for ESBL febrile UTIs in children. As shown in the current and other studies¹⁸, most children have a clinical response regardless of the initial treatment given. As

1
2
3
4 such, the estimated effect size between effective and ineffective groups is so low that the sample
5
6 size required to power future trials would need to be very high in order to detect significant
7
8 differences in outcomes³³. These data suggest that further randomized trials on ESBL febrile
9
10 UTIs comparing the clinical efficacy of empiric prescribing strategies may not be straightforward
11
12 or, indeed, feasible as the great majority of children become afebrile in less than two days¹⁸.
13
14 Unlike adults where fever is not mandatory for patient's inclusion, as fever may be absent
15
16 especially in the elderly³⁴, fever is currently the main diagnostic criterion, but not a suitable
17
18 endpoint, for future pediatric UTI studies. Recent studies have evaluated the relapse rate of UTI
19
20 after the ineffective treatment of an ESBL upper UTI in adults and children younger than 2 years
21
22 old, respectively^{28,30}. Larger-scale observational studies are needed to explore UTI relapse and
23
24 selection of resistance in all age-groups after the treatment of resistant UTIs with ineffective
25
26 regimens. Furthermore, potential long-term adverse events, such as renal scarring, should further
27
28 be explored.
29
30
31
32
33
34
35
36
37

38 **CONCLUSION**

39
40 Time to defervescence and clinical failure did not differ among children receiving
41
42 microbiologically effective or ineffective antibiotics for ESBL UTIs. Common beta-lactam, non-
43
44 carbapenem, antibiotics may empirically be used for SBL febrile UTIs treatment, until
45
46 susceptibility testing results become available. Future research should evaluate the continuation
47
48 of microbiologically ineffective treatments after susceptibility results are available, as well as the
49
50 possibility of long-term complications.
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

CONTRIBUTIONS

KV and RB conceptualized and designed the study and performed data management and analysis. KV, RB, LF, JB and YS contributed to the design of data collection tools. KV and RB wrote the manuscript. KV, RB, YH, JB, LF and MS interpreted the data and critically revised the manuscript for important intellectual content. KV, EA, AlB, AnB, MC, RC, MDL, JK, AMC, TP, AR, ER, TPS, AV and EV contributed data voluntarily and critically revised the manuscript for important intellectual content. All authors reviewed and approved the final manuscript before submission and agree to be accountable for all aspects of the work.

ACKNOWLEDGEMENTS

We would like to thank all colleagues who assisted to data collection: Kirkira Banou, Microbiology Department, “Aghia Sophia Children’s Hospital”, Athens, Greece; Maria Bitsori, Heraklion University Hospital, University of Crete, Greece; Emmanuel Roilides, Third Department of Pediatrics, Aristotle University, Medical Faculty, School of Health Sciences, Hippokration Hospital, Thessaloniki, Greece, Thessaloniki, Greece; Livia Gargiullo, University Hospital Pediatric Department, Bambino Gesù Children's Hospital, Rome, Italy; Pablo Rojo, Hospital 12 de Octubre, Facultad de Medicina, Universidad Complutense, Madrid, Spain; Breda Zakotnik, Department of Infectious Diseases, University Medical Centre, Ljubljana, Slovenia; Begoña Carazo Gallego, Regional University Hospital of Malaga, Spain; Fernanda Rodrigues, Hospital Pediátrico, Centro Hospitalar e Universitário de Coimbra, Coimbra, Portugal; Fiona Shaw, Alder Hey Children's NHS Foundation Trust, Liverpool, UK; Garth Dixon, Great Ormond Street Hospital for Children NHS Trust, London; Bogdan Cojocaru, Hôpital Louis-Mourier, Assistance Publique – Hôpitaux de Paris, F-92700 Colombes, France; Nora Poey, Service

pédiatrie générale, Hôpital Robert Debré, F-75019 Paris, France; Viktoras Sutkus, Faculty of Medicine, Vilnius University, Vilnius, Lithuania

DECLARATION OF INTERESTS

JB's husband is a senior corporate counsel at Novartis International AG (Basel, Switzerland) and holds Novartis stock and stock options. AV has collaborated with Pfizer and GSK in vaccine conferences. TP is the clinical lead of a NHS diagnostic microbiology laboratory at South West London Pathology. He is on advisory boards for Roche, Pfizer, and Singulex for diagnostics. All other authors declare no competing interests.

FUNDING

No funding was received for this study.

REFERENCES

1. Centers for Disease Control and Prevention, Office of Infectious Disease Antibiotic Resistance Threats in the United States, 2013. Apr, 2013. Available at: <http://Www.Cdc.Gov/Drugresistance/Threat-Report-2013>. Accessed March 31, 2019.
2. World Health Organization. Antimicrobial Resistance: Global Report on Surveillance. 2014. Available at: <http://Www.Who.Int/Drugresistance/Documents/Surveillancereport/En> (Accessed 31 March 2019).
3. Paterson DL, Bonomo RA. Extended-Spectrum -Lactamases: a Clinical Update. Clin Microbiol Rev. 2005;18(4):657-686.

4. Pana ZD, Zaoutis T. Treatment of extended-spectrum β -lactamase-producing Enterobacteriaceae (ESBLs) infections: what have we learned until now? F1000Research. 2018;7:1347. doi:10.12688/f1000research.14822.1
5. Hsu AJ, Tamma PD. Treatment of Multidrug-Resistant Gram-Negative Infections in Children. Clin Infect Dis. 2014;58(10):1439-1448.
6. European Centre for Disease Prevention and Control. Surveillance of Antimicrobial Resistance in Europe – Annual Report of the European Antimicrobial Resistance Surveillance Network (EARS-Net) 2017. Stockholm: ECDC; 2018. Available from: <https://ecdc.europa.eu/sites/portal/files/documents/ears-net-report-2017-update-jan-2019.pdf> (Last Accessed 21st April 2019).
7. Fedler KA, Biedenbach DJ, Jones RN. Assessment of pathogen frequency and resistance patterns among pediatric patient isolates: report from the 2004 SENTRY Antimicrobial Surveillance Program on 3 continents. Diagn Microbiol Infect Dis. 2006;56(4):427-436.
8. Morrissey I, Hackel M, Badal R, Bouchillon S, Hawser S, Biedenbach D. A Review of Ten Years of the Study for Monitoring Antimicrobial Resistance Trends (SMART) from 2002 to 2011. Pharmaceuticals. 2013;6(11):1335-1346.
9. Lukac PJ, Bonomo RA, Logan LK. Extended-Spectrum -Lactamase-Producing Enterobacteriaceae in Children: Old Foe, Emerging Threat. Clin Infect Dis. January 2015.
10. Flokas ME, Detsis M, Alevizakos M, Mylonakis E. Prevalence of ESBL-producing Enterobacteriaceae in paediatric urinary tract infections: A systematic review and meta-analysis. J Infect. 2016;73(6):547-557.

11. MacVane SH, Tuttle LO, Nicolau DP. Impact of extended-spectrum β -lactamase-producing organisms on clinical and economic outcomes in patients with urinary tract infection: Outcomes of ESBL-EK UTI. *J Hosp Med*. 2014;9(4):232-238.
12. Roberts KB; Subcommittee on Urinary Tract Infection., Steering Committee on Quality Improvement and Management. Urinary Tract Infection: clinical practice guideline for the diagnosis and management of an initial UTI in febrile infants and young children. *PEDIATRICS*. 2011;128(3):e749-e770.
13. National Institute for Health and Care Excellence (NICE). Urinary Tract Infection in under 16s: Diagnosis and Management. 2007. Available at: <https://www.nice.org.uk/Guidance/Cg54/Evidence/Full-Guideline-Pdf-196566877>. Accessed April 22, 2019.
14. World Health Organization Model List of Essential Medicines for Children, 7th List, 2019. Geneva: World Health Organization; 2019.
15. Jerardi KE, Auger KA, Shah SS, et al. Discordant antibiotic therapy and length of stay in children hospitalized for urinary tract infection. *J Hosp Med*. 2012;7(8):622-627.
16. European Committee on Antimicrobial Susceptibility Testing (EUCAST) Clinical breakpoints and dosing of antibiotics. Available at: http://www.eucast.org/clinical_breakpoints/.
17. Clinical and Laboratory Standards Institute , Performance standards for antimicrobial susceptibility testing. Available at: <https://clsi.org/standards/products/microbiology/documents/>.
18. Madhi F, Jung C, Timsit S, et al. Febrile urinary-tract infection due to extended-spectrum beta-lactamase-producing Enterobacteriaceae in children: A French prospective multicenter study. Huang Y-C, ed. *PLOS ONE*. 2018;13(1):e0190910. doi:10.1371/journal.pone.0190910

- 1
2
3
4 19. Seo YB, Lee J, Kim YK, et al. Randomized controlled trial of piperacillin-tazobactam,
5
6 cefepime and ertapenem for the treatment of urinary tract infection caused by extended-spectrum
7
8 beta-lactamase-producing *Escherichia coli*. *BMC Infect Dis*. 2017;17(1):404.
9
10 doi:10.1186/s12879-017-2502-x
11
12
13
14 20. Vazouras K, Basmaci R, Bielicki J, et al. Antibiotics and Cure Rates in Childhood Febrile
15
16 Urinary Tract Infections in Clinical Trials: A Systematic Review and Meta-analysis. *Drugs*.
17
18 2018;78(15):1593-1604.
19
20
21 21. Zykov IN, Sundsfjord A, Småbrekke L, Samuelsen Ø. The antimicrobial activity of
22
23 mecillinam, nitrofurantoin, temocillin and fosfomycin and comparative analysis of resistance
24
25 patterns in a nationwide collection of ESBL-producing *Escherichia coli* in Norway 2010–2011.
26
27 *Infect Dis*. 2016;48(2):99-107.
28
29
30
31 22. Koçak M, Büyükkaragöz B, Çelebi Tayfur A, et al. Causative pathogens and antibiotic
32
33 resistance in children hospitalized for urinary tract infection: Urinary tract infection in children.
34
35 *Pediatr Int*. 2016;58(6):467-471.
36
37
38 23. Pérez Heras I, Sanchez-Gomez JC, Beneyto-Martin P, Ruano-de-Pablo L, Losada-Pinedo
39
40 B. Community-onset extended-spectrum β -lactamase producing *Escherichia coli* in urinary tract
41
42 infections in children from 2015 to 2016: Prevalence, risk factors, and resistances. *Medicine*
43
44 (Baltimore). 2017;96(50):e8571.
45
46
47
48 24. Vardakas KZ, Tansarli GS, Rafailidis PI, Falagas ME. Carbapenems versus alternative
49
50 antibiotics for the treatment of bacteraemia due to Enterobacteriaceae producing extended-
51
52 spectrum β -lactamases: a systematic review and meta-analysis. *J Antimicrob Chemother*.
53
54 2012;67(12):2793-2803.
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4 25. Perez F, Bonomo RA. Can We Really Use ss-Lactam/ss-Lactam Inhibitor Combinations
5
6 for the Treatment of Infections Caused by Extended-Spectrum ss-Lactamase-Producing
7
8 Bacteria? Clin Infect Dis. 2012;54(2):175-177.
9
- 10
11 26. Bingen E, Bidet P, Birgy A, Sobral E, Mariani P, Cohen R. In vitro interaction between
12
13 cefixime and amoxicillin-clavulanate against extended-spectrum-beta-lactamase-producing
14
15 Escherichia coli causing urinary tract infection. J Clin Microbiol. 2012;50(7):2540-2541.
16
17
18 27. Ramos-Martínez A, Alonso-Moralejo R, Ortega-Mercader P, Sánchez-Romero I, Millán-
19
20 Santos I, Romero-Pizarro Y. Pronóstico de las infecciones urinarias con tratamiento antibiótico
21
22 discordante. Rev Clínica Esp. 2010;210(11):545-549.
23
24
25 28. Kim S-H, Oh S, Huh K, et al. Inappropriate empirical antibiotic therapy does not
26
27 adversely affect the clinical outcomes of patients with acute pyelonephritis caused by extended-
28
29 spectrum β -lactamase-producing Enterobacteriales. Eur J Clin Microbiol Infect Dis.
30
31
32 2019;38(5):937-944.
33
34
35 29. Wang ME, Lee V, Greenhow TL, et al. Clinical Response to Discordant Therapy in Third-
36
37 Generation Cephalosporin-Resistant UTIs. Pediatrics. 2020;145(2):e20191608.
38
39
40 doi:10.1542/peds.2019-1608
41
42
- 43 30. Hyun HS, Kim JH, Cho MH, et al. Low relapse rate of urinary tract infections from
44
45 extended-spectrum beta-lactamase-producing bacteria in young children. Pediatr Nephrol.
46
47 July 2019.
48
49
- 50 31. Tratselas A, Iosifidis E, Ioannidou M, et al. Outcome of Urinary Tract Infections Caused by
51
52 Extended Spectrum β -Lactamase-producing Enterobacteriaceae in Children: Pediatr Infect
53
54 Dis J. 2011;30(8):709-712.
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4 32. Long SS, Pickering LK, Prober CG. Principles and Practice of Pediatric Infectious
5 Diseases. 3rd Ed. New York, NY: Churchill Livingstone/ Elsevier; 2009.
6
7
8
9 33. Bhalerao S, Kadam P. Sample size calculation. Int J Ayurveda Res. 2010;1(1):55.
10
11 34. European Medicines Agency. Guideline on the evaluation of medicinal products indicated
12 for treatment of bacterial infections. Final guidance. 2012. Available at:
13
14 www.ema.europa.eu/docs/en_
15
16 GB/document_library/Scientific_guideline/2009/09/WC500003417.pdf. Accessed July 17,
17
18 2019.
19
20
21
22 35. Tratselas A, Simitsopoulou M, Giannakopoulou A, et al. Effect of Ceftriaxone on the
23 Outcome of Murine Pyelonephritis Caused by Extended-Spectrum- β -Lactamase-Producing
24 Escherichia coli. Antimicrob Agents Chemother. 2014;58(12):7102-7111.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1. Study Demographics

Demographics/Characteristics	Overall (n=142)	Initial Effective (n=61)*	Initial Ineffective (n=77)*	p-value
Age in years (Median, IQR)	1.1 (0.5-4.0)	1.3 (0.7 – 5.0)	1.0 (0.3-2.0)	0.034
Sex				
Male	57 (40.1)	22 (36.1)	34 (44.2)	0.385
Female	85 (59.9)	41 (63.9)	43 (55.8)	
Temperature in °C (Median, IQR)	39.0 (38.4 – 39.5)	39.0 (38.4 – 39.5)	39.0 (38.4 – 39.5)	0.9182
Type of UTI				
Bacteraemic UTI	7 (4.9)	3/59 (5.1)	3/74 (4.1)	0.102
Catheter-related UTI	7 (4.9)	0/59 (0)	6/74 (8.2)	
Vesicostomy or ureterostomy related	5 (3.5)	4/59 (6.8)	1/74 (1.3)	
Kidney Stone Infection	1 (0.7)	0/59 (0)	1/74 (1.3)	
DMSA-confirmed UTI	5 (3.5)	2/59 (3.4)	3/74 (4.1)	
None of the above	117 (82.4)	52/59 (88.1)	63/74 (85.1)	
Medical History				
Urinary Tract (UT)-abnormality	54 (38.0)	31 (50.8)	22/75 (29.3)	0.013
Other comorbidity**	51 (35.9)	31 (50.8)	17 (22.1)	< 0.001
Critically ill/immunosuppressed	18 (12.7)	11/61(18.0)	5 (6.5)	0.058

Prophylactic antibiotics	27 (19.0)	18 (29.5)	8/75 (10.7)	0.008
History of recurrent UTIs	17 (12.0)	8 (13.1)	9 (11.7)	0.801
Pathogens				
<i>E. coli</i>	122 (85.9)	53 (86.9)	68 (88.3)	0.801
Non <i>E.coli</i>	20 (14.1)	8 (13.1)	9 (11.7)	
Laboratory markers				
CRP serum level (mg/L) (Median, IQR)	74.0 (32.0- 152.0)	46.7 (14.2- 126.0)	67.5 (17.4- 152.0)	0.723
Abnormal creatinine	22/97 (22.7)	11/41 (26.8)	10/53 (18.9)	0.455
Initial Treatment (route)				
Parenteral	101 (71.1)	46 (75.4)	52 (67.5)	0.500
Oral	40 (28.2)	15 (24.6)	24 (31.2)	
Parenteral and oral	1 (0.7)	0 (0)	1 (1.3)	
No. of initial antibiotic(s) prescribed				
One antibiotic	124 (87.3)	51 (83.6)	71 (92.2)	0.180
> 1 antibiotics	18 (12.7)	10 (16.4)	6 (7.8)	
Outcomes				
Time to defervescence in days (Median, IQR)	1.0*** (0-2.0)	1.0 (0-2.0)	1.0 (0-2.0)	0.722
Early clinical failure	6/125(4.8)	2/59(3.4)	4/66 (6.1)	0.683
Late clinical failure	14/113 (12.4)	8/51 (15.7)	6/58 (10.3)	0.568
Microbiological failure	12/43 (27.9)	5/16 (31.3)	7/27 (25.9)	0.737
Complications****	5/137 (3.6)	5/61 (8.2)	0/72 (0)	0.019
Ward of Admission				

General Paediatrics	57 (40.1)	17/61 (27.9)	39/77 (50.7)	0.028
Outpatients Clinic	37 (26.1)	15/61 (24.5)	22/77 (28.6)	
Paediatric Surgical/Urology	11 (7.7)	7/61 (11.5)	4/77 (5.2)	
Paediatric Intensive Care Unit	8 (5.6)	5/61 (8.2)	2/77 (2.6)	
Neonatal Intensive Care Unit	7 (4.9)	3/61 (4.9)	3/77 (3.9)	
Paediatric Infectious Diseases	4 (2.8)	3/61 (4.9)	1/77 (1.3)	
Haematology-Oncology	2 (1.4)	2/61 (3.3)	0/77 (0)	
Nephrology	2 (1.4)	2/61 (3.3)	0/77 (0)	
Other Special Wards	14 (9.9)	7/61 (11.5)	6/77 (7.7)	

NOTE: Values are listed as number (percentage) or “Median, IQR” or “Mean, SD” if continuous variable. Percentages are calculated with the total number of patients mentioned in the head of each column, except when specified.

Abbreviations: UTI, urinary tract infection; CRP, C reactive protein; SD, standard deviation; IQR, interquartile range

* One-hundred forty-two patients were included in the study. Two patients were excluded from outcomes analysis, while two more could not be classified as effective/ineffective as the initial antibiotic given was not tested in the antibiogram.

** cardiovascular, respiratory, gastrointestinal, neurological, endocrine, metabolic, hematological, immune, perinatal or surgical comorbidities

*** on 99 patients, 45 (45.4%) effective and 54 (54.6%) ineffective

**** secondary sepsis, pyonephrosis, renal failure, admission to an intensive care unit and death after care was withdrawn

Table 2. Antimicrobial resistance rates of *Escherichia coli* and *Klebsiella* spp. in children with febrile ESBL UTIs.

Species	<i>E.coli</i>		<i>Klebsiella</i> spp	
Antibiotic	I-R/total	% of Resistance	I-R/total	% of Resistance
Co-amoxiclav	66/119	55.5%	10/15	66.7%
Temocillin	3/33	9.1%	0/1	-
Mecillinam	2/39	5.1%	0/1	-
TZP	19/100	19.0%	6/13	46.1%
Cefoxitin	23/64	35.9%	4/9	-
Ertapenem	6/89	6.7%	0/9	-
Imipenem	1/92*	1.1%	0/11	0%
Nalidixic acid	24/34	70.6%	1/4	-
Ciprofloxacin	60/107	56.1%	9/15	60.0%
Trimethoprim	59/83	71.1%	4/7	-
Co-trimoxazole	71/110	64.6%	12/15	80.0%
Nitrofurantoin	3/101	3.0%	3/5	-
Gentamicin	37/120	30.8%	13/15	86.7%
Amikacin	14/98	14.3%	5/13	38.5%
Fosfomycin	5/59	8.5%	2/10	20.0%

Abbreviations: I, intermediate; R, resistant; TZP, piperacillin-tazobactam

*intermediate resistance reported

Table 3. Early clinical failure and time to defervescence in ESBL febrile UTIs

Patients' Characteristics	Early clinical failure				Time to defervescence (in days)		
	Total	IET	IIT	p-value	IET, Mean (SD)	IIT, Mean (SD)	p-value
Age							
<29 days	0/3 (0%)	0/1 (0%)	0/2 (0%)	1.000	-	-	-
1 month- 2 years	2/67 (3.0%)	0/25(0%)	2/42 (4.8%)	0.525	1.56 (1.34)	1.03 (0.85)	0.188
>2 years	4/55 (7.3%)	2/33(6.1%)	2/22 (9.1%)	1.000	1.23 (1.48)	1.58 (1.50)	0.309
Sex							
Male	2/55 (3.6%)	0/21 (0%)	2/34 (5.9%)	0.519	1.13 (1.02)	0.68 (0.69)	0.115
Female	4/70 (5.7%)	2/38 (5.3%)	2/32 (6.3%)	1.000	1.48 (1.57)	1.62 (1.29)	0.404
Co-morbidities							
UT-related	3/50 (6.0%)	1/31 (3.2%)	2/19 (10.5%)	0.549	1.40 (1.47)	0.91 (0.70)	0.462
Non UT-related	4/46 (8.7%)	2/30 (6.7%)	2/16 (12.5%)	0.602	1.30 (1.34)	1.24 (1.25)	0.936
Pathogens							

<i>E. coli</i>	4/107 (3.7%)	1/50 (2.0%)	3/57 (5.3%)	0.621	1.32 (1.38)	1.21 (1.16)	0.930
Non <i>E.coli</i>	2/18 (11.1%)	1/9 (11.1%)	1/9 (11.1%)	1.000	1.57 (1.62)	1.00 (1.15)	0.459
Initial Treatment (by route)							
IV/IM	2/91 (2.2%)	1/43 (2.3%)	1/48 (2.1%)	1.000	1.36 (1.25)	1.07 (0.90)	0.388
Oral (single)	4/33 (12.1%)	1/16 (6.3%)	3/17 (17.6%)	0.601	1.33 (2.34)	1.78 (1.99)	0.460
Initial treatment (by antibiotic class)							
Single	6/109 (5.5%)	2/49 (4.1%)	4/60(6.7%)	0.689	1.44 (1.46)	1.20 (1.19)	0.561
<i>Aminopenicillins</i> + <i>BLI</i> ¹	2/21 (9.5%)	1/14 (7.1%)	1/7 (14.3%)	1.000	1.43 (2.15)	1.17 (1.47)	0.940
<i>Cephalosporins</i> ²	2/43 (4.7%)	-	2/43 (4.7%)	-	-	1.25 (1.20)	-
<i>Aminoglycosides</i> ³	0/20 (0%)	0/16 (0%)	0/4 (0%)	1.000	1.79 (1.37)	1.50 (1.00)	0.726
<i>Carbapenems</i> ⁴	1/11 (9.1%)	1/11 (9.1%)	-	-	1.36 (1.29)	-	-
<i>Other</i> ⁵	1/14 (7.1%)	0/8 (0%)	1/6 (16.7%)	0.429	0.50 (0.58)	0.33 (0.58)	0.683
Combinations ⁶	0/16 (0%)	0/10	0/6	1.000	1.00	1.00 (0.71)	0.833

					(1.12)		
--	--	--	--	--	--------	--	--

¹ *Co-amoxiclav, piperacillin-tazobactam*

² *Cefotaxime, ceftriaxone, cefuroxime, cefixime, cefalexin*

³ *Gentamicin, amikacin*

⁴ *Ertapenem, imipenem/cilastatin, meropenem*

⁵ *Co-trimoxazole, Nitrofurantoin, ciprofloxacin, fosfomycin, amoxicillin*

⁶ *Ampicillin + cefotaxime, ampicillin + gentamicin, ampicillin + amikacin, co-amoxiclav + amikacin, ceftriaxone + amikacin, cefoxitin + amikacin, piperacillin-tazobactam + ciprofloxacin, piperacillin-tazobactam + amikacin, meropenem + Vancomycin, meropenem + amikacin, meropenem + gentamicin, imipenem/cilastatin + amikacin, ceftriaxone + oral ofloxacin*

Abbreviations: IET, Initial Effective Treatment; IIT, Initial Ineffective Treatment; BLI – beta-lactamase inhibitor; UT – urinary tract; SD, standard deviation




Figure 1. Formation of the cohort

Supplemental Table 1. Participating Centres and Records

Hospital	Country	Records (n=142)
University Medical Centre of Ljubljana	Slovenia	14
Vilnius University Children's Hospital	Lithuania	15
Great Ormond Street Hospital	United Kingdom	10
Alder Hey Children's Hospital	United Kingdom	16
St George's University of London	United Kingdom	5
Hôpital Robert Debré	France	14
Hôpital Louis Mourier	France	8
Ospedale Pediatrico Bambino Gesù	Italy	9
Hippokration Hospital	Greece	1
Agia Sophia Children's Hospital	Greece	17
University Hospital of Herakleion	Greece	7
Hospital Universitario 12 de Octubre	Spain	10
Hospital Regional Universitario de Málaga	Spain	8
Hospitais da Universidade de Coimbra	Portugal	8

Supplemental Table 2. Initial and Switched Regimens for ESBL febrile UTIs

Antibiotics	Initial N=142 (%)	Switched (1st) N=100	Switched (2nd) N=32	Switched (3rd) N=3	Switched (4th) N=1
Single treatments	124 (87.3)	95 (95.0)	29 (90.6)	-	-
Amoxicillin	1 (0.7)	-	-	-	-
Co-amoxiclav	17 (12.0)	21 (21.0)	6 (18.8)	-	-
Ampicillin-sulbactam	-	1 (1.0)	-	-	-
Piperacillin-tazobactam	5 (3.5)	5 (5.0)	1 (3.1)	-	-
Pivmecillinam	-	2 (2.0)	-	-	-
Temocillin	-	2 (2.0)	-	-	-
Cefalexin	3 (2.1)	-	-	-	-
Cefuroxime	12 (8.5)	2 (2.0)	-	-	-
Cefixime	11 (7.8)	2 (2.0)	-	-	-
Cefotaxime	16 (11.3)	1 (1.0)	-	-	-
Ceftriaxone	11 (7.8)	-	-	-	-
Ertapenem	1 (0.7)	1 (1.0)	-	-	-
Imipenem/cilastatin	1 (0.7)	2 (2.0)	-	1	-
Meropenem	10 (7.0)	8 (8.0)	1 (3.1)	-	-
Gentamicin	16 (11.3)	5 (5.0)	-	-	-
Amikacin	7 (4.9)	10 (10.0)	1 (3.1)	1	-
Ciprofloxacin	4 (2.8)	11 (11.0)	2 (6.3)	-	-
Levofloxacin	-	1 (1.0)	-	-	-
Trimethoprim	-	1 (1.0)	3 (9.4)	-	-

Co-trimoxazole	4 (2.8)	9 (9.0)	10 (31.3)	-	-
Nitrofurantoin	4 (2.8)	10 (10.0)	3 (9.4)	1	-
Fosfomycin	1 (0.7)	1 (1.0)	2 (6.3)	-	-
Combinations	18 (12.7)	5 (5.0)	3 (9.4)	-	-
Ampicillin + Cefotaxime	3 (2.1)	-	-	-	-
Ampicillin + Gentamicin	3 (2.1)	-	-	-	-
Ampicillin + Amikacin	1 (0.7)	1 (1.0)	-	-	-
Piperacillin-tazobactam + Ciprofloxacin	1 (0.7)	-	-	-	-
Piperacillin-tazobactam + Amikacin	1 (0.7)	-	-	-	-
Co-amoxiclav + Amikacin	2 (1.4)	1 (1.0)	-	-	-
Ceftriaxone + Amikacin	1 (0.7)	1 (1.0)	-	-	-
Cefoxitin + Amikacin	1 (0.7)	-	-	-	-
Ceftriaxone + Ofloxacin	1 (0.7)	-	-	-	-
Meropenem + Vancomycin	1 (0.7)	-	-	-	-
Meropenem + Amikacin	1 (0.7)	-	-	-	-
Meropenem + Gentamicin	1 (0.7)	2 (2.0)	-	-	-
Imipenem/cilastatin + Amikacin	1 (0.7)	-	-	-	-
Co-amoxiclav +	-	-	2 (6.3)	-	1

Cefixime					
Pivmecillinam + Nitrofurantoin	-	-	1 (3.1)	-	-

Supplemental Table 3. Characteristics and outcomes of patients treated with an initial ineffective treatment.

Antibiotic	Dose (mg/kg/24 h)	Route	Vulnerable	Pathogen	MIC	EUCAST breakpoints
Early clinical success						
Amikacin	15	IV	No	<i>E. coli</i>	-	>16
Amoxicillin	47	Oral	No	<i>E. coli</i>	-	
Cefalexin	-	Oral	No	<i>E. coli</i>	-	
Cefalexin	-	Oral	No	<i>E. coli</i>	-	
Cefalexin	22	Oral	No	<i>E. coli</i>	-	
Cefixime	-	Oral	No	<i>E. coli</i>	-	>1
Cefixime	8	Oral	No	<i>E. coli</i>	2	>1
Cefixime	8	Oral	No	<i>E. coli</i>	>64	>1
Cefixime	8	Oral	No	<i>E. coli</i>	-	>1
Cefotaxime	196	IV	No	<i>E. coli</i>	-	>2
Cefotaxime	147	IV	No	<i>E. coli</i>	-	>2
Cefotaxime	200	IV	No	<i>E. coli</i>	-	>2
Cefotaxime	150	IV	No	<i>E. coli</i>	-	>2
Cefotaxime	-	IV	No	<i>E. coli</i>	-	>2
Cefotaxime	-	IV	No	<i>E. coli</i>	-	>2
Cefotaxime	155	IV	No	<i>E. coli</i>	-	>2
Cefotaxime	147	IV	No	<i>E. coli</i>	-	>2

Cefotaxime	155	IV	No	<i>E. coli</i>	-	>2
Cefotaxime	-	IV	No	<i>E. coli</i>	-	>2
Cefotaxime	151	IV	No	<i>E. coli</i>	-	>2
Cefotaxime	198	IV	No	<i>E. coli</i>	-	>2
Cefotaxime	160	IV	No	<i>E. coli</i>	-	>2
Cefotaxime	208	IV	Yes	<i>E. coli</i>	-	>2
Cefotaxime	200	IV	No	<i>E. coli</i>	-	>2
Cefotaxime	94	IV	No	<i>E. coli</i>	-	>2
Ceftriaxone	80.5	IV	No	<i>E. coli</i>	128	>2
Ceftriaxone	200	IV	No	<i>E. coli</i>	32	>2
Ceftriaxone	50	IV	No	<i>E. coli</i>	-	>2
Ceftriaxone	54	IV	No	<i>E. coli</i>	-	>2
Ceftriaxone	50	IV	No	<i>E. coli</i>	-	>2
Ceftriaxone	49	IM	No	<i>E. coli</i>	-	>2
Ceftriaxone	83.9	IV	No	<i>Klebsiellapneumoniae</i>	128	>2
Ceftriaxone	50	IV	No	<i>E. coli</i>	-	>2
Ceftriaxone	78	IV	No	<i>Enterobacteraerogenes</i>	8	>2
Cefuroxime	150	IV	No	<i>E. coli</i>	-	
Cefuroxime	31	Oral	No	<i>E. coli</i>	-	
Cefuroxime	31	Oral	No	<i>E. coli</i>	-	
Cefuroxime	32	Oral	No	<i>E. coli</i>	-	
Cefuroxime	50	IV	No	<i>E. coli</i>	-	

Cefuroxime	94	IV	Yes	<i>E. coli</i>	-	
Cefuroxime	147	IV	No	<i>E. coli</i>	-	
Cefuroxime	89	IV	No	<i>E. coli</i>	-	
Cefuroxime	142	IV	No	<i>E. coli</i>	-	
Cefuroxime	-	IV	No	<i>E. coli</i>	-	
Ciprofloxacin	30	IV	No	<i>E. coli</i>	> 4	>0.5
Ciprofloxacin	30	IV	Yes	<i>E. coli</i>	-	>0.5
Co-amoxiclav	120	IV	No	<i>E. coli</i>	-	>8
Co-amoxiclav	101.3	IV	No	<i>E. coli</i>	16	>8
Co-amoxiclav	-	IV	No	<i>Klebsiellapneumonia</i> <i>e</i>	-	>8
Co-amoxiclav	Adult dosing	Oral	No	<i>Morganellamorganii</i>	≥ 32	-
Co-trimoxazole	46	Oral	No	<i>E. coli</i>	-	>4
Gentamicin	4.8	IV	Yes	<i>E. coli</i>	-	>4
Gentamicin	6.4	IV	No	<i>E. coli</i>	-	>4
Gentamicin	4	IM	No	<i>Klebsiellapneumonia</i> <i>e</i>	-	>4
Nitrofurantoin	7	Oral	No	<i>Klebsiellapneumonia</i> <i>e</i>	-	
Piperacillin- tazobactam	209	IV	Yes	<i>E. coli</i>	-	>16
Piperacillin- tazobactam	350	IV	No	<i>Klebsiellapneumonia</i> <i>e</i>	32	>16

Ampicillin &Cefotaxime	209 & 209	IV	No	<i>E. coli</i>	-	Na
Ampicillin &Cefotaxime	207 & 207	IV	No	<i>E. coli</i>	-	Na
Ampicillin &Cefotaxime	200 & 150	IV	No	<i>E. coli</i>	(-) & 64	Na
Ampicillin & Gentamicin	205 & 5	IV	No	<i>E. coli</i>	-	Na
Ampicillin & Gentamicin	153.2 & 3.9	IV	No	<i>Klebsiellaoxytoca</i>	(-) & $\geq$ 16	Na
Ceftriaxone &Ofloxacin	Adult dosing	IV & oral	No	<i>E. coli</i>	-	Na
Early clinical failure						
Cefixime	7.8	Oral	No	<i>E. coli</i>	24	>1
Cefuroxime	29	Oral	No	<i>E. coli</i>	-	
Co-amoxiclav	88.2	IV	No	<i>Klebsiellapneumonia</i> <i>e</i>	> 32	>8
Co-trimoxazole	23	Oral	No	<i>E. coli</i>	-	>4

Vulnerable patients: critically ill, immunocompromised

Abbreviations: Na, not applicable