**Evolution and phenotypic characterization of HBV
quasispecies in compliant patients experiencing
unexplained entecavir treatment failure Running title:
Entecavir, viral resistance & whole genome**

Julien Marlet, Emmanuelle Roch, Clément Lier, Alain Moreau, Benjamin Combe, Lynda Handala, Sandrine Lefeuvre, Morgan Maugey, Laure Elkrief, Louis D'alteroche, et al.

► **To cite this version:**

Julien Marlet, Emmanuelle Roch, Clément Lier, Alain Moreau, Benjamin Combe, et al.. Evolution and phenotypic characterization of HBV quasispecies in compliant patients experiencing unexplained entecavir treatment failure Running title: Entecavir, viral resistance & whole genome. Antiviral Research, 2021, 192, pp.105106. 10.1016/j.antiviral.2021.105106 . inserm-03285455

HAL Id: inserm-03285455

<https://www.hal.inserm.fr/inserm-03285455>

Submitted on 13 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Evolution and phenotypic characterization of HBV quasispecies in compliant patients
experiencing unexplained entecavir treatment failure**

Running title: Entecavir, viral resistance & whole genome

Julien Marlet ^{a,b,#}, Clément Lier ^{a,b}, Emmanuelle Roch ^a, Alain Moreau ^a, Benjamin Combe ^a,
Sandrine Lefeuvre ^c, Morgan Maugey ^a, Laure Elkrief ^d, Louis d'Alteroche ^d, Pascal Potier ^e,
Denys Brand ^{a,b}, Catherine Gaudy-Graffin ^{a,b}.

^a INSERM U1259, Université de Tours et CHRU de Tours, France

^b Service de Bactériologie-Virologie-Hygiène, CHRU de Tours, France

^c Laboratoire de Biochimie, CHR d'Orléans, France

^d Service d'Hépatogastroentérologie, CHRU de Tours, France

^e Service d'Hépatogastroentérologie, CHR d'Orléans, France

Corresponding author: Julien Marlet, Julien.marlet@univ-tours.fr

ABSTRACT

Entecavir treatment failure can be observed in compliant patients despite an absence of detectable resistance mutations by Pol/RT Sanger sequencing. We hypothesized that these unexplained treatment failures could rely on other mechanisms of viral resistance, especially on mutations selected outside of the Pol/RT domain. Partial virological response to entecavir was observed in three patients treated with immunosuppressive drugs, without selection of Pol/RT resistance mutations. Mutations selected in the whole HBV genome during entecavir treatment and potentially associated with resistance were searched for using deep sequencing and characterized using a phenotypic resistance assay. Mutations Q206K (pre-core/core), Q120K (pre-S1/pre-S2, T-cell epitope) and A300E (spacer domain) were selected during entecavir treatment in patient #1 but were not associated with an increased level of resistance to entecavir or an increase in HBV replication capacity. Core promoter mutations T1753G, A1762T and G1764A were present before treatment in patient #1. HBs Ag immune escape mutations were present before treatment in patients #2 (sK122R, sT126I, sP127S and sG145R) and #3 (sM133I). These results provided a unique insight into the evolution of HBV quasispecies during entecavir treatment, addressing the whole HBV genome and not only the Pol/RT domain. Our results suggest that mutations selected outside of the Pol/RT domain could contribute to entecavir treatment failure, not by an increased level of resistance, but by an immune escape mechanism, in synergy with immunosuppressive drugs.

INTRODUCTION

Chronic hepatitis B virus infection affects 292 million people worldwide and is associated with an increase in the risk of cirrhosis and hepatocellular carcinoma (HCC) (Razavi-Shearer et al., 2018). Nucleoside or nucleotide analogs are used as a first-line treatment to prevent these complications (Li et al., 2020; Papatheodoridis et al., 2020; Xu et al., 2015). Two of these molecules, entecavir and tenofovir (disoproxil fumarate or alafenamide) have been associated with higher rates of viral response and a higher drug genetic barrier than previous antiviral drugs (lamivudine, adefovir). Entecavir resistance mutations in the HBV Pol/RT domain are rarely selected ($< 1.2\%$, after five years of treatment) (Tenney et al., 2009) except in patients with a history of lamivudine treatment (Hayashi et al., 2015; L. Liu et al., 2019; Y. Liu et al., 2019; Rose et al., 2018). These mutations have been associated with primary non-response and viral breakthrough on entecavir treatment, but not with a partial virological response (PVR) to entecavir (Lampertico et al., 2017). This last form of treatment failure is defined as a decrease in HBV DNA levels of more than 1 log₁₀ IU/mL, with HBV DNA detectable after at least 12 months of treatment in compliant patients (Lampertico et al., 2017). It occurs in 10-33% of patients treated with entecavir (Chang et al., 2006; Lai et al., 2006) and has been associated with HBe Ag positivity, and with low levels of ALAT and a high HBV DNA viral load at baseline (Chen et al., 2014; Lee et al., 2015; Li et al., 2015; Zoutendijk et al., 2011).

The management of patients experiencing entecavir treatment failure is based on the identification of known resistance mutations by Pol/RT Sanger sequencing (Lampertico et al., 2017; Lok et al., 2007) and, if relevant, the characterization of new resistance mutations in phenotypic assays (Zoulim and Locarnini, 2013). However, this conventional strategy has several limitations. First, it does not take into account polymorphisms selected outside of the

Pol/RT domain and associated with different clinical phenotypes of chronic hepatitis B (Huang et al., 2013; Peng et al., 2017; Podlaha et al., 2019). Second, unlike deep sequencing, it is unsuitable for the sensitive detection of minor variants before (Chevaliez et al., 2019) or after treatment failure (Lee et al., 2018; Lowe et al., 2016).

This conventional approach fails to detect any Pol/RT resistance mutations in some patients experiencing treatment failure on entecavir. We hypothesized that a more thorough exploration, based on deep sequencing of the whole HBV genome before and after treatment failure, would provide a more comprehensive overview of the viral factors associated with treatment failure.

We applied this strategy to compliant patients experiencing entecavir treatment failure without the selection of mutations of the HBV Pol/RT domain. Relevant mutations selected during entecavir treatment were characterized in a phenotypic resistance assay, based on whole HBV genome sequences from clinical isolates.

MATERIALS AND METHODS

Patients and samples

Plasma samples were collected between 2011 and 2018, at Tours University Hospital, for the diagnosis or follow-up of HBV infection (Marlet et al., 2020). We included patients experiencing entecavir treatment failure with no detectable Pol/RT resistance mutations (Lampertico et al., 2017). HBV infection and liver disease were followed, and therapeutic drug monitoring was performed, as previously described (Marlet et al., 2020). This study was approved by the Tours University ethics committee (no. 2019 055).

Genotypic drug resistance testing

Genotypic drug resistance testing was performed by amplifying the Pol/RT domain (aa rt1-rt344) and subjecting it to Sanger sequencing, as previously described (Villeneuve et al., 2003). We searched for resistance mutations as previously described (Marlet et al., 2020) and interpreted them in accordance with the latest EASL guidelines (Lampertico et al., 2017).

Deep sequencing of the whole HBV genome

Two overlapping regions of the HBV genome were amplified by PCR with primer P1 (CCGGAAAGCTTGAGCTCTTCTTTTTCACCTCTGCCTAATCA) plus primer 786R (GGACTCAAGATGTTGTACAGACTTGGCC), and with primer P2 (CCGGAAAGCTTGAGCTCTTCAAAAAGTTGCATGGTGCTGG) plus primer BB (CTGGATGTGTCTGCGGCG), described elsewhere (Günther et al., 1995; Marlet et al., 2020). These PCRs were performed with 300 nM of each primer, 2.5 µL of DNA extract and 12.5 µL of Platinum™ PCR SuperFi PCR Master Mix 2X (Thermo Fisher Scientific, Massachusetts, USA). Cycling reactions were performed on a T3000 (Biometra) with the following cycle parameters: 2 min at 94°C, followed by 40 cycles of 40 s at 94°C, 90 s at 60°C, 90 s at 68°C, with an increment of 12 s after each cycle and a final extension step of 10 min at 68°C. We fused these two overlapping regions of the HBV genome together by PCR, using 1 µL of each diluted PCR product (equimolar) obtained in the first step and 12.5 µL of Platinum™ SuperFi PCR Master Mix 2X. Fusion PCR was performed on the T3000 with the following cycle parameters: 30 s at 98°C, followed by 10 cycles of 10 s at 98°C, 45 s at 72°C and 2 min at 72°C. The whole genome of HBV was amplified by PCR with 300 nM primers P1 and P2, and 25 µL of fusion PCR product. Cycling reactions were performed on a T3000 Biometra with the following cycle parameters: 2 min at 94°C, followed by 40 cycles of 40 s at 94°C, 90 s at

110 60°C, 180 s at 68°C, with an increment of 12 s after each cycle and a final extension step of 10
111 min at 68°C.

112 Next-generation sequencing was performed on the Miseq platform (Illumina, San Diego, USA),
113 with a previously described workflow (Guinoiseau et al., 2017). Briefly, the sequencing library
114 was prepared with the Nextera XT DNA sample preparation kit (Illumina, San Diego, USA),
115 in accordance with the manufacturer's protocol. Paired-end sequencing reads of 150 base pairs
116 were obtained with the Miseq (Illumina, San Diego, USA) platform. Illumina sequencer output
117 files matching of up to 151-base pair sequencing reads were processed with the Galaxy platform
118 after filtering and checking read quality (FastQC algorithm). Trinity software was used for *de*
119 *novo* assembly. The constructed sequences were checked with the Blast option from the
120 National Center for Biotechnology Information (NCBI). Reads were mapped with the Bowtie2
121 tool onto the best sequence given by the *de novo* assembly sequences. Nucleotide analysis was
122 performed position-by-position using the mpileup program. Positions with a sequencing depth
123 of over 100X were retained for further analysis. Only variants representing 10% or more of the
124 HBV quasispecies were considered for analysis. For each patient, consensus sequences of the
125 whole HBV genome were compared between samples collected before treatment and at
126 treatment failure. We searched for known HBs Ag escape mutations (Colagrossi et al., 2018;
127 Coppola, 2015; Lazarevic, 2014; Weber, 2005), core promoter and pre-core mutations (Parekh
128 et al., 2003) in these sequences, after alignment with other sequences from the same genotype
129 and reference sequences: HBV/E (AY738145), HBV/D (AF121240) and HBV/I (AB231908)
130 extracted from HBV database (HBVdb) (Hayer et al., 2013). HBV strain X02763 from HBVdb
131 (Hayer et al., 2013), with the usual numbering, was used as the reference sequence for the
132 annotation of mutations. Mutations were identified on the basis of their position in the pre-S1,

HBe, HBx and polymerase proteins or their position in the HBs Ag (s) or reverse transcriptase domain (rt), if relevant. Core promoter mutations (CPM) were annotated with their nucleic acid position in the X02763 HBV genome.

Phenotypic drug resistance testing

We had 1.1 units of the HBV genome (pCI_HBV1.1x), corresponding to a consensus sequence identified by NGS on isolates collected before treatment, synthesized *in vitro* (by GenScript Biotech, the Netherlands) and inserted between MluI and SalI restriction sites of a pCI vector (Promega, Toulon, France), downstream from a CMV promoter. This construct allowed HBV replication after transfection in HuH7 cells. Relevant mutations selected during entecavir treatment were introduced into these clones by site-directed mutagenesis (GenScript Biotech).

Phenotypic resistance assays were performed as previously described (Marlet et al., 2020). Briefly, Huh7 cells were transfected with pCI_HBV1.1x. Entecavir monohydrate (Merck, Missouri, USA) was added to the cells at six different final concentrations, ranging from 2.5 to 50 000 nM. After four days of culture, intracapsid viral DNA was quantified by duplex real-time PCR. We determined 50% inhibitory concentrations (IC₅₀) by least squares regression in GraphPad Prism v5.0 (GraphPad Software, California, USA). The fold-change in resistance corresponded to the IC₅₀ of entecavir for a given isolate, relative to the IC₅₀ of entecavir for the wild-type HBV genotype D *ayw3* strain (pCI_HBVpg1820), which was tested in parallel. Entecavir IC₅₀ and the fold-change in resistance were determined in three independent experiments and compared in Mann-Whitney tests. Replication capacity relative to the same wild-type strain as before was determined by quantifying HBV DNA levels in six different wells after four days of culture without antiviral drugs. Results are represented as the means and standard errors.

156

157 **Data availability**

158 All sequences were deposited in the GenBank database (Accession numbers: [SAMN16263252-](#)
159 [SAMN16263257](#) for MiSeq data and MW082636-MW082641 for consensus sequences).

160 **RESULTS**

161 **Patients**

162 During the study period, HBV genotypic drug resistance testing was performed on 110 clinical
163 isolates. We identified seven isolates associated with entecavir treatment failure but with no
164 detectable Pol/RT resistance mutations. Four isolates were not included because of poor patient
165 compliance with treatment ($n=3$) or because viral DNA load was too low for amplification by
166 PCR ($n=1$, 1.3 log IU/mL). The three remaining isolates (P1, P2 and P3) were associated with
167 a partial virological response (PVR) to entecavir in immunocompromised patients (Figure 1).

168 Patient #1 underwent liver transplantation in the context of decompensated alcohol-related
169 cirrhosis. He was negative for HBs Ag, HBs Ab and HBc Ab before transplantation, but he
170 received a graft from a donor with positive HBc Ab. He received one dose of anti-HBs
171 immunoglobulins (IVHEBEX, 10 000 IU, LFB Biomédicaments) during the anhepatic phase,
172 to prevent HBV infection from the graft. Despite this prophylaxis, he experienced HBV
173 infection leading to acute hepatitis one year later, whilst on mycophenolate mofetil (MMF) and
174 tacrolimus treatment.

175 Patient #2 underwent kidney transplantation and was treated with MMF, sirolimus and
176 prednisolone. She was anti-HBc Ab-positive and HBs Ag-negative before transplantation. She

177 was diagnosed with asymptomatic HBV infection six years later (she had tested negative for
178 HBs Ag the year before).

179 Patient #3 had a history of resolved HBV infection (HBc Ab-positive, HBs Ag-negative) before
180 he was diagnosed with chronic lymphocytic leukemia (CLL). He was treated with one dose of
181 rituximab, and with cyclophosphamide and fludarabine for two years before remission of CLL
182 was achieved and treatment stopped. HBV reactivation was diagnosed six years later, in the
183 context of a relapse of CLL with cytotoxicity. Rituximab is known to be associated with a high
184 risk of HBV reactivation and could have triggered HBV reactivation in this patient (Loomba
185 and Liang, 2017). Interestingly, this patient remained negative for HBc Ab throughout follow-
186 up, probably due to the strong B-cell depletion induced by rituximab (Feeney et al., 2013).
187 Patients #1, #2 and #3 reported good treatment compliance, which was confirmed by
188 therapeutic drug monitoring (entecavir plasma concentrations = 0.45, 0.78 and 0.45 ng/mL,
189 respectively). No relevant drug interactions between HBV treatment and concomitant
190 medication were detected. Retrospective genotypic drug resistance testing identified the same
191 Pol/RT sequences in pretreatment (P1_D0, P2_D0 and P3_D0) and post-treatment (P1, P2 and
192 P3) isolates (GenBank MW082636-MW082641). These findings suggest that the PVR to
193 entecavir in these three patients was not associated with the selection of resistance mutations in
194 the Pol/RT domain, or with poor patient compliance.

Figure 1: Biological and clinical follow-up of patients experiencing a partial virological response to entecavir

HBV viral loads are represented from the diagnosis of HBV infection (month 0) to the end of follow-up. The clinical isolates included in this study are identified with red diamonds and characters in bold typeface. ETV, entecavir; TNF, tenofovir disoproxil fumarate; MMF, mycophenolate mofetil. HBV viral loads were quantified with the Abbott HBV real-time assay. Its lower limit of detection (10 IU/mL) is represented with a dashed line. Patients #2 and #3 died from graft rejection and acute complications of hemopathy, respectively.

204 Table 1: Clinical and virological profiles of patients before entecavir treatment

Patient	Sex	Age (years)	Immunosuppression		HBV genotype	Serotype	Metavir score	ALAT (IU/L)	HBV DNA ^a (log IU/mL)	HBs Ag ^b	HBs Ab ^b	HBe Ag ^b
			Context	Delay before HBV infection (years)								
#1	M	41	Liver transplant	1	E	<i>ayw4</i>	A2F2	400	7.4	3760	0	924
#2	F	75	Kidney transplant	6	I ^c	<i>ayw^c</i>	A0F2	26	6.6	2051	5	2094
#3	M	73	Chronic lymphocytic leukemia	17	D	<i>ayw3</i>	F2	57	8.3	320	0	1622

205 ^a HBV real-time assay (Abbott); ^b Architect HBs Ag qualitative, HBs Ab and HBe Ag assays (Abbott); ^c This rare genotype originates from
206 a recombination between genotypes A, C and G, and circulates in Asian countries, including Laos, the patient's home country (Hannoun et
207 al., 2000; Huy et al., 2008). The subtype of this isolate remained undetermined, because aa s127S was not included in the current
208 classification (Kay and Zoulim, 2007; Purdy et al., 2007).

209 **Mutations detected before entecavir treatment**

210 Clinically relevant mutations (HBs immune escape or core promoter mutations, CPM) were detected
211 in the consensus sequence of the three HBV isolates collected before entecavir treatment (Figure 2).
212 Isolate P1_D0 harbored core promoter mutations (CPM) T1753G, A1762T and G1764A (Figure 2).
213 The last two of these mutations are known to be associated with poor HBe Ag secretion and an
214 increased HBV replication capacity, with no impact on susceptibility to entecavir (Leng et al., 2015;
215 Parekh et al., 2003). Interestingly, this unique combination of three CPMs has been described once
216 before, in a patient experiencing HBV reactivation following liver transplantation (Kim et al., 2003).
217 Mutation T1753G has not been previously characterized, but mutation T1753C increased the level of
218 replication associated with mutations A1762T and G1764A (Parekh et al., 2003).
219 Isolates P2_D0 and P3_D0 harbored HBs Ag immune escape mutations, which are frequently
220 detected in this context of HBV reactivation (Salpini et al., 2015). Isolate P2_D0 harbored a unique
221 set of mutations in the “a” determinant of the HBs Ag: sK122R, sT126I, sP127S and sG145R. The
222 sK122R, sT126I and sG145R mutations have been associated with immune escape and the restoration
223 of viral fitness in lamivudine-resistant isolates (Amini-Bavil-Olyaei et al., 2010; Bock et al., 2002).
224 The sP127S mutation has been observed only once before, in an isolate from Laos (Olinger et al.,
225 n.d.) but another mutation at this position (sP127L) has been described as an immune escape mutation
226 (Colagrossi et al., 2018). This suggests that the sP127S mutation may also be an immune escape
227 mutation. Isolate P3_D0 harbored the sM133I mutation of the “a” determinant of the HBs Ag MHR
228 region. This mutation has been associated with HBV reactivation (Pei et al., 2014), mother-to-child
229 HBV transmission despite vaccination (Latthaphasavang et al., 2019) and major antigenic changes in
230 the HBs Ag (Ie et al., 2010).

Figure 2: Mutations detected before entecavir treatment

Mutations of the consensus sequence detected in HBV isolates collected before entecavir treatment. Clinically relevant mutations (HBs Ag escape mutations and core promoter mutations CPM T1753G, A1762T and G1764A) are represented with orange symbols and labeled in bold typeface, whereas mutations of unknown significance are shown in white. Mutations rarely (<3%) found in other isolates of the same genotype are represented with a triangle, whereas more frequent mutations are represented with a diamond. Mutations associated with a change in two ORFs are indicated with a dashed line between the two ORFs. Shades of gray correspond to the pre-C, Core, pre-S1, pre-S2, S and X genes (upper row) and the terminal protein, spacer, polymerase/reverse transcriptase and RNase domains of the polymerase gene (lower row). MHR, major hydrophilic region. HBV strain X02763, with the usual numbering, was used as the reference sequence for annotation.

Mutations selected during entecavir treatment

We analyzed the evolution of HBV quasispecies by studying the whole HBV genome by NGS and comparing isolates collected before (P1_D0, P2_D0 and P3_D0) and after (P1, P2 and P3) entecavir treatment failure. Twenty-five non-synonymous mutations had been selected in isolates P1 ($n=9$), P2 ($n=2$) and P3 ($n=14$) (Figure 3). Two of these mutations, both in isolate P1, were considered to be major variants. Their proportion in the HBV quasispecies increased from 0-6 % before treatment to 41-57 % during treatment (Figure 3). The first mutation resulted in simultaneous changes in the spacer domain of the polymerase (A300E) and in the pre-S1 protein (Q120K) (Figure 3). The latter mutation was localized at the N-terminus of pre-S2 (Q2K) in a T-cell epitope (Barnaba et al., 1989; Pajot et al., 2006). This mutation has been observed in 1.2% of HBV/E isolates (Abdou Chekaraou et al., 2010; Candotti et al., 2016; Mulders et al., 2004). The second mutation, in the pre-core (Q206K) and core (Q177K) proteins, was located at the C-terminal end of the core protein (Figure 3), in a highly conserved protamine-like DNA-binding domain crucial for HBV pgRNA encapsidation and DNA replication (Jung et al., 2012). Interestingly, this is the first description of a mutation at this position. These two mutations were not known to be associated with resistance to entecavir. Phenotypic assays were performed on recombinant viruses harboring one or both of these mutations, to determine whether they had any impact on the level of resistance to entecavir.

The other 23 mutations were selected during entecavir treatment, but only as minor variants (frequency <40% at treatment failure), suggesting that they were probably not required for a PVR to entecavir. Eight of these mutations were located in the sequence corresponding to the Pol/RT domain, all as minor variants (<25%). None was detected with Sanger sequencing, coherent with their distribution as minor variants. Three mutations were of clinical interest. The

267 first was pre-core mutation W28*, selected in patient #1. The other two were basal core
268 promoter mutations A1762T and G1764A, selected in patient #3 (Figure 3). These pre-
269 core/basal core promoter mutations are known to be associated with increases in viral
270 replication and decreases in HBe Ag expression (Parekh et al., 2003). However, as minor
271 variants, they were considered unlikely to account for entecavir treatment failure and were not
272 further characterized.

273

Figure 3: HBV mutations selected or repressed during entecavir treatment

Mutations selected or repressed during entecavir treatment are indicated with pie charts representing their proportion in the HBV quasispecies, together with their position in the HBV genome. Clinically relevant mutations are represented with orange symbols and in bold typeface: core promoter mutations (CPM) A1762T and G1764A and pre-core mutation W28*. Mutations rarely (<3%) found in other isolates of the same genotype are represented with a triangle, whereas more frequent mutations are represented with a diamond. Mutations

associated with a change in two ORFs are indicated with a dashed line between the two ORFs. Shades of gray correspond to the pre-C, Core, pre-S1, pre-S2, S and X genes (upper row) and the terminal protein, spacer, polymerase/reverse transcriptase and RNase domains of the polymerase gene (lower row). MHR, major hydrophilic region. HBV strain X02763, with the usual numbering, was used as the reference sequence for annotation.

Phenotypic characterization of mutations selected during entecavir treatment

Recombinant viruses harboring the whole genome of isolate P1_D0, with or without the two relevant mutations selected during entecavir treatment in patient #1, were obtained by *in vitro* synthesis and site-directed mutagenesis, followed by transfection of the Huh7 cell line. The level of resistance to entecavir and the replication capacity of these recombinant viruses are shown in Figure 4. Variants harboring one or both of these mutations were as susceptible to entecavir and had a similar replicative capacity to the pretreatment isolate P1_D0 (Figure 4). This suggests that these mutations, alone or in combination, played no direct role in entecavir treatment failure.

298

299 **Figure 4: Phenotypic resistance testing of HBV isolates selected during entecavir**
300 **treatment in patient #1.**

301 Major variant before treatment (P1_D0, plain bar) and variants detected at treatment failure
302 (hatched bars) harboring the A300E mutation in the polymerase and the Q120K mutation in
303 the pre-S1/S2 proteins (P1_PS*), the Q206K mutation in the Core protein (P1_C*) or both
304 (P1_PS*C*). NS, non-significant ($p > 0.05$). Six drug concentrations, ranging from 2.5 to 50
305 000 nM, were tested in triplicate, in three independent experiments.

306

DISCUSSION

In a series of 110 patients tested for HBV drug resistance, we identified three patients experiencing entecavir treatment failure despite having good treatment compliance and no detectable Pol/RT resistance mutations. These three patients experienced a PVR to entecavir in a context of immunosuppression.

Previous studies demonstrated that PVR to entecavir does not require selection of resistance mutations in the Pol/RT domain but is associated with HBe Ag positivity, low levels of ALAT and a high HBV DNA viral load at baseline (Chen et al., 2014; Lee et al., 2015; Li et al., 2015; Zoutendijk et al., 2011). Despite these findings, most cases of PVR to entecavir remain unexplained. We hypothesized that mutations selected outside of the Pol/RT domain could contribute to PVR to entecavir. Mutations Q206K (pre-core/core), Q120K (pre-S1/pre-S2, in a T-cell epitope) and A300E (spacer domain) were selected during entecavir treatment in patient #1 but were not associated with an increased level of resistance to entecavir or an increase in HBV replication capacity. No mutations were selected in the whole HBV genome of patients #2 and #3. This study provided a unique insight into the evolution of HBV quasispecies during entecavir treatment, addressing the whole HBV genome and not only the Pol/RT domain. We demonstrated that PVR to entecavir does not require selection of any resistance mutation in the whole HBV genome. Our results demonstrate that mutations can be selected outside of the Pol/RT domain during entecavir treatment. These mutations could contribute to entecavir treatment failure by other mechanisms than an increased level of resistance.

Mutations of unknown significance can be selected in the whole HBV genome during entecavir treatment. In this study, mutation Q120K in the pre-S1/pre-S2 protein was selected during entecavir treatment in patient #1. This mutation was localized in a T-cell epitope (pre-S2 1-15)

(Barnaba et al., 1989; Pajot et al., 2006) and in a domain required for viral secretion (pre-S2 1-5) (Le Seyec et al., 1998). Mutations and deletions at this position tend to be selected in asymptomatic carriers and are associated with reduced secretion of HBs Ag *in vitro* and *in vivo* (Fan et al., 2001; Taghiabadi et al., 2019; Wang et al., 2018). This mutation could have been associated with viral escape from cellular immune responses and could have contributed to the partial virological response observed in patient #1. In contrast, in patients #2 and #3, no mutations were selected in the whole HBV genome during treatment, suggesting that PVR to entecavir did not rely on viral escape or resistance. HBs Ag immune escape mutations were selected before treatment in these two patients, as previously observed in the context of HBV reactivation (Corti et al., 2018; Rehmann and Nascimbeni, 2005).

It is unclear whether treatment with immunosuppressive drugs, such as rituximab, is associated with PVR to entecavir. These drugs have been associated with an increased risk of HBV reactivation (Loomba and Liang, 2017) and delayed response to entecavir treatment in a minority of patients (Brost et al., 2010; Yap et al., 2014). In addition, low ALAT levels at baseline have been associated with an increase in the risk of PVR to entecavir. It has been hypothesized by others that these low ALAT levels were the result of an impaired host immune response, which could increase the risk of PVR to entecavir (Lee et al., 2015; Li et al., 2015). Our findings support this hypothesis and suggest that mutations selected outside of the Pol/RT domain could contribute to entecavir treatment failure, not by an increased level of resistance, but by an immune escape mechanism, in synergy with immunosuppressive drugs.

We developed an original approach to study entecavir treatment failure, based on whole HBV genome analysis, combining deep sequencing and a phenotypic resistance assay. This made it possible to perform a comprehensive evaluation of mutations selected before or during entecavir treatment, potentially associated with treatment failure. The main limitation of this

study is the small sample size, which made it impossible to draw reliable conclusions on the association between immune escape mutations and PVR to entecavir.

This study argues for a systematic approach to entecavir treatment failure. In addition to Pol/RT sequencing, therapeutic drug monitoring seems to be useful for detecting poor compliance with treatment or underdosing in some patients. Screening for HBs immune escape mutations may be relevant in immunocompromised patients experiencing HBV reactivation, to confirm the association between these mutations and the PVR to entecavir and to guide treatment adaptation more effectively. By contrast, deep sequencing does not seem more relevant than Sanger sequencing for detecting resistance mutations at treatment failure, as they can be detected as major variants in this context (Lee et al., 2018; Lowe et al., 2016). However, in the near future, HBV treatment strategies will target other steps of the viral cycle and/or aim to modulate the host immune response (Cornberg et al., 2020; Fanning et al., 2019). In this context, deep sequencing and phenotypic characterization of whole HBV genomes from sequential clinical isolates, as performed in this study, will probably be essential, for the identification of viral factors associated with the response to these new antiviral treatments.

ACKNOWLEDGMENTS

We thank the Thellie endowment fund and our generous donors for funding this project. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

REFERENCES

- Abdou Chekaraou, M., Brichler, S., Mansour, W., Le Gal, F., Garba, A., Dény, P., Gordien, E., 2010. A novel hepatitis B virus (HBV) subgenotype D (D8) strain, resulting from recombination between genotypes D and E, is circulating in Niger along with HBV/E strains. *J. Gen. Virol.* 91, 1609–1620. <https://doi.org/10.1099/vir.0.018127-0>
- Amini-Bavil-Olyaei, S., Vucur, M., Luedde, T., Trautwein, C., Tacke, F., 2010. Differential impact of immune escape mutations G145R and P120T on the replication of lamivudine-resistant hepatitis B virus e antigen-positive and -negative strains. *J. Virol.* 84, 1026–1033. <https://doi.org/10.1128/JVI.01796-09>
- Barnaba, V., Franco, A., Alberti, A., Balsano, C., Benvenuto, R., Balsano, F., 1989. Recognition of hepatitis B virus envelope proteins by liver-infiltrating T lymphocytes in chronic HBV infection. *J. Immunol. Baltim. Md* 143, 2650–2655.
- Bock, C., Thomas, Tillmann, H.L., Torresi, J., Klempnauer, J., Locarnini, S., Manns, M.P., Trautwein, C., 2002. Selection of hepatitis B virus polymerase mutants with enhanced replication by lamivudine treatment after liver transplantation. *Gastroenterology* 122, 264–273. <https://doi.org/10.1053/gast.2002.31015>
- Brost, S., Schnitzler, P., Stremmel, W., Eisenbach, C., 2010. Entecavir as treatment for reactivation of hepatitis B in immunosuppressed patients. *World J. Gastroenterol.* 16, 5447–5451. <https://doi.org/10.3748/wjg.v16.i43.5447>
- Candotti, D., Diarra, B., Bisseye, C., Tao, I., Pham Quang, K., Sanou, M., Laperche, S., Sanogo, R., Allain, J.-P., Simporé, J., 2016. Molecular characterization of hepatitis B virus in blood donors from Burkina Faso: Prevalence of quasi-subgenotype A3, genotype E, and mixed infections. *J. Med. Virol.* 88, 2145–2156. <https://doi.org/10.1002/jmv.24589>
- Chang, T.-T., Gish, R.G., de Man, R., Gadano, A., Sollano, J., Chao, Y.-C., Lok, A.S., Han, K.-H., Goodman, Z., Zhu, J., Cross, A., DeHertogh, D., Wilber, R., Colonno, R., Apelian, D., BEHoLD A1463022 Study Group, 2006. A comparison of entecavir and lamivudine for HBeAg-positive chronic hepatitis B. *N. Engl. J. Med.* 354, 1001–1010. <https://doi.org/10.1056/NEJMoa051285>
- Chen, C.-H., Hu, T.-H., Hung, C.-H., Wang, J.-H., Lu, S.-N., Lee, C.-M., 2014. Antiviral effect of entecavir in nucleos(t)ide analogue-naïve and nucleos(t)ide analogue-experienced chronic hepatitis B patients without virological response at week 24 or 48 of therapy. *J. Viral Hepat.* 21, e55–e64. <https://doi.org/10.1111/jvh.12239>
- Chevaliez, S., Rodriguez, C., Poiteau, L., Soulier, A., Donati, F., Darty-Mercier, M., Pioche, C., Leroy, V., Brodard, V., Zoulim, F., Brouard, C., Larsen, C., Semaille, C., Roudot-Thoraval, F., Pawlotsky, J.-M., 2019. Primary resistance of hepatitis B virus to nucleoside and nucleotide analogues. *J. Viral Hepat.* 26, 278–286. <https://doi.org/10.1111/jvh.13025>
- Colagrossi, L., Hermans, L.E., Salpini, R., Di Carlo, D., Pas, S.D., Alvarez, M., Ben-Ari, Z., Boland, G., Bruzzzone, B., Coppola, N., Seguin-Devaux, C., Dyda, T., Garcia, F., Kaiser, R., Köse, S., Krarup, H., Lazarevic, I., Lunar, M.M., Maylin, S., Micheli, V., Mor, O., Paraschiv, S., Paraskevis, D., Poljak, M., Puchhammer-Stöckl, E., Simon, F., Stanojevic, M., Stene-Johansen, K., Tihic, N., Trimoulet, P., Verheyen, J., Vince, A., Lepej, S.Z., Weis, N., Yalcinkaya, T., Boucher, C.A.B., Wensing, A.M.J., Perno, C.F., Svisch, V., HEPVIR working group of the European Society for translational antiviral research (ESAR), 2018. Immune-escape mutations and stop-codons in HBsAg develop in a large proportion of patients with chronic HBV infection exposed to anti-HBV drugs in Europe. *BMC Infect. Dis.* 18, 251. <https://doi.org/10.1186/s12879-018-3161-2>
- Coppola, N., 2015. Clinical significance of hepatitis B surface antigen mutants. *World J. Hepatol.* 7, 2729. <https://doi.org/10.4254/wjh.v7.i27.2729>
- Cornberg, M., Lok, A.S.-F., Terrault, N.A., Zoulim, F., Berg, T., Brunetto, M.R., Buchholz, S., Buti, M., Chan, H.L.Y., Chang, K.-M., Dandri, M., Dusheiko, G., Feld, J.J., Ferrari, C., Ghany, M., Janssen, H.L.A., Kennedy, P., Lampertico, P., Liang, J., Locarnini, S., Maini, M.K., Mishra, P., Papatheodoridis, G., Petersen, J., Schlottmann, S., Wang, S., Wedemeyer, H., 2020. Guidance for design and endpoints of clinical trials in chronic hepatitis B - Report from the 2019 EASL-

- AASLD HBV Treatment Endpoints Conference. *J. Hepatol.* 72, 539–557.
<https://doi.org/10.1016/j.jhep.2019.11.003>
- Corti, D., Benigni, F., Shouval, D., 2018. Viral envelope-specific antibodies in chronic hepatitis B virus infection. *Curr. Opin. Virol.*, Antiviral strategies 30, 48–57.
<https://doi.org/10.1016/j.coviro.2018.04.002>
- Fan, Y.-F., Lu, C.-C., Chen, W.-C., Yao, W.-J., Wang, H.-C., Chang, T.-T., Lei, H.-Y., Shiau, A.-L., Su, I.-J., 2001. Prevalence and significance of hepatitis B virus (HBV) pre-S mutants in serum and liver at different replicative stages of chronic HBV infection. *Hepatology* 33, 277–286.
<https://doi.org/10.1053/jhep.2001.21163>
- Fanning, G.C., Zoulim, F., Hou, J., Bertoletti, A., 2019. Therapeutic strategies for hepatitis B virus infection: towards a cure. *Nat. Rev. Drug Discov.* 18, 827–844.
<https://doi.org/10.1038/s41573-019-0037-0>
- Feeney, S.A., McCaughey, C., Watt, A.P., Agnaf, M.R.E., McDougall, N., Wend, U.C., Gerlich, W.H., Coyle, P.V., 2013. Reactivation of occult hepatitis B virus infection following cytotoxic lymphoma therapy in an anti-HBc negative patient. *J. Med. Virol.* 85, 597–601.
<https://doi.org/10.1002/jmv.23513>
- Guinoiseau, T., Moreau, A., Hohnadel, G., Ngo-Giang-Huong, N., Brulard, C., Vourc'h, P., Goudeau, A., Gaudy-Graffin, C., 2017. Deep sequencing is an appropriate tool for the selection of unique Hepatitis C virus (HCV) variants after single genomic amplification. *PLOS ONE* 12, e0174852.
<https://doi.org/10.1371/journal.pone.0174852>
- Günther, S., Li, B.-C., Miska, S., Krüger, D.H., Meisel, H., Will, H., 1995. A novel method for efficient amplification of whole hepatitis B virus genomes permits rapid functional analysis and reveals deletion mutants in immunosuppressed patients. *J. Virol.* 69, 5437–5444.
- Hannoun, C., Norder, H., Lindh, M., 2000. An aberrant genotype revealed in recombinant hepatitis B virus strains from Vietnam The GenBank accession numbers of the sequences reported in this paper are AF241407–AF241411. *J. Gen. Virol.* 81, 2267–2272.
<https://doi.org/10.1099/0022-1317-81-9-2267>
- Hayashi, S., Murakami, S., Omagari, K., Matsui, T., Iio, E., Isogawa, M., Watanabe, T., Karino, Y., Tanaka, Y., 2015. Characterization of novel entecavir resistance mutations. *J. Hepatol.* 63, 546–553.
- Hayer, J., Jadeau, F., Deléage, G., Kay, A., Zoulim, F., Combet, C., 2013. HBVdb: a knowledge database for Hepatitis B Virus. *Nucleic Acids Res.* 41, D566–D570.
<https://doi.org/10.1093/nar/gks1022>
- Huang, C.-J., Wu, C.-F., Lan, C.-Y., Sung, F.-Y., Lin, C.-L., Liu, C.-J., Liu, H.-F., Yu, M.-W., 2013. Impact of Genetic Heterogeneity in Polymerase of Hepatitis B Virus on Dynamics of Viral Load and Hepatitis B Progression. *PLoS ONE* 8, e70169. <https://doi.org/10.1371/journal.pone.0070169>
- Huy, T.T.T., Ngoc, T.T., Abe, K., 2008. New Complex Recombinant Genotype of Hepatitis B Virus Identified in Vietnam. *J. Virol.* 82, 5657–5663. <https://doi.org/10.1128/JVI.02556-07>
- Ie, S.I., Thedja, M.D., Roni, M., Muljono, D.H., 2010. Prediction of conformational changes by single mutation in the hepatitis B virus surface antigen (HBsAg) identified in HBsAg-negative blood donors. *Virol. J.* 7, 326. <https://doi.org/10.1186/1743-422X-7-326>
- Jung, J., Kim, H.-Y., Kim, T., Shin, B.-H., Park, G.-S., Park, S., Chwae, Y.-J., Shin, H.-J., Kim, K., 2012. C-terminal substitution of HBV core proteins with those from DHBV reveals that arginine-rich 167RRRSQSPRR175 domain is critical for HBV replication. *PLoS One* 7, e41087.
<https://doi.org/10.1371/journal.pone.0041087>
- Kay, A., Zoulim, F., 2007. Hepatitis B virus genetic variability and evolution. *Virus Res.* 127, 164–176.
<https://doi.org/10.1016/j.virusres.2007.02.021>
- Kim, K.-H., Lee, K.-H., Chang, H.-Y., Ahn, S.H., Tong, S., Yoon, Y.J., Seong, B.L., Kim, S.I., Han, K.-H., 2003. Evolution of hepatitis B virus sequence from a liver transplant recipient with rapid breakthrough despite hepatitis B immune globulin prophylaxis and lamivudine therapy. *J. Med. Virol.* 71, 367–375. <https://doi.org/10.1002/jmv.10503>

479 Lai, C.-L., Shouval, D., Lok, A.S., Chang, T.-T., Cheinquer, H., Goodman, Z., DeHertogh, D., Wilber, R.,
 480 Zink, R.C., Cross, A., Colonna, R., Fernandes, L., BEHoLD A1463027 Study Group, 2006.
 481 Entecavir versus lamivudine for patients with HBeAg-negative chronic hepatitis B. *N. Engl. J.*
 482 *Med.* 354, 1011–1020. <https://doi.org/10.1056/NEJMoa051287>
 483 Lampertico, P., Agarwal, K., Berg, T., Buti, M., Janssen, H.L.A., Papatheodoridis, G., Zoulim, F., Tacke,
 484 F., 2017. EASL 2017 Clinical Practice Guidelines on the management of hepatitis B virus
 485 infection. *J. Hepatol.* 67, 370–398. <https://doi.org/10.1016/j.jhep.2017.03.021>
 486 Latthaphasavang, V., Vanhems, P., Ngo-Giang-Huong, N., Sibounlang, P., Paboriboune, P., Malato, L.,
 487 Keoluangkhot, V., Thammasack, S., Salvadori, N., Khamduang, W., Steeneste, N., Trépo, C.,
 488 Dény, P., Jourdain, G., 2019. Perinatal hepatitis B virus transmission in Lao PDR: A
 489 prospective cohort study. *PLoS ONE* 14. <https://doi.org/10.1371/journal.pone.0215011>
 490 Lazarevic, I., 2014. Clinical implications of hepatitis B virus mutations: Recent advances. *World J.*
 491 *Gastroenterol.* WJG 20, 7653–7664. <https://doi.org/10.3748/wjg.v20.i24.7653>
 492 Le Seyec, J., Chouteau, P., Cannie, I., Guguen-Guillouzo, C., Gripon, P., 1998. Role of the Pre-S2
 493 Domain of the Large Envelope Protein in Hepatitis B Virus Assembly and Infectivity. *J. Virol.*
 494 72, 5573–5578.
 495 Lee, G.-H., Inoue, M., Chong, R.H.-H., Toh, J., Wee, S.Y., Loh, K.-S., Lim, S.-G., 2018. Pyrosequencing
 496 method for sensitive detection of HBV drug resistance mutations. *J. Med. Virol.* 90, 1071–
 497 1079. <https://doi.org/10.1002/jmv.25066>
 498 Lee, H.W., Kwon, J.-C., Oh, I.S., Chang, H.Y., Cha, Y.J., Choi, I.-S., Kim, H.J., 2015. Prolonged Entecavir
 499 Therapy Is Not Effective for HBeAg Seroconversion in Treatment-Naïve Chronic Hepatitis B
 500 Patients with a Partial Virological Response. *Antimicrob. Agents Chemother.* 59, 5348–5356.
 501 <https://doi.org/10.1128/AAC.01017-15>
 502 Leng, X.-H., Chen, E.-Q., Du, L.-Y., Bai, L., Gong, D.-Y., Cheng, X., Huang, F.-J., Tang, H., 2015. Biological
 503 characteristics of the A1762T/G1764A mutant strain of hepatitis B virus in vivo. *Mol. Med.*
 504 *Rep.* 12, 5141–5148. <https://doi.org/10.3892/mmr.2015.4072>
 505 Li, M., Lv, T., Wu, S., Wei, W., Wu, X., Ou, X., Ma, H., Chow, S.-C., Kong, Y., You, H., Jia, J., 2020.
 506 Tenofovir versus entecavir in lowering the risk of hepatocellular carcinoma development in
 507 patients with chronic hepatitis B: a critical systematic review and meta-analysis. *Hepatol. Int.*
 508 14, 105–114. <https://doi.org/10.1007/s12072-019-10005-0>
 509 Li, X., Li, F., Zhang, Y., Kang, Y., Yu, J., Yang, F., Liu, H., Qin, Y., Huang, Y., Mao, R., Zhang, J., 2015.
 510 Partial virological response to entecavir treatment in nucleos(t)ide-naïve patients with
 511 HBeAg-positive chronic hepatitis B is not caused by reduced sensitivity. *Biochem. Biophys.*
 512 *Res. Commun.* 464, 1185–1191. <https://doi.org/10.1016/j.bbrc.2015.07.101>
 513 Liu, L., Liu, Y., Chen, R., Li, X., Luo, D., Zhao, Y., Li, Q., Huang, B., Wang, F.-S., Liu, X., Xu, D., 2019.
 514 Prevalence of the entecavir-resistance-inducing mutation rtA186T in a large cohort of
 515 Chinese hepatitis B virus patients. *Antiviral Res.*
 516 <https://doi.org/10.1016/j.antiviral.2019.02.012>
 517 Liu, Y., Zhou, Y., Li, X., Niu, M., Chen, R., Shao, J., Si, L., Luo, D., Lin, Y., Li, L., Zhang, K., Xiao, X., Xu, Z.,
 518 Liu, M., Lu, M., Zoulim, F., Xu, D., 2019. Hepatitis B virus mutation pattern
 519 rtL180M+A181C+M204V may contribute to entecavir resistance in clinical practice. *Emerg.*
 520 *Microbes Infect.* 8, 354–365. <https://doi.org/10.1080/22221751.2019.1584018>
 521 Lok, A.S., Zoulim, F., Locarnini, S., Bartholomeusz, A., Ghany, M.G., Pawlotsky, J.-M., Liaw, Y.-F.,
 522 Mizokami, M., Kuiken, C., Hepatitis B Virus Drug Resistance Working Group, 2007. Antiviral
 523 drug-resistant HBV: Standardization of nomenclature and assays and recommendations for
 524 management. *Hepatology* 46, 254–265. <https://doi.org/10.1002/hep.21698>
 525 Loomba, R., Liang, T.J., 2017. Hepatitis B Reactivation Associated With Immune Suppressive and
 526 Biological Modifier Therapies: Current Concepts, Management Strategies, and Future
 527 Directions. *Gastroenterology* 152, 1297–1309. <https://doi.org/10.1053/j.gastro.2017.02.009>
 528 Lowe, C.F., Merrick, L., Harrigan, P.R., Mazzulli, T., Sherlock, C.H., Ritchie, G., 2016. Implementation
 529 of Next-Generation Sequencing for Hepatitis B Virus Resistance Testing and Genotyping in a

Clinical Microbiology Laboratory. *J. Clin. Microbiol.* 54, 127–133.
<https://doi.org/10.1128/JCM.02229-15>

Marlet, J., Lier, C., Roch, E., Maugey, M., Moreau, A., Combe, B., Lefeuvre, S., d'Alteroche, L., Barbereau, D., Causse, X., Bastides, F., Bachelier, M.-N., Brand, D., Gaudy-Graffin, C., 2020. Revisiting HBV resistance to entecavir with a phenotypic approach. *Antiviral Res.* 104869. <https://doi.org/10.1016/j.antiviral.2020.104869>

Mulders, M.N., Venard, V., Njayou, M., Edorh, A.P., Bola Oyefolu, A.O., Kehinde, M.O., Muyembe Tamfum, J.-J., Nebie, Y.K., Maiga, I., Ammerlaan, W., Fack, F., Omilabu, S.A., Le Faou, A., Muller, C.P., 2004. Low genetic diversity despite hyperendemicity of hepatitis B virus genotype E throughout West Africa. *J. Infect. Dis.* 190, 400–408. <https://doi.org/10.1086/421502>

Olinger, C.M., Jutavijittum, P., Hübschen, J.M., Yousukh, A., Samountry, B., Thammavong, T., Toriyama, K., Muller, C.P., n.d. Possible New Hepatitis B Virus Genotype, Southeast Asia - Volume 14, Number 11—November 2008 - *Emerging Infectious Diseases journal* - CDC. <https://doi.org/10.3201/eid1411.080437>

Pajot, A., Michel, M.-L., Mancini-Bourgine, M., Ungeheuer, M.-N., Ojcius, D.M., Deng, Q., Lemonnier, F.A., Lone, Y.-C., 2006. Identification of novel HLA-DR1-restricted epitopes from the hepatitis B virus envelope protein in mice expressing HLA-DR1 and vaccinated human subjects. *Microbes Infect.* 8, 2783–2790. <https://doi.org/10.1016/j.micinf.2006.08.009>

Papatheodoridis, G.V., Sypsa, V., Dalekos, G.N., Yurdaydin, C., Van Boemmel, F., Buti, M., Calleja, J.L., Chi, H., Goulis, J., Manolakopoulos, S., Loglio, A., Voulgaris, T., Gatselis, N., Keskin, O., Veelken, R., Lopez-Gomez, M., Hansen, B.E., Savvidou, S., Kourikou, A., Vlachogiannakos, J., Galanis, K., Idilman, R., Esteban, R., Janssen, H.L.A., Berg, T., Lampertico, P., 2020. Hepatocellular carcinoma prediction beyond year 5 of oral therapy in a large cohort of Caucasian patients with chronic hepatitis B. *J. Hepatol.* <https://doi.org/10.1016/j.jhep.2020.01.007>

Parekh, S., Zoulim, F., Ahn, S.H., Tsai, A., Li, J., Kawai, S., Khan, N., Trepo, C., Wands, J., Tong, S., 2003. Genome Replication, Virion Secretion, and e Antigen Expression of Naturally Occurring Hepatitis B Virus Core Promoter Mutants. *J. Virol.* 77, 6601–6612. <https://doi.org/10.1128/JVI.77.12.6601-6612.2003>

Pei, R., Grund, S., Verheyen, J., Esser, S., Chen, X., Lu, M., 2014. Spontaneous reactivation of hepatitis B virus replication in an HIV coinfecting patient with isolated anti-Hepatitis B core antibodies. *Virol. J.* 11, 9. <https://doi.org/10.1186/1743-422X-11-9>

Peng, Y., Li, Yutang, Hou, J., Sun, J., Su, M., Li, Yao, Xiang, K., Yan, L., Zhuang, H., Li, T., 2017. The nucleotide changes within HBV core promoter/precore during the first 12 weeks of nucleos(t)ide treatment might be associated with a better virological response. *Infect. Genet. Evol.* 49, 116–121. <https://doi.org/10.1016/j.meegid.2017.01.009>

Podlaha, O., Gane, E., Brunetto, M., Fung, S., Chuang, W.-L., Pan, C.Q., Jiang, Z., Liu, Y., Bhardwaj, N., Mukherjee, P., Flaherty, J., Gaggar, A., Subramanian, M., Izumi, N., Shalimar, Lim, Y.-S., Marcellin, P., Buti, M., Chan, H.L.Y., Agarwal, K., 2019. Large-scale viral genome analysis identifies novel clinical associations between hepatitis B virus and chronically infected patients. *Sci. Rep.* 9, 1–9. <https://doi.org/10.1038/s41598-019-46609-7>

Purdy, M.A., Talekar, G., Swenson, P., Araujo, A., Fields, H., 2007. A New Algorithm for Deduction of Hepatitis B Surface Antigen Subtype Determinants from the Amino Acid Sequence. *Intervirology* 50, 45–51. <https://doi.org/10.1159/000096312>

Razavi-Shearer, D., Gamkrelidze, I., Nguyen, M.H., Chen, D.-S., Damme, P.V., Abbas, Z., Abdulla, M., Rached, A.A., Adda, D., Aho, I., Akarca, U., Hasan, F., Lawati, F.A., Naamani, K.A., Al-Ashgar, H.I., Alavian, S.M., Alawadhi, S., Albillos, A., Al-Busafi, S.A., Aleman, S., Alfaleh, F.Z., Aljumah, A.A., Anand, A.C., Anh, N.T., Arends, J.E., Arkkila, P., Athanasakis, K., Bane, A., Ben-Ari, Z., Berg, T., Bizri, A.R., Blach, S., Mello, C.E.B., Brandon, S.M., Bright, B., Bruggmann, P., Brunetto, M., Buti, M., Chan, H.L.Y., Chaudhry, A., Chien, R.-N., Choi, M.S., Christensen, P.B.,

Chuang, W.-L., Chulanov, V., Clausen, M.R., Colombo, M., Cornberg, M., Cowie, B., Craxi, A., Croes, E.A., Cuellar, D.A., Cunningham, C., Desalegn, H., Drazilova, S., Duberg, A.-S., Egeonu, S.S., El-Sayed, M.H., Estes, C., Falconer, K., Ferraz, M.L.G., Ferreira, P.R., Flisiak, R., Frankova, S., Gaeta, G.B., García-Samaniego, J., Genov, J., Gerstoft, J., Goldis, A., Gountas, I., Gray, R., Pessôa, M.G., Hajarizadeh, B., Hatzakis, A., Hézode, C., Himatt, S.M., Hoepelman, A., Hrstic, I., Hui, Y.-T.T., Husa, P., Jahis, R., Janjua, N.Z., Jarčuška, P., Jaroszewicz, J., Kaymakoglu, S., Kershenovich, D., Kondili, L.A., Konysbekova, A., Krajden, M., Kristian, P., Laleman, W., Lao, W.C., Layden, J., Lazarus, J.V., Lee, M.-H., Liakina, V., Lim, Y.-S.S., Loo, C.K., Lukšić, B., Malekzadeh, R., Malu, A.O., Mamatkulov, A., Manns, M., Marinho, R.T., Maticic, M., Mauss, S., Memon, M.S., Correa, M.C.M., Mendez-Sanchez, N., Merat, S., Metwally, A.M., Mohamed, R., Mokhbat, J.E., Moreno, C., Mossong, J., Mourad, F.H., Müllhaupt, B., Murphy, K., Musabaev, E., Nawaz, A., Nde, H.M., Negro, F., Nersesov, A., Nguyen, V.T.T., Njouom, R., Ntagirabiri, R., Nurmatov, Z., Obekpa, S., Ocamo, P., Oguiche, S., Omede, O., Omuemu, C., Opore-Sem, O., Opio, C.K., Örmeci, N., Papatheodoridis, G., Pasini, K., Pimenov, N., Poustchi, H., Quang, T.D., Qureshi, H., Ramji, A., Razavi-Shearer, K., Redae, B., Reesink, H.W., Rios, C.Y., Rjaskova, G., Robbins, S., Roberts, L.R., Roberts, S.K., Ryder, S.D., Safadi, R., Sagalova, O., Salupere, R., Sanai, F.M., Sanchez-Avila, J.F., Saraswat, V., Sarrazin, C., Schmelzer, J.D., Schröter, I., Scott, J., Seguin-Devaux, C., Shah, S.R., Sharara, A.I., Sharma, M., Shiha, G.E., Shin, T., Sievert, W., Sperl, J., Stärkel, P., Stedman, C., Sypsa, V., Tacke, F., Tan, S.S., Tanaka, J., Tomasiewicz, K., Urbanek, P., Meer, A.J. van der, Vlierberghe, H.V., Vella, S., Vince, A., Waheed, Y., Waked, I., Walsh, N., Weis, N., Wong, V.W., Woodring, J., Yaghi, C., Yang, H.-I., Yang, C.-L., Yesmembetov, K., Yosry, A., Yuen, M.-F., Yusuf, M.A.M., Zeuzem, S., Razavi, H., 2018. Global prevalence, treatment, and prevention of hepatitis B virus infection in 2016: a modelling study. *Lancet Gastroenterol. Hepatol.* 3, 383–403. [https://doi.org/10.1016/S2468-1253\(18\)30056-6](https://doi.org/10.1016/S2468-1253(18)30056-6)

Rehermann, B., Nascimbeni, M., 2005. Immunology of hepatitis B virus and hepatitis C virus infection. *Nat. Rev. Immunol.* 5, 215–229. <https://doi.org/10.1038/nri1573>

Rose, R.E., Hernandez, D., Falk, P.J., Ericson, K., Zhou, N., Thiry, A., McPhee, F., 2018. Discovery of the Novel Entecavir-Resistant Hepatitis B Virus Reverse Transcriptase A181C Substitution From an Integrated Genotypic Analysis. *Hepatol. Commun.* 2, 1123–1135. <https://doi.org/10.1002/hep4.1231>

Salpini, R., Colagrossi, L., Bellocchi, M.C., Surdo, M., Becker, C., Alteri, C., Aragri, M., Ricciardi, A., Armenia, D., Pollicita, M., Di Santo, F., Carioti, L., Louzoun, Y., Mastroianni, C.M., Lichtner, M., Paoloni, M., Esposito, M., D’Amore, C., Marrone, A., Marignani, M., Sarrecchia, C., Sarmati, L., Andreoni, M., Angelico, M., Verheyen, J., Perno, C.-F., Svicher, V., 2015. Hepatitis B surface antigen genetic elements critical for immune escape correlate with hepatitis B virus reactivation upon immunosuppression. *Hepatol. Baltim. Md* 61, 823–833. <https://doi.org/10.1002/hep.27604>

Taghiabadi, M., Hosseini, S.Y., Gorzin, A.A., Taghavi, S.A., Monavari, S.H.R., Sarvari, J., 2019. Comparison of pre-S1/S2 variations of hepatitis B virus between asymptomatic carriers and cirrhotic/hepatocellular carcinoma-affected individuals. *Clin. Exp. Hepatol.* 5, 161–168. <https://doi.org/10.5114/ceh.2019.84781>

Tenney, D.J., Rose, R.E., Baldick, C.J., Pokornowski, K.A., Eggers, B.J., Fang, J., Wichroski, M.J., Xu, D., Yang, J., Wilber, R.B., Colonno, R.J., 2009. Long-term monitoring shows hepatitis B virus resistance to entecavir in nucleoside-naïve patients is rare through 5 years of therapy. *Hepatology* 49, 1503–1514. <https://doi.org/10.1002/hep.22841>

Villeneuve, J.-P., Durantel, D., Durantel, S., Westland, C., Xiong, S., Brosgart, C.L., Gibbs, C.S., Parvaz, P., Werle, B., Trépo, C., Zoulim, F., 2003. Selection of a hepatitis B virus strain resistant to adefovir in a liver transplantation patient. *J. Hepatol.* 39, 1085–1089.

- Wang, T., Dai, Y., Zhang, M., Cui, D., Xu, X., Sun, C., Cheng, J., 2018. Sequence analysis of the Pre-S gene in chronic asymptomatic HBV carriers with low-level HBsAg. *Int. J. Mol. Med.* 42, 2689–2699. <https://doi.org/10.3892/ijmm.2018.3831>
- Weber, B., 2005. Genetic variability of the S gene of hepatitis B virus: clinical and diagnostic impact. *J. Clin. Virol.* 32, 102–112. <https://doi.org/10.1016/j.jcv.2004.10.008>
- Xu, Y., Zhang, Y.-G., Wang, X., Qi, W.-Q., Qin, S.-Y., Liu, Z.-H., Jiao, J., Wang, J.-B., 2015. Long-term antiviral efficacy of entecavir and liver histology improvement in Chinese patients with hepatitis B virus-related cirrhosis. *World J. Gastroenterol. WJG* 21, 7869–7876. <https://doi.org/10.3748/wjg.v21.i25.7869>
- Yap, D.Y.H., Yung, S., Tang, C.S.O., Seto, W.K., Ma, M.K.M., Mok, M.M.Y., Kwan, L.P.Y., Chan, G.C.W., Choy, B.Y., Yuen, M.F., Chan, T.M., 2014. Entecavir treatment in kidney transplant recipients infected with hepatitis B. *Clin. Transplant.* 28, 1010–1015. <https://doi.org/10.1111/ctr.12410>
- Zoulim, F., Locarnini, S., 2013. Optimal management of chronic hepatitis B patients with treatment failure and antiviral drug resistance. *Liver Int.* 33, 116–124. <https://doi.org/10.1111/liv.12069>
- Zoutendijk, R., Reijnders, J.G.P., Brown, A., Zoulim, F., Mutimer, D., Deterding, K., Petersen, J., Hofmann, W.P., Buti, M., Santantonio, T., van Bömmel, F., Pradat, P., Oo, Y., Luetgehetmann, M., Berg, T., Hansen, B.E., Wedemeyer, H., Janssen, H.L.A., VIRGIL Surveillance Study Group, 2011. Entecavir treatment for chronic hepatitis B: adaptation is not needed for the majority of naïve patients with a partial virological response. *Hepatol. Baltim. Md* 54, 443–451. <https://doi.org/10.1002/hep.24406>