



HAL
open science

Crossed brainstem syndrome revealing bleeding brainstem cavernous malformation: an illustrative case

Nathan Beucler, Sébastien Boissonneau, Aurélia Ruf, Stéphane Fuentes,
Romain Carron, Henry Dufour

► To cite this version:

Nathan Beucler, Sébastien Boissonneau, Aurélia Ruf, Stéphane Fuentes, Romain Carron, et al.. Crossed brainstem syndrome revealing bleeding brainstem cavernous malformation: an illustrative case. BMC Neurology, 2021, 21 (1), pp.204. 10.1186/s12883-021-02223-7 . inserm-03310546

HAL Id: inserm-03310546

<https://inserm.hal.science/inserm-03310546>

Submitted on 30 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CASE REPORT

Open Access



Crossed brainstem syndrome revealing bleeding brainstem cavernous malformation: an illustrative case

Nathan Beucler^{1,2*} , Sébastien Boissonneau^{1,3}, Aurélia Ruf⁴, Stéphane Fuentes¹, Romain Carron^{3,5} and Henry Dufour^{1,6}

Abstract

Background: Since the nineteenth century, a great variety of crossed brainstem syndromes (CBS) have been described in the medical literature. A CBS typically combines ipsilateral cranial nerves deficits to contralateral long tracts involvement such as hemiparesis or hemianesthesia. Classical CBS seem in fact not to be so clear-cut entities with up to 20% of patients showing different or unnamed combinations of crossed symptoms. In terms of etiologies, acute brainstem infarction predominates but CBS secondary to hemorrhage, neoplasm, abscess, and demyelination have been described. The aim of this study was to assess the proportion of CBS caused by a bleeding episode arising from a brainstem cavernous malformation (BCM) reported in the literature.

Case presentation: We present the case of a typical Foville syndrome in a 65-year-old man that was caused by a pontine BCM with extralesional bleeding. Following the first bleeding episode, a conservative management was decided but the patient had eventually to be operated on soon after the second bleeding event.

Discussion: A literature review was conducted focusing on the five most common CBS (Benedikt, Weber, Foville, Millard-Gubler, Wallenberg) on Medline database from inception to 2020. According to the literature, hemorrhagic BCM account for approximately 7 % of CBS. Microsurgical excision may be indicated after the second bleeding episode but needs to be carefully weighted up against the risks of the surgical procedure and openly discussed with the patient.

Conclusions: In the setting of a CBS, neuroimaging work-up may not infrequently reveal a BCM requiring complex multidisciplinary team management including neurosurgical advice.

Keywords: Foville syndrome, Crossed brainstem syndrome, Intracranial hemorrhage, Brainstem cavernous malformation, Developmental venous anomaly

* Correspondence: nathan.beucler@neurochirurgie.fr

¹Department of Neurosurgery, Timone University Hospital, APHM, 264 rue Saint-Pierre, 13005 Marseille, France

²Ecole du Val-de-Grâce, French Military Health Service Academy, 1 place Alphonse Laveran, 75230 Paris Cedex 5, France

Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

The anatomy of the brainstem is notable for comprising the nuclei and fibers of cranial nerves III to XII, long motor and sensory tracts, and crucial vegetative structures for cardio-respiratory functions and wakefulness. As a consequence, the clinical manifestations of brainstem injury vary from focal symptoms such as cranial nerves deficits to signs of long tracts involvement with motor or sensory impairment, and even vegetative state or death. The most frequent etiology of brainstem damage appears to be ischemic stroke [1]. Less frequent causes include multiple sclerosis, brainstem gliomas, brainstem abscesses, and vascular malformations just to cite a few. Among vascular malformations, a brainstem cavernous malformation (BCM) consists of a mulberry-like assembly of thin-walled vascular sinusoids which growth is self-sustained by repeated intralesional micro-bleed episodes. Yet, BCM may also be responsible for symptomatic extralesional bleedings which can be life threatening. The aim of this report is to present an original case of a genuine crossed brainstem syndrome (CBS) that turned out to be the mode of revelation of a bleeding BCM, and to discuss its frequency and its management.

Case presentation

A 65-year-old man presented with a 10-day history of sudden onset binocular diplopia and gait disturbance; he

also complained of tinnitus. His medical history consisted in chronic glaucoma treated with latanoprost eye droplets. The patient was on daily acetylsalicylic acid for primary prevention of cardiovascular disease. His family medical history revealed an ischemic stroke in one of his sisters and an unexpected death during her sleep in another sister. He also reported a fifty pack-year smoking and admitted chronic alcohol intake. His general practitioner introduced candesartan 4 mg daily upon symptoms onset. Careful neurological examination revealed a left abducens nerve (CN VI) palsy, a left peripheral facial nerve (CN VII) palsy, and a contralateral face-sparing hemiparesis (Fig. 1, Video 1). Right-sided mild dysesthesiae were also reported. There was no other cranial nerve deficit, no other focal neurological deficit (FND). There was no headache, no fever, no meningismus. Lab tests did not reveal inflammatory reaction. Magnetic resonance imaging of the brain revealed a BCM located on the left side of the floor of the fourth ventricle with evidence of recent extralesional bleeding. There was no other cerebral cavernous malformation on gradient-echo sequences. The BCM was associated with a developmental venous anomaly (DVA) draining both sides of the cerebellum directly into the vein of Galen (Fig. 2). The co-existence of an ipsilateral deficit of CN VI and VII and a contralateral face-sparing hemiparesis was highly suggestive of the inferior medial pontine syndrome, also known as Foville syndrome. The patient was admitted to




Fig. 1 Clinical examination reveals (a) an abducens nerve palsy and (b) a peripheral facial nerve palsy on the left side, associated with (c,d) a contralateral face-sparing hemiparesis. This crossed brainstem syndrome involves the inferior medial pons and was originally described by Achille Louis Foville in 1859




Fig. 2 The cerebral MRI shows (a, white arrowhead, post-contrast T1-weighted sequence) a brainstem cavernous malformation (BCM) associated with (a,b,c, black arrowheads, post-contrast T1-weighted sequence) a bilateral cerebellar developmental venous anomaly (DVA) prevailing on the right side and draining into the vein of Galen. (b, black arrow, post-contrast T1-weighted sequence and D, black arrow, gradient-echo sequence) The BCM was responsible for a medial pontine hematoma

the neurosurgery department for close follow-up. Acetylsalicylic acid was stopped. Considering this first bleeding episode, the non-exophytic character of the pontine hemorrhage, and the mild degree of disability of the patient (Glasgow Outcome Scale [GOS] of 5), a conservative management was decided in the first place. The option of stereotactic radiosurgery was deemed unnecessary at the acute phase and in the setting of a first bleeding. Five months later, the patient was admitted for recurrence of the symptoms with a grade V House-Brackman peripheral facial palsy and complete abducens nerve palsy on the left side, associated with contralateral face-sparing paresthesia. The CT scan of the brain showed evidence of rebleeding. After 2 weeks of close monitoring in the intensive care unit, surgical excision of the BCM was performed. The patient was operated on in a right park-bench position, the head being slightly rotated on the right to better expose the left side of the posterior fossa. Following a median incision and a median posterior fossa craniotomy, a telovelar approach was used to gain access to the rhomboid fossa. The exophytic hematoma appeared clearly on the left side at the level of the *striae medullares*, thus enabling us to

remove the hematoma and the adjoining cavernoma through the infrafacial triangle. The DVA was left intact (Figs. 3, 4, Video 2). The postoperative course was complicated by a surgical site infection requiring surgical revision, placement of a temporary external ventricular drain and combined antibiotic therapy (meropenem and linezolid). The patient suffered from a left-sided grade VI House-Brackmann peripheral facial nerve palsy, further complicated by a corneal ulcer which was managed with local treatment. He also presented postoperatively with a non-pre-existing left-sided glossopharyngeal nerve (CN IX) palsy responsible for dysphagia and aspiration pneumonia, requiring a temporary gastroplasty. The patient was finally sent to neurological rehabilitation 3 months after the procedure.

Discussion

Scope of the review

In the light of this case report, our aim was to evaluate the proportion of CBS caused by hemorrhagic BCMs. We purposely chose to restrict the search to the five most frequent and widely recognized CBS, namely Benedikt (paramedian midbrain syndrome), Weber (superior




Fig. 3 Postoperative MRI of the brain in post-contrast T1-weighted sequence. We performed (d) a suboccipital telovelar approach to gain access to the rhomboid fossa. Then we used (c) the infrafacial triangle as an entry point to the pons to perform microsurgical excision of (a) the BCM and (b,c) the pontine hematoma. a,b,c,d The DVA was left intact




Fig. 4 Important surgical sequences. **a** Hockey stick fashion opening of the dura mater, discovering the cerebellar notch medially, and the left superior and inferior semilunar lobules. **b** The floor of the fourth ventricle appears after passing through the inferior medullary velum, and displays crucial anatomical landmarks such as the *striae medullares* (SM), the infrafacial triangle (IFT) just above and the hypoglossal triangle (HT) underneath. The exophytic hematoma appears clearly on the left side at the level of the *striae medullares*. The cerebellar developmental venous anomaly (DVA) appears in blue under a thin layer of nervous tissue. **c** Evacuation of the pontine hematoma (H) at the level of the infrafacial triangle (IFT), just above the *striae medullares* (SM). The midline is marked with a dotted line. **d** En-bloc excision of the cavernous malformation (CAV) using a tumor's clamp. **e** Resection cavity

alternating hemiplegia), Foville (inferior medial pontine syndrome), Millard-Gubler (ventral pontine syndrome), and Wallenberg (lateral medullary syndrome) syndromes.

Database research

We conducted a comprehensive literature review on Medline database (<https://pubmed.ncbi.nlm.nih.gov/>) from inception to 2020. We used the advanced search mode with the following Mesh terms in the title or in the text: Benedikt, Weber, Foville, Millard-Gubler, Wallenberg.

Inclusion and exclusion criteria

In the first instance, all the articles describing a CBS were retained regardless of the language and were screened in a systematic manner. The following information was extracted as previously planned: author, year, patient's age, name of the crossed brainstem syndrome, and etiology. When the full text was not available, the abstract was analyzed in search of the same information. Exclusion criteria consisted in articles with no genuine or dubious CBS, no patient's age, or no clear reference as to the underlying etiology.

Results of database research

The primary database research yielded 234 articles, among which 168 met the exclusion criteria after careful reading of the text or the abstract. Sixty-six articles were finally retained for a total of 69 patients [2]. There were 14 cases of Benedikt syndrome [3], three cases of Weber syndrome [4], 15 cases of Foville syndrome [5], nine cases of Millard-Gubler syndrome [6, 7], and 28 cases of Wallenberg syndrome [8] (Table 1).

Causes of crossed brainstem syndromes

At the level of the midbrain, Benedikt syndrome was usually caused by ischemic stroke ($n = 6/14$), followed by hemorrhage ($n = 4/14$) and direct nervous compression ($n = 3/14$) [9–22]. Weber syndrome was mainly caused by hemorrhage ($n = 1/3$) or infectious etiologies ($n = 2/3$) [4, 23, 24]. At the level of the pons, Foville syndrome was frequently caused by hemorrhage ($n = 8/15$), followed by ischemic stroke ($n = 4/15$) and brain metastases ($n = 2/15$) [25–37]. Conversely, Millard-Gubler syndrome was mostly related to an ischemic stroke ($n = 7/9$), and rarely brought about by hemorrhage ($n = 1/9$) or brain abscess ($n = 1/9$) [38–46]. At the level of the *medulla oblongata*, Wallenberg syndrome was

Table 1 Medline based literature review on the five most common crossed brainstem syndromes

Region	Midbrain			Pons			Medulla oblongata				
	Benedikt syndrome (Paramedian midbrain syndrome)	Weber syndrome (Superior alternating hemiplegia)	Foville syndrome (Inferior medial pontine syndrome)	Millard-Gubler syndrome (Ventral pontine syndrome)	Wallenberg syndrome (Lateral medullary syndrome)	Year	age	Etiology	Year	age	Etiology
Crossed brainstem syndrome	CN III	CN III	CN VI and VII	CN VII	CN V, VIII, IX, X, Horner's sd, cerebellar sd						
Ipsilateral symptoms	cerebellar ataxia, hemiparesis, hyperactive reflexes	hemiparesis	hemiparesis and hemianesthesia sparing the face	hemiparesis and hemianesthesia sparing the face	thermoalgesic anesthesia						
Contralateral symptoms											
Year	1889	1863	1859	1856	1895						
author	Benedikt Weber	Foville	Millard	Wallenberg	Amantini						
Year	1974	2014	1939	1856	1982						
age	44	62	Stroke	Stroke	70						
Etiology	Stroke	Encephalitis (HSV)	Stroke	Stroke	Stroke						
Year	1981	2018	1943	1993	1984						
author	Loseke	Parija	Freeman	Yasuda	Dharmoon						
Year	1992	2018	1943	1993	1999						
age	60	28	60	60	56						
Etiology	Stroke	Brain abscess (tuberculosis)	Hemorrhage (autopsy)	Probable hemorrhage	Stroke						
Year	1994	2018	1994	2005	2000						
author	Ono	Ono	Ono	Onbas	Faust						
Year	1995	1995	1957	2010	2000						
age	51	Stroke	48	45	49						
Etiology	Stroke	Multiloculated cyst	Hemorrhage (pheochromocytoma)	Stroke (infectious thrombosis)	Stroke (Wegener vasculitis)						
Year	1997	1997	1958	2011	2000						
author	Borras	Borras	Leslie	Kesikburun	Miyazaki						
Year	1999	2005	1981	2012	2004						
age	38	53	48	7	48						
Etiology	Hemorrhage (after CN V vascular decompression)	Hemorrhage (after stereotactic biopsy)	Stroke (basilar aneurysm clipping)	Abscess (neurocysticercosis)	Stroke						
Year	2005	2008	1996	2013	2004						
author	Akdal	Akdal	Hubloue	Ahdab	Kim						
Year	2008	2008	2000	2019	2005						
age	55	Stroke	88	58	44						
Etiology	Stroke	Stroke	Hemorrhage (hypertension)	Stroke	Stroke						
Year	2011	2011	2009	2019	2008						
author	Sturiale	Sturiale	Nakaso	Li	Zhang						
Year	2013	2013	2013	2019	2009						
age	49	Hemorrhage (BCM)	61	49	61						
Etiology	Hemorrhage (BCM)	Hemorrhage (BCM)	Hemorrhage	Stroke	Stroke						
Year	2015	2015	2014	2009	2009						
author	Koskela	Koskela	Canepa	Ceballos	Porta-Etessam						
Year	2018	2018	2015	2019	2009						
age	66	Stroke (basilar aneurysm clipping)	44	52	56						
Etiology	Stroke (basilar aneurysm clipping)	Stroke (basilar aneurysm clipping)	Brain metastasis	Multiple sclerosis	Stroke						
Year	2018	2018	2016	2016	2009						
author	Yamanaka	Yamanaka	Massi	Yeh	Yeh						
Year	2018	2018	2016	2009	2009						
age	16	Hemorrhage (BCM)	68	38	38						
Etiology	Hemorrhage (BCM)	Hemorrhage (telangiectasia)	Hemorrhage	Stroke (traumatic vertebral artery dissection)	Stroke						
Year	2013	2013	2016	2009	2013						
author	Cheng	Cheng	Selvadurai	Zabaleta	Zhu						
Year	2013	2013	2016	2009	2013						
age	73	Stroke (Horton giant cell)	73	53	73						
Etiology	Stroke (Horton giant cell)	Stroke (Horton giant cell)	Stroke (Horton giant cell)	Stroke	Stroke						

Table 1 Medline based literature review on the five most common crossed brainstem syndromes (Continued)

Region	Pons			Medulla oblongata
	Midbrain	Weber syndrome (Superior alternating hemiplegia)	Foville syndrome (Inferior medial pontine syndrome)	
Crossed brainstem syndrome	Benedikt syndrome (Paramedian midbrain syndrome)			Wallenberg syndrome (Lateral medullary syndrome)
				Stengl arteritis
				2013 Ueda 48 Hemorrhage (multiple BCM)
				72 Hemorrhage (antiplatelet and anticoagulant therapy)
			2014 Wu	43 Stroke
			2015 Koskela	Unruptured aneurysm
			2015 Das	86 Stroke
			2015 Ehresmann	7 Stroke
			2015 Louis	30 Stroke (2 weeks post-partum)
			2015 Ospino Quiroz	48 Stroke
			2018 Kornbluh	14 Stroke
			2018 Oks	58 Stroke (sarcooidosis)
			2018 Sivakumar	62 Stroke (PICA aneurysm clipping)

predominantly caused by ischemic stroke ($n = 23/28$), more rarely by hemorrhage ($n = 2/28$) or multiple sclerosis ($n = 1/28$) [18, 47–71]. The complete data is provided in Table 2.

Brainstem hemorrhage was responsible for approximately one quarter of the cases of CBS ($n = 15/66$). As for the underlying condition responsible for the brainstem bleeding, hypertension was the most frequently encountered etiology ($n = 6/15$), closely followed by BCM ($n = 5/15$). Extralesional bleeding arising from BCM was responsible for one-seventh of the cases of Benedikt syndrome ($n = 2/14$), one out of ten cases of Millard-Gubler syndrome ($n = 1/9$), one-fifteenth of the cases of Foville syndrome ($n = 1/15$), and approximately one out of thirty cases of Wallenberg syndrome ($n = 1/28$). There was also one case of Foville syndrome caused by a hemorrhage imputed to a telangiectasia.

It is to note that posterior circulation aneurysms were frequently encountered in this review ($n = 5/69$). Two unruptured aneurysms were responsible for nervous compression, the first one (probably arising from the

posterior communicating artery) leading to a case of Benedikt syndrome and the second one (arising from the posterior inferior cerebellar artery) at the origin of a Wallenberg syndrome. Three aneurysms clipping resulted in infarction of perforating arteries, causing respectively a Benedikt syndrome, a Foville syndrome, and a Wallenberg syndrome.


Similarly, two cases of Benedikt syndrome were caused by a midbrain hematoma which occurred immediately after a neurosurgical procedure: one was secondary to a third ventricle tumor biopsy, and the other one was secondary to microvascular decompression for trigeminal neuralgia.

Physiopathology of cerebral cavernous malformations

Cerebral cavernous malformations (CCM) are mulberry-like fragile vascular malformations that are encountered in the cerebral hemispheres, brainstem and cerebellum, or in the spinal cord. Their structure consists in endothelial lined vascular sinusoids with no tight junctions and even gaps between the endothelial cells, forming

Table 2 Etiologies reported for the five most common crossed brainstem syndromes

	Total	Benedikt	Weber	Foville	Millard-Gubler	Wallenberg
Total	69	14	3	15	9	28
Stroke	40	6	0	4	7	23
Embollic event	30	5		1	6	18
Aneurysm clipping	3	1		1		1
Artery dissection	2			1		1
Transient ischemic attack	1			1		
Vasculitis	2					2
Infectious thrombosis	1				1	
Sarcoidosis	1					1
Hemorrhage	16	4	1	8	1	2
Hypertension	6			6		
Brainstem cavernous malformation	5	2		1	1	1
Telangiectasia	1			1		
Post-operative complication	2	2				
Anticoagulant therapy	1					1
Compression	5	3				2
Unruptured aneurysm	2	1				1
Cyst / Virchow-Robin spaces	2	2				
Occipital fracture	1					1
Brain metastasis	3	1		2		
Infection	3		2		1	
Brain abscess	2		1		1	
Encephalitis	1		1			
Multiple sclerosis	1					1
Unknown	1			1		



caverns within a dense collagen matrix clustered without intervening normal parenchyma [72].

CCM are often associated with venous drainage anomalies, ranging from solitary trans-cerebral or subpial draining veins to genuine DVAs [73]. DVA constitute an extreme anatomical variation draining normal cerebral tissue into an extra-parenchymatous collector; they reflect a variation of the well-known anastomosis between the superficial and the deep venous drainage systems of the brain which respond to a hemodynamic equilibrium [74].

The combination of inherently fragile sinusoids walls in the absence of blood-brain barrier and DVAs with raised venous pressure results in repeated intralesional micro hemorrhages which, in turn, leads to neoangiogenesis [75]. This “hemorrhagic angiogenic proliferation” mechanism results over time in the self-sustained growth of CCM, which is why they appear on neuroimaging as multilobulated vascular and calcified “pop-corn” lesions as the type 2 described by Zabramski [76]. Although half of the CCM are discovered incidentally on neuroimaging, the other half may cause seizures related to the hemosiderin deposit around the lesion causing cortical irritation (25%), focal neurological symptoms related to mass effect (15%), or intracranial hemorrhage (ICH) (12%) [77].

Specific considerations for brainstem CCM

It comes as no surprise that in the brainstem the most feared complication of CCM turns out to be bleeding which is also the main indication of excisional surgery [78]. The two main risk factors for the occurrence of an ICH ascribable to a CCM are history of a previous bleeding episode and the location in the brainstem [79]. Indeed, the estimated 5-year risk of ICH for an untreated CCM is 3.8% in case of non-brainstem CCM without ICH or FND, 8% in case of BCM without ICH or FND, 18.4% for non-brainstem CCM with ICH or FND, and increases up to 30.8% for BCM with ICH or FND [80]. In the brainstem, the estimated annual rate of extralesional bleeding is 8.7% for asymptomatic CCM, and rises to 12.4% for CCM with asymptomatic ICH, and up to 15.9% for CCM with symptomatic ICH [81].

Relevant surgical anatomy of the pons

At the middle pons, corticospinal tract fibers are scattered anteriorly; motor neurons transit through transverse pontine fibers to merge the contralateral pontine nuclei and then join the middle cerebellar peduncle. The spinothalamic tract is located just posteriorly and lies within the medial lemniscus. The floor of the fourth ventricle provides a few surface reliefs that constitute

important landmarks for neurosurgeons. The medial sulcus is bordered by the medial longitudinal fasciculi on both sides. The nucleus of the facial nerve is located laterally at the inferior part of the pons. The fibers of the future CN VII loop superiorly and medially around the abducens nerve nucleus. This peculiar anatomical configuration creates a bulging within the floor of the fourth ventricle known as the facial colliculus. Inferiorly, the *striae medullares* define the superior limit of the hypoglossal (CN XII), ambiguous (CN IX, X, XI) and vagus (CN X) *nuclei*. Pontine arterial supply is mainly anterior and lateral; no major artery is to be found near the floor of the fourth ventricle floor (Fig. 5).

Surgical approaches to the pons

The facial colliculus along with the fibers of future CN VII represent an important surgical landmark within the rhomboid fossa. They constitute the inferior limit of the suprafacial triangle which superior border are the superior and the middle cerebellar peduncles. On the same way, they constitute the superior limit of the infrafacial triangle which inferior borders are the *striae medullares*. These two triangles are known to be relatively safe entry corridors entry corridors for a surgical approach to the floor of the fourth ventricle as only scarce nerve fibers are encountered there [82, 83].

Surgical considerations for brainstem cavernous malformations

Recent literature does not provide sufficient evidence regarding the optimal timing for the surgical excision of a brainstem CCM with symptomatic extralesional bleeding, which is still a matter of debate. Zaidi et al. presented a series of 397 patients operated on for brainstem CCM, among which 96% percent presented history of prior ICH [84]. Thirty-five percent of the patients presented persistent postoperative neurological deficits (mainly CN deficits), and the mean GOS was unchanged at last follow-up compared with the GOS upon admission (4.47 vs 4.46, median follow-up 35.5 months). They reported that early surgery within 6 weeks after ICH and smaller lesion size were associated with improved outcome. Garcia et al. presented a series of 104 patients operating on for brainstem CCM, among which 99% presented history of prior ICH [85]. The mean modified Rankin scale upon admission was 2.23 compared to 1.58 at final follow-up. The most frequent perioperative complications were cerebrospinal fluid leakage (12.5%), infection (9.6%) and surgical site hematoma (6%). Older age, large size lesions, lesions crossing the midline, delay between last bleeding event and surgery, and the association with a DVA were associated with a poorer prognosis.

Based on these retrospective series, surgical excision of a BCM may be deemed reasonable soon after the second symptomatic bleeding. In such case, the high operative morbidity inherent to brainstem surgery is warranted by the aggressive natural course of the disease.

Surgical considerations for associated developmental venous anomalies

Until the 2000s, there have only been sporadic reports on the treatment of DVA. Some reported cases supported the surgical excision of the DVA [86, 87], whereas intraoperative complications such as brain swelling after DVA coagulation have been reported [88]. Campeau et al. neuro-imaging study seemed to confirm the hypothesis that repeated microbleeding episodes and neoangiogenesis led to the formation of CCM in the vicinity of DVA [89]. In accordance with that theory, Wurm et al. reported a series of 15 patients who benefited from microsurgical excision of a CCM [90]. The associated DVA was coagulated in six patients and left intact in nine of them. Three patients from the group with intact DVA presented the recurrence of a CCM and benefited from a second microsurgical excision with simultaneous coagulation of the DVA. The authors did not report any venous complication in the patients who benefited from the treatment of the DVA, with a mean follow-up of 29 months. Nevertheless, this series, made up of only 15 patients, lacks long-term follow-up. Besides, six patients whose DVA had been left intact did not present recurrence of CCM. More recent reports continue to support the elective microsurgical excision of symptomatic CCM without touching the associated DVA [91]. Venous sacrifice in cranial neurosurgical procedures has always been considered hazardous for fear of the potential disastrous consequences of venous infarction [92, 93], which are very difficult to predict [94]. Consequently, we tend to recommend leaving the DVA intact during the microsurgical excision of CCM.

Specific considerations for crossed pontine syndromes

The specific vascular supply of the pons may explain the difference of etiology that we have observed between Foville syndrome (the inferior medial pontine syndrome) and Millard-Gubler syndrome (the ventral pontine syndrome). Pontine hemorrhage caused by high blood pressure is usually located more medially and damages both CN VI nucleus and CN VII fibers, leading to Foville syndrome. By contrast, ischemic stroke involves rather the paramedian branches or the short circumferential branches of the basilar artery which supply more lateral structures such as CN VII nucleus, leading thus to Millard-Gubler syndrome [95].

If we closely examine the clinical nuances reported throughout the history concerning Foville syndrome, the

Table 3 The inferior medial pontine syndrome of Foville: clinical nuances reported since its first description

Author - year	Journal	Age	Cause	VI palsy	Ipsilateral superior VII palsy	Ipsilateral inferior VII palsy	Lateral gaze palsy	Face-sparing hemiparesis / plegia	Proportional hemiparesis / plegia	Contralateral hemianesthesia	Contralateral sympathetic symptoms
Foville 1859 [5]	Gaz Hebd Med Chir					yes	ipsilateral	contralateral			
Mutch 1939 [25]	Brit J Ophthalmology	56		ipsilateral	yes	yes	ipsilateral	ipsilateral lower limb			
Freeman 1943 [26]	Arch Neurology & Psychiatry	25		ipsilateral	yes	yes	ipsilateral	contralateral		yes	
		60	pontine hemorrhage	both sides	yes	yes	both sides	contralateral upper limb		yes	
Melkild 1957 [27]	Acta Med Scand	48	hemorrhage (pheochromocytoma)	ipsilateral	yes	yes	ipsilateral	contralateral			
Leslie 1958 [28]	J Am Geriatrics Soc	76	cerebellar tumor	ipsilateral	yes	yes	ipsilateral	ipsilateral pyramidal			
Takase 1981 [29]	Shinkei Neurol Surg	48	basilar aneurysm clipping				ipsilateral deviation		yes		
Hubloue 1996 [30]	Eur J Emerg Med	74	transient ischemic attack	ipsilateral	yes	yes	ipsilateral	contralateral		yes	
Sato 2000 [31]	Rinsho Shinkeigaku Clin Neurol	88	pontine hemorrhage	ipsilateral	yes	yes		contralateral		proportional	yes
Nakaso 2009 [32]	Internal Medicine	61	pontine hemorrhage (cavernoma)				ipsilateral	contralateral			
Cheng 2013 [33]	Taiwan Journal of Ophthalmology		pontine hemorrhage	ipsilateral internuclear ophthalmoplegia	yes	yes		contralateral		yes	
Canepa-Raggio 2014 [34]	BMI Case reports		infarction (vertebral a. dissection)		numbness	numbness		contralateral upper limb		yes	
Man 2015 [35]	BMI Case reports	44	pontine lung metastasis		yes	yes	ipsilateral	contralateral			
Massi 2016 [36]	Pan Afr Med J	20	pontine hemorrhage	ipsilateral	yes	yes	ipsilateral	contralateral			
Selvaudurai 2016 [37]	Neurology	68	pontine hemorrhage (telangiectasia)				ipsilateral	contralateral			

different forms of oculomotor palsies that were observed led to the distinction between a “superior Foville syndrome” characterized by the presence of a CN VI palsy and an “inferior Foville syndrome” with lateral conjugate gaze palsy due to the involvement of the medial longitudinal fasciculus or the paramedian pontine reticular formation (Table 3).

Limitations of the study

This review presents some limits inherent to its retrospective nature. Purposely or not case reports unconsciously select patients with favorable outcome; thus, their compilation may lead to a reporting bias which may underestimate the mortality rate. The literature review was deliberately restricted to the five most common CBS which may constitute a limit but still enabled us to collect a great number of articles. To the best of our knowledge, this is the first study attempting to provide a clear and updated picture of the proportion of BCMs responsible for or revealed by a genuine CBS.

Conclusions

Pure crossed brainstem syndromes are rarely encountered in clinical practice. They remarkably illustrate the anatomical peculiarity of the brainstem, which represents a crossroad between the cranial nerves, the long tracts and key vegetative structures. In the light of this review, brainstem cavernous malformations with extraleSIONAL bleeding appear to account for approximately 7 % of all crossed brainstem syndromes. The indication and timing of the surgical excision of a symptomatic brainstem cavernous malformation remains a complex decision to make and requires multidisciplinary team expertise. It has to be discussed openly between neurosurgeons and their patient, taking into consideration the existing evidence in favor of surgery but also the substantial risks associated with such a delicate procedure. Multicentric prospective trials will be very difficult to conduct on such rare entities. Robust knowledge in brainstem anatomy along with thorough neurological examination skills will remain pivotal to the initial management of these patients.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12883-021-02223-7>.

Video 1
Video 2

Acknowledgements

None.

Authors' contributions

NB and HD conceptualized the article. NB, SB, AR, SF, RC, HD participated to the literature review. NB, SB, AR, SF, RC, HD participated to the clinical care of the patient. NB, SB, AR, SF, RC, HD participated to the drafting of the manuscript. NB, RC, HD participated to the critical revision of the manuscript. All authors have read and approved the manuscript.

Funding

The authors did not receive any funding for this work. The publication charges of this article were gracefully covered by the "neuro-oncology and neuro-endocrinology research group" (Grenone) of Timone University Hospital, 13010 Marseille, France.

Availability of data and materials

All the relevant data is included in the manuscript. There is no data deposit for this work.

Declarations

Ethics approval and consent to participate

Informed written consent was obtained from the patient whose case report is included in the manuscript. He has been given the opportunity to review the manuscript and the attached files. This work was conducted in accordance with the Declaration of Helsinki of 1964 or its further amendments (2013).

Consent for publication

Written informed consent was obtained from the patient for the publication of this manuscript and any accompanying figure and video. A copy of the written consent is attached to the manuscript.

Competing interests

All authors certify that they have no affiliations with or involvement in any organization or entity with any financial interest (such as honoraria; educational grants; participation in speakers' bureaus; membership, employment, consultancies, stock ownership, or other equity interest; and expert testimony or patent-licensing arrangements), or non-financial interest (such as personal or professional relationships, affiliations, knowledge or beliefs) in the subject matter or materials discussed in this manuscript.

Author details

¹Department of Neurosurgery, Timone University Hospital, APHM, 264 rue Saint-Pierre, 13005 Marseille, France. ²Ecole du Val-de-Grâce, French Military Health Service Academy, 1 place Alphonse Laveran, 75230 Paris Cedex 5, France. ³Aix Marseille Univ, INSERM, INS, Inst Neurosci Syst, Marseille, France. ⁴Emergency Department, Timone University Hospital, APHM, 264 Rue Saint-Pierre, 13005 Marseille, France. ⁵Department of Stereotactic and Functional Neurosurgery, Timone University Hospital, APHM, 264 rue Saint-Pierre, 13005 Marseille, France. ⁶Aix-Marseille Univ, INSERM, MMG, Marseille, France.

Received: 7 December 2020 Accepted: 4 May 2021

Published online: 20 May 2021

References

1. Querol-Pascual MR. Clinical approach to brainstem lesions. *Semin Ultrasound CT MRI*. 2010;31(3):220–9. <https://doi.org/10.1053/j.sult.2010.03.004>.
2. Marx JJ, Thömke F. Classical crossed brain stem syndromes: myth or reality? *J Neurol*. 2009;256(6):898–903. <https://doi.org/10.1007/s00415-009-5037-2>.
3. Benedikt M. lecture. 1889;
4. Weber H. A contribution to the pathology of the Crura Cerebri. *J R Soc Med*. 1863;MCT-46(1):121–39. <https://doi.org/10.1177/095952876304600112>.
5. Foville A. Note sur une paralysie peu connue de certains muscles de l'oeil et sa liaison avec quelques points de l'anatomie et la physiologie de la protubérance annulaire. *Bull Soc Anat. (Paris)*. 1858;33:373-405.
6. Cenac H, Millard A. Hémorragie de la protubérance annulaire. *Bull Soc Anat. (Paris)*. 1856;36:206-21.
7. Gubler A. De l'hémiplégie alterne envisagée comme signe de lésion de la protubérance annulaire et comme preuve de la décuSSation. *Gaz Hebd Méd Chir*. 1856;3(43):749–54.

8. Wallenberg A. Akute Bulbäreffektion (Embolie der Arteria cerebelli post inf sinistra). *Arch Psychiatry*. 1895;27(2):504–40. <https://doi.org/10.1007/BF02075799>.
9. Fujieda T, Yamauchi T, Takahashi S, Moroji T. Letter: effect of levodopa on tremor in Benedikt's syndrome. *Br Med J*. 1974;1(5905):456–7. <https://doi.org/10.1136/bmj.1.5905.456-b>.
10. Loseke N, Retif J, Noterman J, Flament-Durand J. Inferior red nucleus syndrome (Benedikt's syndrome) due to a single intramesencephalic metastasis from a prostatic carcinoma. *Case Rep Acta Neurochir (Wien)*. 1981;56(1–2):59–64.
11. Ono Y, Suzuki M, Kayama T, Yoshimoto T. Multilobulated cystic formation in the brain stem with Benedikt's syndrome: case report. *Neurosurgery*. 1994;34(4):726–9 discussion 729.
12. Borrás JM, Salazar FG, Grandas F. Oculomotor palsy and contralateral tremor (Benedikt's syndrome) following a stereotactic procedure. *J Neurol*. 1997;244(4):272–4. <https://doi.org/10.1007/s004150050085>.
13. Fernandez HH, Friedman JH, Centofanti JV. Benedikt's syndrome with delayed-onset rubral tremor and hemidystonia: a complication of tic douloureux surgery. *Mov Disord Off J Mov Disord Soc*. 1999;14(4):695–7. [https://doi.org/10.1002/1531-8257\(199907\)14:4<695::AID-MDS1024>3.0.CO;2-3](https://doi.org/10.1002/1531-8257(199907)14:4<695::AID-MDS1024>3.0.CO;2-3).
14. Akdal G, Kutluk K, Men S, Yaka E. Benedikt and "plus-minus lid" syndromes arising from posterior cerebral artery branch occlusion. *J Neurol Sci*. 2005;228(1):105–7. <https://doi.org/10.1016/j.jns.2004.09.029>.
15. Bandt SK, Anderson D, Biller J. Deep brain stimulation as an effective treatment option for post-midbrain infarction-related tremor as it presents with Benedikt syndrome. *J Neurosurg*. 2008;109(4):635–9. <https://doi.org/10.3171/JNS/2008/109/10/0635>.
16. Sturiale CL, Albanese A, Lofrese G, Frassanito P, Sabatino G, Marchese E, et al. Pathological enlargement of midbrain Virchow–Robin spaces: a rare cause of obstructive hydrocephalus. *Br J Neurosurg*. 2011;25(1):130–1. <https://doi.org/10.3109/02688697.2010.504050>.
17. Maduri R, Barbagallo G, Iofrida G, Signorelli M, Signorelli F. Regression of Benedikt's syndrome after single-stage removal of mesencephalic cavernoma and temporal meningioma: a case report. *Clin Neurol Neurosurg*. 2013;115(6):748–50. <https://doi.org/10.1016/j.clineuro.2012.06.033>.
18. Koskela E, Setälä K, Kivisaari R, Hernesniemi J, Laakso A. Neuro-ophthalmic presentation and surgical results of unruptured intracranial aneurysms—prospective Helsinki experience of 142 patients. *World Neurosurg*. 2015;83(4):614–9. <https://doi.org/10.1016/j.wneu.2014.12.017>.
19. Cheng G, Yang Y, Wang Y, Tan H, Zhang S. Deep brain stimulation of the thalamic ventral intermediate nucleus for Benedikt's syndrome mainly present as tremor: a long-term case observation. *Acta Neurochir*. 2018;160(7):1349–53. <https://doi.org/10.1007/s00701-018-3526-8>.
20. Mateos V, Campos DM, Colosía VP, Salas-Puig J, Fernández JM, Lahoz CH. Nuclear syndrome of the oculomotor nerve caused by a mesencephalic infarction confirmed by MRI. *Arch Neurobiol (Madr)*. 1992;55(4):183–7.
21. Duncan GW, Weindling SM. Posterior cerebral artery stenosis with midbrain infarction. *Stroke*. 1995;26(5):900–2. <https://doi.org/10.1161/01.STR.26.5.900>.
22. Yamanaka Y, Shinohara T, Kozano I, Yoshizumi T, Kawasaki T. Benedikt syndrome associated with neck clipping of ruptured basilar-superior cerebellar artery aneurysms: a case report. *No Shinkei Geka*. 2018;46(11):1007–12. <https://doi.org/10.11477/mf.1436203855>.
23. Ballaekere JS, Chebbi PP, Sundarmurthy H, Parameshwarappa A. Weber syndrome: herpes simplex virus brainstem encephalitis as an etiology. *Am J Med*. 2014;127(12):e5–6. <https://doi.org/10.1016/j.amjmed.2014.08.005>.
24. Parija S, Lalitha CS, Naik S. Weber syndrome secondary to brain stem tuberculoma. *Indian J Ophthalmol*. 2018;66(7):1036–9. https://doi.org/10.4103/ijo.IJO_1040_17.
25. Mutch JR. FOVILLE'S syndrome record of a CASE. *Br J Ophthalmol*. 1939;23(4):225–38. <https://doi.org/10.1136/bjo.23.4.225>.
26. Freeman W. SYNDROMES OF, THE PONTILE TEGMENTUM. FOVILLE'S syndrome: report of three cases. *Arch Neurol Psychiatr*. 1943;50(4):462. <https://doi.org/10.1001/archneurpsyc.1943.02290220092008>.
27. Melkild A, Mörstád KS. Foville's syndrome as a complication of Phaeochromocytoma. *Acta Med Scand*. 1957;159(6):471–4. <https://doi.org/10.1111/j.0954-6820.1957.tb00155.x>.
28. Levene LJ. Benign cerebellar tumor presenting a modified Foville syndrome. *J Am Geriatr Soc*. 1958;6(4):346–51. <https://doi.org/10.1111/j.1532-5415.1958.tb00726.x>.
29. Takase M, Saeki N, Oka N, Satoh A, Otaki M, Yamaura A. Superior Foville syndrome after clipping of basilar bifurcation aneurysm—case report (author's transl). *No Shinkei Geka*. 1981;9(3):343–7.
30. Hubloue I, Laureys S, Michotte A. A rare case of diplopia: medial inferior pontine syndrome or Foville's syndrome. *Eur J Emerg Med*. 1996;3(3):194–8.
31. Sato K, Nitta E. Pontine hemorrhage presenting with Foville syndrome and transient contralateral hyperhidrosis. *Rinsho Shinkeigaku*. 2000;40(3):271–3.
32. Nakaso K, Sasaki K. Fluctuating Foville's syndrome caused by a pontine Angioma in a patient with a polycystic kidney. *Intern Med*. 2009;48(15):1335–6. <https://doi.org/10.2169/internalmedicine.48.2314>.
33. Cheng H-C, Yen M-Y, Wang A-G. Foville's syndrome with ipsilateral internuclear ophthalmoplegia due to spontaneous pontine hemorrhage. *Taiwan J Ophthalmol*. 2013;3(2):75–7. <https://doi.org/10.1016/j.tjjo.2012.12.001>.
34. Canepa Raggio C, Dasgupta A. Three cases of Spontaneous Vertebral Artery Dissection (SVAD), resulting in two cases of Wallenberg syndrome and one case of Foville syndrome in young, healthy men. *Case Rep*. 2014;2014(apr28 2):bcr2014203945.
35. Man BL, Fu YP. Bronchogenic carcinoma presented as Foville's syndrome. *Case Rep*. 2015;2015(mar03 2):bcr2014205025.
36. Massi DG, Nyassinde J, Ndiaye MM. Superior Foville syndrome due to pontine hemorrhage: a case report. *Pan Afr Med J*. 2016;25:215.
37. Selvadurai C, Rondeau MW, Colorado RA, Feske SK, Prasad S. Teaching video neuro *Images* : Foville syndrome. *Neurology*. 2016;86(19):e203. <https://doi.org/10.1212/WNL.0000000000002658>.
38. Matlis A, Kleinman Y, Korn-Lubetzki I. Millard-Gubler syndrome. *AJNR Am J Neuroradiol*. 1994;15(1):179–81.
39. Yasuda Y, Matsuda I, Sakagami T, Kobayashi H, Kameyama M. Pontine infarction with pure Millard-Gubler syndrome: precise localization with magnetic resonance imaging. *Eur Neurol*. 1993;33(4):333–4. <https://doi.org/10.1159/000116965>.
40. Onbas O, Kantarci M, Alper F, Karaca L, Okur A. Millard-Gubler syndrome: MR findings. *Neuroradiology*. 2005;47(1):35–7. <https://doi.org/10.1007/s00234-004-1312-1>.
41. Rose DZ, Parra-Herran C, Petito CK, Post MJD. Restricted diffusion of pus in the subarachnoid space: MRSA Meningo-Vasculitis and progressive brainstem ischemic strokes - a case report. *Case Rep Neurol*. 2010;2(2):101–10. <https://doi.org/10.1159/000319691>.
42. Kesikburun S, Safaz İ, Alaca R. Pontine cavernoma hemorrhage leading to millard-gubler syndrome. *Am J Phys Med Rehabil*. 2011;90(3):263. <https://doi.org/10.1097/PHM.0b013e3181e29e8e>.
43. Prasad R, Kapoor K, Mishra OP, Srivastava A. Neurocysticercosis presenting as Millard Gubler syndrome. *J Neurosci Rural Pract*. 2012;03(03):375–7.
44. Ahdab R, Saade HS, Kikano R, Ferzli J, Tarcha W, Riachi N. Pure ipsilateral central facial palsy and contralateral hemiparesis secondary to ventromedial medullary stroke. *J Neurol Sci*. 2013;332(1–2):154–5. <https://doi.org/10.1016/j.jns.2013.06.028>.
45. Ceballos-Lizarraga R, Palomino-Díaz C, Romero-Figueroa JA. Wall-eyed monocular Internuclear Ophthalmoplegia (WEMINO) and Millard-Gubler syndromes in a patient with isolated pontine infarction: topographic, oculomotor, and radiological analysis of two very uncommon conditions. *Case Rep Neurol*. 2019;11(2):230–7. <https://doi.org/10.1159/000501794>.
46. Li X-T, Yuan J-L, Hu W-L. Vertebrobasilar artery dissection manifesting as Millard-Gubler syndrome in a young ischemic stroke patient: a case report. *World J Clin Cases*. 2019;7(1):73–8. <https://doi.org/10.12998/wjcc.v7.i1.73>.
47. Amantini A, Arnetoli G, Rossi L, Fenzi, Salviati A, Rizzuto N, et al. BAEP and autopsy findings in Wallenberg syndrome. *Ital J Neurol Sci*. 1982;3(3):237–40. <https://doi.org/10.1007/BF02043316>.
48. Dhamoon SK, Iqbal J, Collins GH. Ipsilateral hemiplegia and the Wallenberg syndrome. *Arch Neurol*. 1984;41(2):179–80. <https://doi.org/10.1001/archneur.1984.04050140077029>.
49. Rousseaux M, Cassim F, Bayle B, Laureau E. Analysis of the perception of and reactivity to pain and heat in patients with Wallenberg syndrome and severe spinothalamic tract dysfunction. *Stroke*. 1999;30(10):2223–9. <https://doi.org/10.1161/01.STR.30.10.2223>.
50. Faust J, Visbeck A, Fitzek S, et al. Vasculitic wallenberg syndrome with detection of anti-proteinase 3 antibodies in the cerebrospinal fluid of a patient with severe Wegener's granulomatosis and only mild kidney involvement. *Nephrol Dial Transplant Off Publ Eur Dial Transpl Assoc Eur Ren Assoc*. 2000;15(6):893–6.
51. Miyazaki C, Katsume M, Yamazaki T, Aoki K, Kuroki T, Takasu N. Unusual occipital condyle fracture with multiple nerve palsies and Wallenberg

- syndrome. *Clin Neurol Neurosurg.* 2000;102(4):255–8. [https://doi.org/10.1016/S0303-8467\(00\)00109-8](https://doi.org/10.1016/S0303-8467(00)00109-8).
52. Hippius WM, Wilhelmus KR. Persistent visual loss from neurotrophic corneal ulceration after dorsolateral medullary infarction (Wallenberg syndrome). *J Neuroophthalmol.* 2004;24(4):345–6. <https://doi.org/10.1097/00041327-200412000-00015>.
 53. Kim JS, Moon SY, Park S-H, Yoon B-W, Roh J-K. Ocular lateropulsion in Wallenberg syndrome. *Neurology.* 2004;62(12):2287. <https://doi.org/10.1212/WNL.62.12.2287>.
 54. Nomoto N, Konno S, Sugimoto H, Kurihara T, Wakata N. Wallenberg syndrome with SIADH. *Eur Neurol.* 2005;53(1):35–6. <https://doi.org/10.1159/000084260>.
 55. Zhang S-Q, Liu M-Y, Wan B, Zheng H-M. Contralateral body half Hypalgesia in a patient with lateral medullary infarction: atypical Wallenberg syndrome. *Eur Neurol.* 2008;59(3–4):211–5. <https://doi.org/10.1159/000114050>.
 56. Porta-Etessam J, Casanova I, Pajuelo B, di Capua D, del Val J, García García ME, et al. See-saw nystagmus in a patient with Wallenberg syndrome. *J Neuroophthalmol.* 2009;29(1):73–4. <https://doi.org/10.1097/WNO.0b013e3181989dc1>.
 57. Qiu W, Wu J-S, Carroll WM, Mastaglia FL, Kermod AG. Wallenberg syndrome caused by multiple sclerosis mimicking stroke. *J Clin Neurosci.* 2009;16(12):1700–2. <https://doi.org/10.1016/j.jocn.2009.04.008>.
 58. Seo WK, Kwon DY, Seo SH, Park MH, Park KW. Neuropathic pruritus following Wallenberg syndrome. *Neurology.* 2009;72(7):676. <https://doi.org/10.1212/01.wnl.0000342484.99298.52>.
 59. Yeh H-F, Seak C-J, Chiu T-F, Chang Y-C. Traumatic vertebral artery dissection and Wallenberg syndrome after a motorcycle collision. *Am J Emerg Med.* 2009;27(1):131.e1–3.
 60. Ayerbe Zabaleta I, Negrevergne M, Ucelay Vicinay I, Catherine M. Vértigo agudo pseudoperiférico y síndrome de Wallenberg. *Acta Otorrinolaringológica Esp.* 2010;61(4):312–4. <https://doi.org/10.1016/j.otorri.2009.06.001>.
 61. Zhu J, Li J, Liu Y, Han Y, Wang Y. Unilateral headache with visual Aura from a Wallenberg syndrome. *CNS Neurosci Ther.* 2012;18(7):598–600. <https://doi.org/10.1111/j.1755-5949.2012.00330.x>.
 62. Stengl KL, Buchert R, Bauknecht H, Sobesky J. A hidden giant: Wallenberg syndrome and aortal wall thickening as an atypical presentation of a giant cell arteritis. *Case Rep.* 2013;2013(mar01 1):bcr2012006994.
 63. Ueda M, Nishiyama Y, Abe A, Katayama Y. Hemorrhagic Wallenberg syndrome. *Intern Med.* 2013;52(20):2383–4. <https://doi.org/10.2169/interia.52.1141>.
 64. Wu S, Li N, Xia F, Sidlauskas K, Lin X, Qian Y, et al. Neurotrophic keratopathy due to dorsolateral medullary infarction (Wallenberg syndrome): case report and literature review. *BMC Neurol.* 2014;14(1):231. <https://doi.org/10.1186/s12883-014-0231-y>.
 65. Das P, Chopra A, Rai A, Kuppuswamy PS. Late-onset recurrent mania as a manifestation of Wallenberg syndrome: a case report and review of the literature. *Bipolar Disord.* 2015;17(6):677–82. <https://doi.org/10.1111/bdi.12318>.
 66. Ehresmann A, Van H, Merlini L, Fluss J. Wallenberg syndrome: an exceptional cause of acute Vertigo in children. *Neuropediatrics.* 2015;47(01):061–3. <https://doi.org/10.1055/s-0035-1566731>.
 67. Louis DW, Dholakia N, Raymond MJ. Wallenberg syndrome with associated motor weakness in a two-week-postpartum female. *Case Rep Neurol.* 2015;7(3):186–90. <https://doi.org/10.1159/000440712>.
 68. Ospino Quiroz JC, Monteagudo CJ. A propósito de un caso de síndrome de Wallenberg. *SEMERGEN Med Fam.* 2016;42(8):e179–80. <https://doi.org/10.1016/6/j.semerg.2015.10.012>.
 69. Kombliuh A, Twanow J-D. Teaching NeuroImages: adolescent Wallenberg syndrome with overlooked signs: Ipsipulsion and ipsilateral facial palsy. *Neurology.* 2018;91(20):e1949–50. <https://doi.org/10.1212/WNL.0000000000006513>.
 70. Oks M, Li A, Makaryus M, Pomeranz HD, Sachdeva M, Pullman J, et al. Sarcoidosis presenting as Wallenberg syndrome and panuveitis. *Respir Med Case Rep.* 2018;24:16–8. <https://doi.org/10.1016/j.mcr.2018.03.002>.
 71. Sivakumar K, Garcha M, Mehta D, Leary MC, Yacoub HA. Central hypoventilation: a rare complication of Wallenberg syndrome. *Case Rep Neurol Med.* 2018;2018:1–3.
 72. Clatterbuck RE. Ultrastructural and immunocytochemical evidence that an incompetent blood-brain barrier is related to the pathophysiology of cavernous malformations. *J Neurol Neurosurg Psychiatry.* 2001;71(2):188–92. <https://doi.org/10.1136/jnnp.71.2.188>.
 73. Dammann P, Wrede KH, Maderwald S, el Hindy N, Mueller O, Chen B, et al. The venous angioarchitecture of sporadic cerebral cavernous malformations: a susceptibility weighted imaging study at 7 T MRI. *J Neurol Neurosurg Psychiatry.* 2013;84(2):194–200. <https://doi.org/10.1136/jnnp-2012-302599>.
 74. Lasjaunias P, Burrows P, Planet C. Developmental venous anomalies (DVA): the so-called venous angioma. *Neurosurg Rev.* 1986;9(3):233–42. <https://doi.org/10.1007/BF01743138>.
 75. Pozzati E, Acciarri N, Tognetti F, Marliani F, Giangaspero F. Growth, subsequent bleeding, and De novo appearance of cerebral cavernous Angiomas. *Neurosurgery.* 1996;38(4):662–70. <https://doi.org/10.1227/00006123-199604000-00006>.
 76. Zabramski JM, Wascher TM, Spetzler RF, Johnson B, Golfinos J, Drayer BP, et al. The natural history of familial cavernous malformations: results of an ongoing study. *J Neurosurg.* 1994;80(3):422–32. <https://doi.org/10.3171/jns.1994.80.3.0422>.
 77. Salman RA-S, Hall JM, Horne MA, et al. Untreated clinical course of cerebral cavernous malformations: a prospective, population-based cohort study. *Lancet Neurol.* 2012;11(3):217–24. [https://doi.org/10.1016/S1474-4422\(12\)70004-2](https://doi.org/10.1016/S1474-4422(12)70004-2).
 78. Al-Shahi Salman R, Berg MJ, Morrison L, Awad IA. Hemorrhage from cavernous malformations of the brain: definition and reporting standards. *Stroke.* 2008;39(12):3222–30. <https://doi.org/10.1161/STROKEAHA.108.515544>.
 79. Akers A, Al-Shahi Salman R, A. Awad I, et al. synopsis of guidelines for the clinical Management of Cerebral Cavernous Malformations: consensus recommendations based on systematic literature review by the Angioma Alliance scientific advisory board clinical experts panel. *Neurosurgery.* 2017;80(5):665–80. <https://doi.org/10.1093/neuros/nyx091>.
 80. Horne MA, Flemming KD, Su L-C, Stapf C, Jeon JP, Li D, et al. Clinical course of untreated cerebral cavernous malformations: a meta-analysis of individual patient data. *Lancet Neurol.* 2016;15(2):166–73. [https://doi.org/10.1016/S1474-4422\(15\)00303-8](https://doi.org/10.1016/S1474-4422(15)00303-8).
 81. Li D, Hao S-Y, Jia G-J, Wu Z, Zhang L-W, Zhang J-T. Hemorrhage risks and functional outcomes of untreated brainstem cavernous malformations. *J Neurosurg.* 2014;121(1):32–41. <https://doi.org/10.3171/2014.3.JNS132537>.
 82. Kyoshima K, Kobayashi S, Gibo H, Kuroyanagi T. A study of safe entry zones via the floor of the fourth ventricle for brain-stem lesions: report of three cases. *J Neurosurg.* 1993;78(6):987–93. <https://doi.org/10.3171/jns.1993.78.6.0987>.
 83. Cavalcanti DD, Preul MC, Kalani MYS, Spetzler RF. Microsurgical anatomy of safe entry zones to the brainstem. *J Neurosurg.* 2016;124(5):1359–76. <https://doi.org/10.3171/2015.4.JNS141945>.
 84. Zaidi HA, Mooney MA, Levitt MR, Dru AB, Abila AA, Spetzler RF. Impact of Timing of Intervention Among 397 Consecutively Treated Brainstem Cavernous Malformations. *Neurosurgery [Internet]* 2017 [cited 2020 May 28]; Available from: <https://academic.oup.com/neurosurgery/article/2981949/Impact>
 85. Garcia RM, Ivan ME, Lawton MT. Brainstem cavernous malformations. *Neurosurgery.* 2015;76(3):265–78. <https://doi.org/10.1227/NEU.0000000000000602>.
 86. Chandra PS, Manjari T, Chandramouli BA, Jayakumar PN, Shankar SK. Cavernous-venous malformation of brain stem—report of a case and review of literature. *Surg Neurol.* 1999;52(3):280–5. [https://doi.org/10.1016/S0090-3019\(99\)00079-8](https://doi.org/10.1016/S0090-3019(99)00079-8).
 87. Yamada S, Liwnicz BH, Thompson JR, Colohan AR, Iacono RP, Tran JT. Pericapillary arteriovenous malformations angiographically manifested as cerebral venous malformations. *Neurol Res.* 2001;23(5):513–21. <https://doi.org/10.1179/016164101101198767>.
 88. Rigamonti D, Spetzler RF. The association of venous and cavernous malformations. Report of four cases and discussion of the pathophysiological, diagnostic, and therapeutic implications. *Acta Neurochir* 1988;92(1–4):100–105, DOI: <https://doi.org/10.1007/BF01401979>.
 89. Campeau NG, Lane JI. De novo development of a lesion with the appearance of a cavernous malformation adjacent to an existing developmental venous anomaly. *AJNR Am J Neuroradiol.* 2005;26(1):156–9.
 90. Wurm G, Schnizer M, Fellner FA. Cerebral Cavernous Malformations Associated with Venous Anomalies: Surgical Considerations. *Oper Neurosurg.* 2005;57(suppl_1):42–58.

91. Georgieva VB, Krastev ED. Surgical treatment of brainstem cavernous malformation with concomitant developmental venous anomaly. *Asian J Neurosurg.* 2019;14(2):557–60. https://doi.org/10.4103/ajns.AJNS_246_18.
92. Masuoka J, Matsushima T, Hikita T, Inoue E. Cerebellar swelling after sacrifice of the superior petrosal vein during microvascular decompression for trigeminal neuralgia. *J Clin Neurosci Off J Neurosurg Soc Australas.* 2009;16(10):1342–4.
93. Piatt JH, Kellogg JX. A hazard of combining the infratentorial supracerebellar and the cerebellomedullary fissure approaches: cerebellar venous insufficiency. *Pediatr Neurosurg.* 2000;33(5):243–8. <https://doi.org/10.1159/000055962>.
94. Ferroli P, Acerbi F, Tringali G, Albanese E, Broggi M, Franzini A, et al. Venous sacrifice in neurosurgery: new insights from venous indocyanine green videoangiography: clinical article. *J Neurosurg.* 2011;115(1):18–23. <https://doi.org/10.3171/2011.3.JNS10620>.
95. Kumral E, Bayülkem G, Eyyapan D. Clinical spectrum of pontine infarction. *J Neurol.* 2002;249(12):1659–70. <https://doi.org/10.1007/s00415-002-0879-x>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

