Overview of aldosterone-related genetic syndromes and recent advances

Maria-Christina Zennaro, Fabio Fernandes-Rosa, Sheerazed Boulkroun

► To cite this version:

Maria-Christina Zennaro, Fabio Fernandes-Rosa, Sheerazed Boulkroun. Overview of aldosterone-related genetic syndromes and recent advances. *Current Opinion in Endocrinology, Diabetes and Obesity*, 2018, 25 (3), pp.147-154. 10.1097/MED.0000000000000409 . inserm-03554660

HAL Id: inserm-03554660

<https://www.hal.inserm.fr/inserm-03554660>

Submitted on 3 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Overview of Aldosterone-related Genetic Syndromes and Recent Advances

Maria-Christina Zennaro^{1,2,3}, Fabio Fernandes-Rosa^{1,2,3}, Sheerazed Boulkroun^{1,2}

Affiliations:

¹INSERM, UMRS_970, Paris Cardiovascular Research Center, Paris, France

²Université Paris Descartes, Sorbonne Paris Cité, Paris, France

³Assistance Publique-Hôpitaux de Paris, Hôpital Européen Georges Pompidou, Service de Génétique, Paris, France

Address correspondence to:

Maria-Christina Zennaro, MD, PhD

INSERM, U970

Paris Cardiovascular Research Center – PARCC

56, rue Leblanc,

75015 Paris – France

Tel : +33 (0)1 53 98 80 42

Fax : + 33 (0)1 53 98 79 52

e-mail: maria-christina.zennaro@inserm.fr

Disclosures

The authors have nothing to disclose.

Abstract

Purpose of review: Primary aldosteronism is the most common form of secondary hypertension. Early diagnosis and treatment are key to cure of hypertension and prevention of cardiovascular complications. Recent genetic discoveries have improved our understanding on the pathophysiology of aldosterone production and triggered the development of new diagnostic procedures and targeted treatments for PA.

Recent findings: Different inherited genetic abnormalities distinguish specific forms of familial hyperaldosteronism. Somatic mutations are found not only in aldosterone producing adenoma leading to primary aldosteronism, but also in aldosterone producing cell clusters of normal and micronodules from image-negative adrenal glands. Genetic knowledge has allowed the discovery of surrogate biomarkers and specific pharmacological inhibitors. Aging appears to be associated with dysregulated and relatively autonomous aldosterone production.

Summary: New biochemical markers and pharmacological approaches may allow pre-operative identification of somatic mutation carriers and use of targeted treatments.

Keywords: hypertension, adrenal, primary aldosteronism; ion channels; calcium

Introduction

Aldosterone-related genetic syndromes encompass diseases related to abnormal aldosterone production or abnormal aldosterone effects in target organs. Given the important role of aldosterone in regulating sodium and potassium balance and blood pressure in the kidney, their main clinical manifestations are arterial hypertension or salt wasting and dehydration. Different diseases affect aldosterone production in the adrenal cortex. There exist several Mendelian forms of hyperaldosteronism, termed familial hyperaldosteronism (FH) type I to IV, which share most of the phenotype but show some peculiarities. They are due to gain of function mutations in ion channels and pumps, leading to increased and autonomous aldosterone production. Somatic mutations in the same genes leading to FH have been identified in aldosterone producing adenoma (APA) leading to primary aldosteronism (PA). Increased aldosterone production can also be due to monogenic forms of adrenal insufficiency affecting cortisol production, while loss-of-function mutations in *CYP11B2*, coding for aldosterone synthase, lead to aldosterone deficiency, also called corticosterone methyl oxidase deficiency; those syndromes will be treated elsewhere in this section. On the other hand, different monogenic disorders affect aldosterone action in target tissues. They include a rare disease of aldosterone resistance in peripheral tissues named pseudohypoaldosteronism type 1 as well as excessive mineralocorticoid action due to different mechanisms either directly or indirectly related to aldosterone. In this review we will focus on primary aldosteronism as genetic disease strictu sensu related to aldosterone biosynthesis in the adrenal gland and then describe some emerging concepts related to pathophysiological aldosterone production.

(Figure 1)

The genetics of primary aldosteronism

25 PA is the most frequent form of secondary arterial hypertension. It is due to aldosterone
26 production from the adrenal cortex relatively autonomous from the renin-angiotensin system,
27 due in most cases to APA or bilateral adrenal hyperplasia. Its prevalence is estimated to be
28 around 4% of all hypertensive patients seen in primary care (1), data recently confirmed in a
29 large prospective study (2), and 10% in patients referred to tertiary care centers (1). Starting
30 from 2011, important progress has been made in our understanding of the genetic mechanisms
31 involved in PA. Currently, four Mendelian forms of PA, transmitted as autosomal dominant
32 traits, have been identified (these are reviewed in detail in (3, 4)). FH-I, also known as
33 glucocorticoid remediable aldosteronism (5), is due to the formation of a chimeric gene
34 between the highly homologous genes *CYP11B1*, coding for 11 β -hydroxylase, and *CYP11B2*,
35 coding for aldosterone synthase (6), leading to ACTH-dependent aldosterone production.
36 Mutations in *KCNJ5* have been identified in patients with FH-III, a severe form of familial
37 hyperaldosteronism associated to massive bilateral hyperplasia (7), but in some cases
38 presenting as a mild form of the disease without adrenal abnormalities (3). Germline
39 heterozygous mutations in *CACNA1D* (encoding the L-type calcium channel Cav1.3 subunit
40 α -1D) have been identified in patients with PA, seizures and neurological abnormalities, a
41 rare disease called PASNA (8). Finally, mutations in *CACNA1H* (encoding the pore-forming
42 α 1 subunit of the T-type voltage-dependent calcium channel Cav3.2) have recently been
43 reported as a cause of a new form of inherited PA, called FH-IV, associated in some cases
44 with developmental disorder (9, 10). In addition, germline mutations in the same gene have
45 been associated with mild forms of PA in patients diagnosed as FH-II (10). FH-II is
46 indistinguishable from apparently non-familial PA and only diagnosed on the basis of two or
47 more affected individuals in the same family (11). Despite linkage to 7p22 in one family (12),
48 no causative locus has been identified so far. Given that germline *KCNJ5* and *CACNA1H*
49 mutations have been identified in patients diagnosed as FH-II, and that the clinical phenotype

of FH-II is heterogeneous, being associated to APA as well as bilateral adrenal hyperplasia, it is reasonable to suspect that the disease may encompass different genetic syndromes.

In addition to germline variants, somatic mutations in *KCNJ5* (7) and *CACNA1D* (8, 13), but also ATPases (*ATP1A1* and *ATP2B3* (13, 14) have been identified in APA. Mutations in all these genes affect aldosterone production by altering pathways involved in maintaining intracellular ionic homeostasis and cell membrane potential, ultimately leading to increased calcium signalling. In particular, mutations in *KCNJ5*, coding for the potassium channel GIRK4, affect the selectivity filter of the channel and lead to increased sodium permeability and sodium influx, cell membrane depolarization and opening of voltage-gated calcium channels, followed by increased intracellular calcium concentrations and activation of calcium signalling, the main trigger for aldosterone production (15). Similarly, mutations in *ATP1A1*, encoding the $\alpha 1$ subunit of the Na^+ , K^+ -ATPase lead to loss of pump activity and inward sodium or proton leak, which were suggested to depolarize the cell membrane and increase aldosterone production through activation of calcium signaling (13, 14). More recently, experiments performed in adrenocortical NCI-H295R cells failed to demonstrate modifications of intracellular calcium concentrations in the presence of mutant Na^+ , K^+ -ATPase $\alpha 1$ subunits and rather suggest a mechanism involving intracellular acidification (16).

The pathogenic mechanisms involved in stimulating aldosterone production in the presence of *ATP1A1* mutations may therefore result from alterations in multiple pathways, possibly in relation with the multiple roles of the Na^+ , K^+ -ATPase as a pump but also a signal transducing platform (17). Mutations in *CACNA1D* and *CACNA1H* directly increase intracellular calcium concentrations, by modifying voltage-sensitivity and other functional properties of these voltage-gated calcium channels (9, 10). Similarly, *ATP2B3* mutations affecting the plasma membrane calcium-transporting ATPase 3 (PMCA3) lead to increased intracellular calcium concentrations due to loss of the physiological pump function leading to reduced calcium

export (18). The mutations are also associated with severely depolarized membrane potential of primary adenoma cells (14), which is explained by a pathological sodium leak associated with or conducted by the mutant PMCA3 (18). This may be a consequence of the location of the majority of PMCA3 mutations, which lie next to a “PEGL” motif in transmembrane helix 4, a region involved in calcium binding and ion gating. Disturbed intracellular ionic homeostasis is suggested to cause plasma membrane depolarization, leading to opening of voltage-gated Ca^{2+} channels and increased Ca^{2+} influx driving increased *CYP11B2* expression and aldosterone biosynthesis. Finally, somatic mutations in *CTNNB1*, encoding β -catenin have been described in 2-5% of APA (19, 20) and mutations in *PRKACA* (coding for the cAMP-dependent protein kinase catalytic subunit alpha) in rare cases (21). Mutations in those genes have previously been involved in cortisol producing adenoma and adrenocortical cancer; the mechanisms whereby the same mutation leads to different hormonal phenotypes remain yet to be established ((15), see below).

Clinical implications of mutation detection in PA

Mutation detection in familial forms of PA is routinely performed in some genetic laboratories and is useful for disease screening in affected families, early diagnosis and targeted treatment. In patients with APA, the challenge is represented by the detection of somatic mutations which are absent in DNA extracted from peripheral blood cells. Correlations of any biological feature associated with a particular mutation would allow to identify mutation carriers who could be suitable candidates for adrenal vein sampling and eventually targeted treatments (22, 23). The most frequent somatic event in APA are *KCNJ5* mutations whose frequency ranges from 13-77%, with higher frequency in East Asian populations ((15) and references therein). *CACNA1D* mutations are found in up to 10% of cases, followed by *ATP1A1* and *ATP2B3* mutations. A meta-analysis of 13 studies

investigating a total of 1636 patients confirmed previous results showing that *KCNJ5* mutations are more frequent in women and associated with younger age, larger tumor size and higher plasma aldosterone levels (24). Genotype-specific steroid profiles have been recently associated with APA (25). By profiling 15 steroids in adrenal venous and peripheral plasma samples from 79 patients by liquid chromatography-tandem mass spectrometry, Williams et al identified a 7-steroid fingerprint (including aldosterone, 18-oxocortisol, 18-hydroxycortisol, 21-deoxycortisol, corticosterone, 11-deoxycorticosterone, and cortisol) in peripheral vein plasma able to correctly classify 92% of the APA according to genotype (25). In addition, specific steroid features were identified, in particular the presence of significantly higher hybrid steroids 18-hydroxycortisol and 18-oxocortisol in patients with APA due to *KCNJ5* mutations. The usefulness of this type of screening is related to the possibility of selecting patients for adrenal vein sampling, but also for developing targeted treatments for specific mutation carriers. Initial studies of mutant *KCNJ5* channels showed that their pharmacology was different from that of wild type channels (22). In particular, *KCNJ5* mutants were blocked by calcium-channel blockers, such as verapamil, and Na-channel blockers, such as amiloride. At high therapeutic doses (3 x 120 mg/d) plasma concentrations of verapamil could be sufficient to block the most sensitive *KCNJ5* mutant p.Leu168Arg (22). More recently, Scholl et al applied a high-throughput screen based on the rescue of cell lethality induced by *KCNJ5* mutations. This strategy identified a series of macrolide antibiotics, including roxithromycin, that potently inhibit the mutant channels (IC₅₀ of 0.22 μM for the *KCNJ5* p.Gly151Arg mutant and 0.69 on the p.Leu168Arg) as well as *CYP11B2* expression and aldosterone production in adrenocortical HAC15 cells (23). The authors suggest that macrolides could be useful in screening tests aimed at identifying carriers of *KCNJ5* mutations, which together with peculiar imaging characteristics of these tumors (larger size, lower pre-contrast Hounsfield units on computed tomography compared with tumors carrying

mutations in other genes (19)) could guide subtype diagnosis for surgery. Those compounds could also be used as targeted treatments in patients who are not candidates for surgery, given their long pharmacological history and safety. In a proof-of-concept Study, Caroccia et al tested the effects of clarithromycin on aldosterone biosynthesis in primary cell lines obtained from APA tissues with or without the two most common *KCNJ5* mutations, p.Gly151Arg and p.Leu168Arg (26). Remarkably, clarithromycin inhibited *CYP11B2* gene expression and aldosterone production in a dose-dependent manner. Further studies are warranted to show whether these effects can be translated to patients and whether macrolides or other compounds can be used to identify and treat patients with PA due to APA with *KCNJ5* mutations or patients with FH-III otherwise requiring bilateral adrenalectomy to control blood pressure (27).

How do APA develop?

While the role of somatic mutations in regulating aldosterone biosynthesis has been clearly established, the origin of APA is still a matter of debate. Different studies have been published in the last few years leading to the emergence of two hypotheses for APA development, the “two hit model” and the “APCC model”.

The two-hit model supports the idea of the occurrence of two consecutive events leading to APA, a first one driving abnormal cell proliferation followed by a second one driving specific hormonal secretion. Adrenal cortex remodelling, reduced vascularization and ZG hyperplasia are major features of APA development (28, 29). In many cases, the resected adrenal gland harbours not only one adenoma but other micro- or macronodules, some of them expressing aldosterone synthase (28, 30, 31). Investigation of genetic abnormalities in multinodular adrenals revealed the presence of different somatic APA driver mutations in different nodules from the same adrenal (31, 32), strongly suggesting that mutations are a second event

occurring in a previously remodelled adrenal cortex. In many cases, APA have been described to be heterogeneous with aldosterone synthase expressed only in some parts of the tumour (33, 34); somatic mutations appear to be present only in the aldosterone synthase positive region of the APA (33). The recent description of a patient with PA harbouring a germline heterozygous *APC* gene mutation associated, in the resected multinodular adrenal, to biallelic *APC* inactivation due to a loss of heterozygosity in all nodules, but a somatic *KCNJ5* mutation only in the aldosterone synthase expressing nodule further supports the two-hit model for APA development (35). *APC* is part of the WNT/ β -catenin pathway which plays a major role in adrenal gland development and tumorigenesis (36). In APA, somatic mutations in *CTNNB1*, encoding β -catenin, are present in a subset of APA (19, 20) and the WNT/ β -catenin pathway is activated in two thirds of cases, independently of the mutation status (37, 38). This pathway may therefore constitute the first event leading to abnormal cell proliferation and adenoma formation. Other signalling pathways and transcriptional cascades may also be involved and create a propitious environment for the occurrence of mutations in one of the APA driver genes.

On the other hand, the APCC model supports the idea that occurrence, in ZG cells, of a somatic mutation in one of the APA driver genes leads to the formation of aldosterone-producing cell clusters (APCC), followed, through the development of APCC-to-APA translational lesions (pAATL), to APA formation (39). Next generation sequencing performed on formalin-fixed, paraffin-embedded tissue DNA identified somatic mutations in some of the known APA driver genes, i.e. *CACNA1D* and *ATP1A1* in 8 of 23 APCC from normal adrenals from kidney donors, most frequently affecting *CACNA1D*. Interestingly, no mutation in the *KCNJ5* gene, the most frequent alteration observed in APA, has been reported in APCC (39, 40). The recent description of pAATL consisting of a subcapsular APCC-like structure (APCC-like) and an inner-APA-like structure (mAPA-like) without well-defined histological

border has suggested a possible APCC to APA transition (41). These specific lesions were first described in adrenals from two patients with unilateral multiple adrenocortical micronodules. To determine whether they could constitute intermediate structures between APCC and APA, search for mutations in APA driver genes was performed in both APCC and mAPA structures. In the first case a *KCNJ5* mutation was identified in frozen tissue from a suspected aldosterone producing nodule. A different *KCNJ5* mutation resulting in the same amino acid change was found in the mAPA-like part of a first pAATL, but not in its neighbouring APCC-like structure, indicating that the aldosterone producing nodule and the mAPA had different origins. The fact that only the mAPA portion had the mutation suggested that the mutation may have led to a transdifferentiation of the cells into a mAPA like phenotype. In a second pAATL, an *ATP1A1* mutation was found in the entire pAATL suggesting a clonal origin of the entire structure. Histologically, the adrenal from the second case revealed no major abnormalities except for the presence of a pAATL (42). In a recent report of 4 additional cases, the same authors identified mutations in *KCNJ5* preferentially found in nodules larger than 3mm (41). Taken together, these results suggest that certain APA, particularly those harbouring *CACNA1D*, *ATP1A1* and *ATP2B3* mutations, could derive from APCC, although the precise mechanism remains to be identified. The progression of APCC to APA could be the result of a single-somatic mutation event or alternatively require a second-hit mutation within the APCC to promote cell proliferation. However, it has to be taken into consideration that not all mutations identified in APCCs or pAATL affect amino acids recurrently mutated in APA; those mutations therefore require functional validation for their implication in promoting aldosterone biosynthesis and eventually cell proliferation.

The new physiology of aldosterone production - autonomous aldosteronism with age?

199 In human adrenals, not all cells express *CYP11B2*, possibly reflecting high salt intake in the
200 general population (34). Two patterns of *CYP11B2* expression have been described in the
201 “normal” adrenal gland: a discontinuous expression in scattered cells underneath the capsule
202 and expression in cell clusters named APCC (28, 29). They are composed of subcapsular zona
203 glomerulosa-like cells and inner large zona fasciculata-like cells, expressing *CYP11B2* but not
204 *CYP11B1* (28, 29). Despite the inner portion resembling to zona fasciculata, their
205 transcriptome very much resembles to the transcriptome of zona glomerulosa cells, but with
206 an enhanced capacity to produce aldosterone (39, 41). It has been postulated that APCC are
207 structures of constitutive aldosterone biosynthesis, independent of renin and angiotensin II
208 stimulation (39). This is supported by the fact that APCC in normal adrenal glands harbour
209 somatic mutations in a subset of APA driver genes (39) and that APCC are present in the
210 normal adrenal as well as in peritumoral tissue of adrenals with APA (28, 29). Of note,
211 increased expression of the melanocortin 2 receptor (MC2R) has been reported in APCC,
212 suggesting that adrenocorticotrophic hormone (ACTH) could play a role in the regulation of
213 aldosterone biosynthesis in APCC (39).

214 To assess the evolution of the adrenal cortex structure and function with time, Nishimoto et al
215 have investigated the presence of APCC in adrenal tissues from 33 autopsied patients aged
216 between 0 and 50 years (43). Adrenals from subjects under 11 years had layered zona
217 glomerulosa and zona fasciculata without apparent APCC. The number and size of APCC
218 increased with age. These results suggest either that a somatic mutation occurring in the zona
219 glomerulosa blocks transdifferentiation to zona fasciculata and promotes APCC formation, or
220 that APCC may develop due to aging and environmental factors, with somatic mutations
221 being induced by unknown factors.

222 Based on the assumption that APCC are structures of constitutive aldosterone biosynthesis,
223 the relation of autonomous aldosterone production, APCC and age has been investigated by

evaluating the relationship between age and adrenal aldosterone synthase expression in 127 adrenal glands from deceased kidney donors aged 9 months to 68 years (44). This study confirmed that continuous expression of CYP11B2 was more frequent in young adrenal glands, while old adrenals very often showed discontinuous or absent zona glomerulosa CYP11B2 expression. Total and normal zona glomerulosa CYP11B2-expressing areas in the adrenal cortex were negatively correlated with age, while the area of CYP11B2 expression in APCC increased. The effect of age on aldosterone production was in parallel investigated in the HyperPATH study population, which consists of normotensive or mildly hypertensive subjects who underwent a high salt and low salt diet (45). On a high salt diet, subjects without PA showed a progressive decline of renin activity with age, while serum and 24-hour urinary aldosterone levels did not change; as a consequence, the aldosterone-to renin ratio was positively associated with older age. Under sodium restriction, the physiological stimulation of both plasma renin activity and aldosterone was blunted with increasing age. Altogether, these findings suggest a progressive pattern of abnormal aldosterone physiology with aging. The presence of APCC, which are supposedly renin-independent aldosterone producing structures, manifests with renin-independent aldosterone secretion and a decreased capacity of the adrenal gland to respond to physiological stimuli with age. The authors suggest that aging may be associated with a subclinical form of aldosterone excess. The fact that total aldosterone-expressing area in the adrenal cortex rather diminishes and aldosterone levels do not significantly change with age in the HyperPATH population rather suggests an inappropriate aldosterone production to salt status or an appropriate relatively autonomous aldosterone production.

To gain insight into how APCC evolve with age in subjects without hypertension, Omata et al investigated 107 autaptic adrenal glands using CYP11B2 immunohistochemistry and NGS and determined the APCC score and the associated somatic mutations in non-hypertensive

patients from Japanese origin (46). The number of APCCs, which was in average 0.6 per gland, increased with age, without difference in sex distribution. 21 of 61 APCCs (34%) showed somatic mutations in one of the APA driver genes except *KCNJ5*, with 16 affecting *CACNA1D* (76%), which therefore appears to be the most frequently mutated gene in APCCs in healthy individuals from different origins (39) as well as micronodules from image-negative adrenals removed on the basis of positive lateralization of aldosterone production at AVS (40). As only one third of the genetic variants identified have previously been described in APA and functionally characterized, the consequences of the remaining somatic mutations detected in APCC warrants further examination. Remarkably, in four adrenal glands different somatic mutations were identified in individual APCCs, supporting a different clonal origin of each structure, similar to what observed in multiple nodules in multinodular adrenals with APA (32). The results reported here suggest that APCC accumulation may be an age-dependent phenomenon in non-hypertensive subjects, possibly explaining some cases of inappropriate aldosterone secretion not associated with hypertension. As their number was lower in normotensive compared to CT-negative PA patients (40), the authors suggest that APCC may possibly contribute to hypertension not related to PA. Whether APCC may explain some cases of low-renin hypertension requires further investigation.

Conclusion

After a large number of studies identifying inherited and somatic mutations in familial hyperaldosteronism and APA, the last years have been devoted to the understanding of the functional consequences of these mutations, their possible clinical relevance and screening, and the search of pharmacological approaches specifically targeting mutated proteins. Exciting new knowledge has also been gathered on normal adrenal cortex structure and function in relation to aldosterone production and the genetics of APCC. It can be anticipated that this knowledge will be applied to patients with inherited or sporadic PA in the next few years. In addition, new genes should be identified in patients with as yet unknown genetic defects, which represent half of the patients harboring an APA and a large proportion of patients with familial hyperaldosteronism. The causes underlying the development of bilateral adrenal hyperplasia and idiopathic hyperaldosteronism remain still to be identified.

Key points

- Four Mendelian forms of hyperaldosteronism, inherited as autosomal dominant traits, have been described
- Inherited mutations of different genes coding for ion channels and ATPases have been identified in familial hyperaldosteronism; similar somatic mutations are found in APA
- Pre-operative detection of somatic mutations may allow improved diagnosis and treatment of PA due to APA
- APCC are autonomous aldosterone-producing structures of the normal adrenal harboring somatic mutations in APA driver genes
- The aging adrenal may progressively show features of autonomous aldosterone production

Acknowledgments

1. Acknowledgements. None.

2. Financial support and sponsorship. This work was funded through institutional support from INSERM and by the Agence Nationale pour la Recherche (ANR Blanc 2011, No.: 11-BSV1005 03, ANR-13-ISV1-0006-01), the Fondation pour la Recherche Médicale (DEQ20140329556), the Programme Hospitalier de Recherche Clinique (PHRC grant AOM06179), and by grants from INSERM and Ministère Délégué à la Recherche et des Nouvelles Technologies. The laboratory of Dr. Maria-Christina Zennaro is also partner of the H2020 project ENSAT-HT grant n° 633983.

3. Conflicts of interest. None.

References

1. Hannemann A, Wallaschofski H. Prevalence of primary aldosteronism in patient's cohorts and in population-based studies--a review of the current literature. *Horm Metab Res.* 2012 Mar;44(3):157-62. PubMed PMID: 22135219. Epub 2011/12/03. eng.
2. Monticone S, Burrello J, Tizzani D, Bertello C, Viola A, Buffolo F, et al. Prevalence and Clinical Manifestations of Primary Aldosteronism Encountered in Primary Care Practice. *J Am Coll Cardiol.* 2017 Apr 11;69(14):1811-20. PubMed PMID: 28385310.
3. Korah HE, Scholl UI. An Update on Familial Hyperaldosteronism. *Horm Metab Res.* 2015 Dec;47(13):941-6. PubMed PMID: 26445452.
4. Fernandes-Rosa FL, Boulkroun S, Zennaro MC. Somatic and inherited mutations in primary aldosteronism. *J Mol Endocrinol.* 2017 Jul;59(1):R47-R63. PubMed PMID: 28400483.
5. Sutherland DJ, Ruse JL, Laidlaw JC. Hypertension, increased aldosterone secretion and low plasma renin activity relieved by dexamethasone. *Can Med Assoc J.* 1966 Nov 26;95(22):1109-19. PubMed PMID: 4288576. Pubmed Central PMCID: 1935810. Epub 1966/11/26. eng.
6. Lifton RP, Dluhy RG, Powers M, Rich GM, Cook S, Ulick S, et al. A chimaeric 11beta-hydroxylase aldosterone synthase gene causes glucocorticoid-remediable aldosteronism and human hypertension. *Nature.* 1992;355:262-5.
7. Choi M, Scholl UI, Yue P, Bjorklund P, Zhao B, Nelson-Williams C, et al. K⁺ channel mutations in adrenal aldosterone-producing adenomas and hereditary hypertension. *Science.* 2011 Feb 11;331(6018):768-72. PubMed PMID: 21311022. Epub 2011/02/12. eng.

- 329 8. Scholl UI, Goh G, Stolting G, de Oliveira RC, Choi M, Overton JD, et al. Somatic and
330 germline CACNA1D calcium channel mutations in aldosterone-producing adenomas and
331 primary aldosteronism. *Nat Genet.* 2013 Sep;45(9):1050-4. PubMed PMID: 23913001.
332 Pubmed Central PMCID: 3876926.
- 333 9. Scholl UI, Stolting G, Nelson-Williams C, Vichot AA, Choi M, Loring E, et al.
334 Recurrent gain of function mutation in calcium channel CACNA1H causes early-onset
335 hypertension with primary aldosteronism. *eLife.* 2015;4:e06315. PubMed PMID: 25907736.
336 Pubmed Central PMCID: 4408447.
- 337 10. *Daniil G, Fernandes-Rosa FL, Chemin J, Blesneac I, Beltrand J, Polak M, et al.
338 CACNA1H Mutations Are Associated With Different Forms of Primary Aldosteronism.
339 *EBioMedicine.* 2016 Nov;13:225-36. PubMed PMID: 27729216. This study describes the
340 presence of germline CACNA1H mutations in patients with early onset PA, familial PA
341 diagnosed as FH-II and PA due to APA, suggesting that CACNA1H might be a susceptibility
342 gene predisposing to PA with different phenotypic presentations.
- 343 11. Stowasser M, Gordon RD. Familial hyperaldosteronism. *J Steroid Biochem Mol Biol.*
344 2001;78(3):215-29.
- 345 12. Lafferty AR, Torpy DJ, Stowasser M, Taymans SE, Lin JP, Huggard P, et al. A novel
346 genetic locus for low renin hypertension: familial hyperaldosteronism type II maps to
347 chromosome 7 (7p22). *J Med Genet.* 2000;37(11):831-5.
- 348 13. Azizan EA, Poulsen H, Tuluc P, Zhou J, Clausen MV, Lieb A, et al. Somatic
349 mutations in ATP1A1 and CACNA1D underlie a common subtype of adrenal hypertension.
350 *Nat Genet.* 2013 Sep;45(9):1055-60. PubMed PMID: 23913004.
- 351 14. Beuschlein F, Boulkroun S, Osswald A, Wieland T, Nielsen HN, Lichtenauer UD, et
352 al. Somatic mutations in ATP1A1 and ATP2B3 lead to aldosterone-producing adenomas and

353 secondary hypertension. Nat Genet. 2013 Apr;45(4):440-4. PubMed PMID: 23416519. Epub
354 2013/02/19. eng.

355 15. Zennaro MC, Boulkroun S, Fernandes-Rosa F. Genetic Causes of Functional
356 Adrenocortical Adenomas. Endocr Rev. 2017 Dec 1;38(6):516-37. PubMed PMID:
357 28973103.

358 16. Stindl J, Tauber P, Sterner C, Tegtmeier I, Warth R, Bandulik S. Pathogenesis of
359 Adrenal Aldosterone-Producing Adenomas Carrying Mutations of the Na(+)/K(+)-ATPase.
360 Endocrinology. 2015 Dec;156(12):4582-91. PubMed PMID: 26418325.

361 17. Reinhard L, Tidow H, Clausen MJ, Nissen P. Na(+),K (+)-ATPase as a docking
362 station: protein-protein complexes of the Na(+),K (+)-ATPase. Cellular and molecular life
363 sciences : CMLS. 2013 Jan;70(2):205-22. PubMed PMID: 22695678.

364 **18. Tauber P, Aichinger B, Christ C, Stindl J, Rhayem Y, Beuschlein F, et al. Cellular
365 Pathophysiology of an Adrenal Adenoma-Associated Mutant of the Plasma Membrane
366 Ca(2+)-ATPase ATP2B3. Endocrinology. 2016 Jun;157(6):2489-99. PubMed PMID:
367 27035656. This is the first study addressing the mechanism of PA associated with mutations
368 in *ATP2B3*.

369 19. Scholl UI, Healy JM, Thiel A, Fonseca AL, Brown TC, Kunstman JW, et al. Novel
370 Somatic Mutations in Primary Hyperaldosteronism are related to the Clinical, Radiological
371 and Pathological Phenotype. Clin Endocrinol (Oxf). 2015 Aug 7. PubMed PMID: 26252618.

372 20. Akerstrom T, Maharjan R, Sven Willenberg H, Cupisti K, Ip J, Moser A, et al.
373 Activating mutations in CTNNB1 in aldosterone producing adenomas. Scientific reports.
374 2016;6:19546. PubMed PMID: 26815163. Pubmed Central PMCID: 4728393.

375 21. Rhayem Y, Perez-Rivas LG, Dietz A, Bathon K, Gebhard C, Riester A, et al.
376 PRKACA Somatic Mutations Are Rare Findings in Aldosterone-Producing Adenomas. J Clin
377 Endocrinol Metab. 2016 Aug;101(8):3010-7. PubMed PMID: 27270477.

378 22. Tauber P, Penton D, Stindl J, Humberg E, Tegtmeier I, Sterner C, et al. Pharmacology
379 and pathophysiology of mutated KCNJ5 found in adrenal aldosterone-producing adenomas.
380 Endocrinology. 2014 Apr;155(4):1353-62. PubMed PMID: 24506072.

381 **23. Scholl UI, Abriola L, Zhang C, Reimer EN, Plummer M, Kazmierczak BI, et al.
382 Macrolides selectively inhibit mutant KCNJ5 potassium channels that cause aldosterone-
383 producing adenoma. J Clin Invest. 2017 Jun 12. PubMed PMID: 28604387. This study
384 identifies a series of macrolide antibiotics, including roxithromycin, that potently inhibit
385 mutated KCNJ5 channel. This discovery could have implication for diagnosis and treatment
386 of patients with APA carriers of KCNJ5 mutations.

387 24. Lenzini L, Rossitto G, Maiolino G, Letizia C, Funder JW, Rossi GP. A Meta-Analysis
388 of Somatic KCNJ5 K(+) Channel Mutations In 1636 Patients With an Aldosterone-Producing
389 Adenoma. J Clin Endocrinol Metab. 2015 Aug;100(8):E1089-95. PubMed PMID: 26066531.

390 **25. Williams TA, Peitzsch M, Dietz AS, Dekkers T, Bidlingmaier M, Riester A, et al.
391 Genotype-Specific Steroid Profiles Associated With Aldosterone-Producing Adenomas.
392 Hypertension. 2016 Jan;67(1):139-45. PubMed PMID: 26573708. In this study, steroid
393 profiling of adrenal vein and peripheral blood of patients with APA allowed to discriminate
394 the mutation status with high sensitivity and specificity.

395 *26. Caroccia B, Prisco S, Seccia TM, Piazza M, Maiolino G, Rossi GP. Macrolides Blunt
396 Aldosterone Biosynthesis: A Proof-of-Concept Study in KCNJ5 Mutated Adenoma Cells Ex
397 Vivo. Hypertension. 2017 Dec;70(6):1238-42. PubMed PMID: 28993452. This study shows

398 that clarithromycin was effective in reducing aldosterone production and CYP11B2
399 expression in primary cell cultures from APA tissues with KCNJ5 mutations.

400 27. Geller DS, Zhang J, Wisgerhof MV, Shackleton C, Kashgarian M, Lifton RP. A novel
401 form of human mendelian hypertension featuring nonglucocorticoid-remediable
402 aldosteronism. *J Clin Endocrinol Metab.* 2008 Aug;93(8):3117-23. PubMed PMID:
403 18505761. Pubmed Central PMCID: 2515083. Epub 2008/05/29. eng.

404 28. Boulkroun S, Samson-Couterie B, Dzib JF, Lefebvre H, Louiset E, Amar L, et al.
405 Adrenal cortex remodeling and functional zona glomerulosa hyperplasia in primary
406 aldosteronism. *Hypertension.* 2010 Nov;56(5):885-92. PubMed PMID: 20937967. Epub
407 2010/10/13. eng.

408 29. Nishimoto K, Nakagawa K, Li D, Kosaka T, Oya M, Mikami S, et al. Adrenocortical
409 zonation in humans under normal and pathological conditions. *J Clin Endocrinol Metab.* 2010
410 May;95(5):2296-305. PubMed PMID: 20200334. Epub 2010/03/05. eng.

411 30. Enberg U, Volpe C, Hoog A, Wedell A, Farnebo LO, Thoren M, et al. Postoperative
412 differentiation between unilateral adrenal adenoma and bilateral adrenal hyperplasia in
413 primary aldosteronism by mRNA expression of the gene CYP11B2. *Eur J Endocrinol.* 2004
414 Jul;151(1):73-85. PubMed PMID: 15248825.

415 31. Dekkers T, ter Meer M, Lenders JW, Hermus AR, Schultze Kool L, Langenhuijsen JF,
416 et al. Adrenal nodularity and somatic mutations in primary aldosteronism: one node is the
417 culprit? *J Clin Endocrinol Metab.* 2014 Jul;99(7):E1341-51. PubMed PMID: 24758183.

418 32. Fernandes-Rosa FL, Giscos-Douriez I, Amar L, Gomez-Sanchez CE, Meatchi T,
419 Boulkroun S, et al. Different Somatic Mutations in Multinodular Adrenals With Aldosterone-
420 Producing Adenoma. *Hypertension.* 2015 Nov;66(5):1014-22. PubMed PMID: 26351028.
421 Pubmed Central PMCID: 4600038.

422 33. Nanba K, Chen AX, Omata K, Vinco M, Giordano TJ, Else T, et al. Molecular
423 Heterogeneity in Aldosterone-Producing Adenomas. *J Clin Endocrinol Metab.* 2016
424 Mar;101(3):999-1007. PubMed PMID: 26765578. Pubmed Central PMCID: 4803171.

425 34. Gomez-Sanchez CE, Gomez-Sanchez EP. Immunohistochemistry of the adrenal in
426 primary aldosteronism. *Curr Opin Endocrinol Diabetes Obes.* 2016 Jun;23(3):242-8. PubMed
427 PMID: 27119751. Pubmed Central PMCID: 4915062.

428 *35. Vouillarmet J, Fernandes-Rosa F, Graeppi-Dulac J, Lantelme P, Decaussin-Petrucci
429 M, Thivolet C, et al. Aldosterone-Producing Adenoma With a Somatic KCNJ5 Mutation
430 Revealing APC-Dependent Familial Adenomatous Polyposis. *J Clin Endocrinol Metab.* 2016
431 Nov;101(11):3874-8. PubMed PMID: 27648962. This study describes the presence of two
432 genetic events associated with APA formation.

433 36. El Wakil A, Lalli E. The Wnt/beta-catenin pathway in adrenocortical development and
434 cancer. *Mol Cell Endocrinol.* 2011 Jan 30;332(1-2):32-7. PubMed PMID: 21094679. Epub
435 2010/11/26. eng.

436 37. Boulkroun S, Samson-Couterie B, Golib-Dzib JF, Amar L, Plouin PF, Sibony M, et al.
437 Aldosterone-producing adenoma formation in the adrenal cortex involves expression of
438 stem/progenitor cell markers. *Endocrinology.* 2011 Dec;152(12):4753-63. PubMed PMID:
439 21971159. Epub 2011/10/06. eng.

440 38. Berthon A, Drelon C, Ragazzon B, Boulkroun S, Tissier F, Amar L, et al. WNT/beta-
441 catenin signalling is activated in aldosterone-producing adenomas and controls aldosterone
442 production. *Hum Mol Genet.* 2014 Feb 15;23(4):889-905. PubMed PMID: 24087794.

443 39. Nishimoto K, Tomlins SA, Kuick R, Cani AK, Giordano TJ, Hovelson DH, et al.
444 Aldosterone-stimulating somatic gene mutations are common in normal adrenal glands. *Proc*
445 *Natl Acad Sci U S A.* 2015 Aug 18;112(33):E4591-9. PubMed PMID: 26240369.

446 *40. Yamazaki Y, Nakamura Y, Omata K, Ise K, Tezuka Y, Ono Y, et al.
447 Histopathological Classification of Cross-Sectional Image-Negative Hyperaldosteronism. J
448 Clin Endocrinol Metab. 2017 Apr 01;102(4):1182-92. PubMed PMID: 28388725. This study
449 describes high frequency somatic mutations in micronodules of image-negative PA.

450 41. Nishimoto K, Koga M, Seki T, Oki K, Gomez-Sanchez EP, Gomez-Sanchez CE, et al.
451 Immunohistochemistry of aldosterone synthase leads the way to the pathogenesis of primary
452 aldosteronism. Mol Cell Endocrinol. 2017 Feb 5;441:124-33. PubMed PMID: 27751767.
453 Pubmed Central PMCID: 5470036.

454 *42. Nishimoto K, Seki T, Kurihara I, Yokota K, Omura M, Nishikawa T, et al. Case
455 Report: Nodule Development From Subcapsular Aldosterone-Producing Cell Clusters Causes
456 Hyperaldosteronism. J Clin Endocrinol Metab. 2016 Jan;101(1):6-9. PubMed PMID:
457 26580238. First report describing possible APCC-to-APA transitional lesions.

458 *43. Nishimoto K, Seki T, Hayashi Y, Mikami S, Al-Eyd G, Nakagawa K, et al. Human
459 Adrenocortical Remodeling Leading to Aldosterone-Producing Cell Cluster Generation.
460 International journal of endocrinology. 2016;2016:7834356. PubMed PMID: 27721827.
461 Pubmed Central PMCID: 5046023 publication of this manuscript. This is the first study
462 reporting increase in number and size of APCC with age.

463 **44. Nanba K, Vaidya A, Williams GH, Zheng I, Else T, Rainey WE. Age-Related
464 Autonomous Aldosteronism. Circulation. 2017 May 31. PubMed PMID: 28566337. This
465 study shows that aging is associated with dysregulated aldosterone synthase expression from
466 APCC, renin-independent aldosterone secretion and a decreased capacity of the adrenal gland
467 to respond to physiological stimuli with age.

468 45. Underwood PC, Chamarthi B, Williams JS, Vaidya A, Garg R, Adler GK, et al.
469 Nonmodulation as the mechanism for salt sensitivity of blood pressure in individuals with

470 hypertension and type 2 diabetes mellitus. J Clin Endocrinol Metab. 2012 Oct;97(10):3775-
471 82. PubMed PMID: 22865897. Pubmed Central PMCID: 3462947.

472 *46. Omata K, Anand SK, Hovelson DH, Liu CJ, Yamazaki Y, Nakamura Y, et al.
473 Aldosterone-Producing Cell Clusters Frequently Harbor Somatic Mutations and Accumulate
474 With Age in Normal Adrenals. Journal of the Endocrine Society. 2017 Jul 1;1(7):787-99.
475 PubMed PMID: 29264530. Pubmed Central PMCID: 5686701. This study shows increased
476 number and size of APCC with aging in non-hypertensive Japanese subjects.

477

478

Figure legends.

Figure 1. Two models for APA development. Foci are composed of zona glomerulosa cells where aldosterone production is under the control of its secretagogues, principally angiotensin II (AngII) and potassium (K⁺). On the left part of the adrenal, the “two-hit model”. Local abnormal cell proliferation occurs in some zona glomerulosa cells, due to activation of specific signaling pathway such as Wnt/β-catenin or shh, leading to nodule formation. Subsequently, recurrent somatic mutations in *KCNJ5*, *CACNA1D*, *ATP1A1* or *ATP2B3* occur, inducing autonomous aldosterone hypersecretion and APA development. On the right part of the adrenal, the “APCC model”. Somatic mutations in *CACNA1D* or *ATP1A1* lead to constitutive aldosterone production in specific structures named APCC. APCC develop into pAATL and then into APA, through abnormal cell proliferation driven by the previously described somatic mutations or by other mechanisms, which remain to be identified.